

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
United States Patent and Trademark
Office
Box PCT
Washington, D.C.20231
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 03 October 2000 (03.10.00)	
International application No. PCT/GB00/00373	Applicant's or agent's file reference PHM70495/WO
International filing date (day/month/year) 08 February 2000 (08.02.00)	Priority date (day/month/year) 10 February 1999 (10.02.99)
Applicant HENNEQUIN, Laurent, François, Andre et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

22 August 2000 (22.08.00)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Zakaria EL KHODARY Telephone No.: (41-22) 338.83.38
---	--

PATENT COOPERATION TREATY

PCT

From the INTERNATIONAL BUREAU

NOTIFICATION OF THE RECORDING
OF A CHANGE

(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

To:

BRYANT, Tracey
AstraZeneca
Global Intellectual Property
P.O. Box 272
Mersey, Alderley Park
Macclesfield, Cheshire SK10 4GR
ROYAUME-UNI

Date of mailing (day/month/year) 16 August 2000 (16.08.00)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference PHM70495/WO	
International application No. PCT/GB00/00373	International filing date (day/month/year) 08 February 2000 (08.02.00)

1. The following indications appeared on record concerning:									
<input type="checkbox"/> the applicant	<input type="checkbox"/> the inventor <input checked="" type="checkbox"/> the agent <input type="checkbox"/> the common representative								
Name and Address BRYANT, Tracey AstraZeneca Global Intellectual Property Mersey, Alderley Park Macclesfield Cheshire SK10 4TG United Kingdom	<table border="1"> <tr> <td>State of Nationality</td> <td>State of Residence</td> </tr> <tr> <td colspan="2">Telephone No. 01625 513 228</td> </tr> <tr> <td colspan="2">Facsimile No. 01625 583 358</td> </tr> <tr> <td colspan="2">Teleprinter No.</td> </tr> </table>	State of Nationality	State of Residence	Telephone No. 01625 513 228		Facsimile No. 01625 583 358		Teleprinter No.	
State of Nationality	State of Residence								
Telephone No. 01625 513 228									
Facsimile No. 01625 583 358									
Teleprinter No.									
2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:									
<input type="checkbox"/> the person	<input type="checkbox"/> the name <input checked="" type="checkbox"/> the address <input type="checkbox"/> the nationality <input type="checkbox"/> the residence								
Name and Address BRYANT, Tracey AstraZeneca Global Intellectual Property P.O. Box 272 Mersey, Alderley Park Macclesfield, Cheshire SK10 4GR United Kingdom	<table border="1"> <tr> <td>State of Nationality</td> <td>State of Residence</td> </tr> <tr> <td colspan="2">Telephone No. 01625 513 228</td> </tr> <tr> <td colspan="2">Facsimile No. 01625 583 358</td> </tr> <tr> <td colspan="2">Teleprinter No.</td> </tr> </table>	State of Nationality	State of Residence	Telephone No. 01625 513 228		Facsimile No. 01625 583 358		Teleprinter No.	
State of Nationality	State of Residence								
Telephone No. 01625 513 228									
Facsimile No. 01625 583 358									
Teleprinter No.									
3. Further observations, if necessary:									
4. A copy of this notification has been sent to:									
<input checked="" type="checkbox"/> the receiving Office	<input checked="" type="checkbox"/> the designated Offices concerned								
<input type="checkbox"/> the International Searching Authority	<input type="checkbox"/> the elected Offices concerned								
<input type="checkbox"/> the International Preliminary Examining Authority	<input type="checkbox"/> other:								

<p>The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland</p> <p>Facsimile No.: (41-22) 740.14.35</p>	<p>Authorized officer</p> <p>Mougamadou ABIDINE</p> <p>Telephone No.: (41-22) 338.83.38</p>
--	---

PATENT COOPERATION TREATY

PCT

From the INTERNATIONAL BUREAU

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

To:

BRYANT, Tracey
AstraZeneca
Global Intellectual Property
P.O. Box 272
Mereside, Alderley Park
Macclesfield, Cheshire SK10 4GR
ROYAUME-UNI

Date of mailing (day/month/year) 16 August 2000 (16.08.00)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference PHM70495/WO	
International application No. PCT/GB00/00373	International filing date (day/month/year) 08 February 2000 (08.02.00)

1. The following indications appeared on record concerning:		
<input checked="" type="checkbox"/> the applicant	<input type="checkbox"/> the inventor	<input type="checkbox"/> the agent
<input type="checkbox"/> the common representative		
Name and Address ASTRAZENECA UK LIMITED 15 Stanhope Gate London W1Y 6LN United Kingdom	State of Nationality GB	State of Residence GB
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	
2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:		
<input checked="" type="checkbox"/> the person	<input type="checkbox"/> the name	<input checked="" type="checkbox"/> the address
<input checked="" type="checkbox"/> the nationality		
<input checked="" type="checkbox"/> the residence		
Name and Address ASTRAZENECA AB S-151 85 Södertälje Sweden	State of Nationality SE	State of Residence SE
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	
3. Further observations, if necessary:		
4. A copy of this notification has been sent to:		
<input checked="" type="checkbox"/> the receiving Office	<input checked="" type="checkbox"/> the designated Offices concerned	
<input type="checkbox"/> the International Searching Authority	<input type="checkbox"/> the elected Offices concerned	
<input type="checkbox"/> the International Preliminary Examining Authority	<input type="checkbox"/> other:	

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland	Authorized officer Mougamadou ABIDINE
Facsimile No.: (41-22) 740.14.35	Telephone No.: (41-22) 338.83.38

09/913020

PCT

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

BRYANT, Tracey
AstraZeneca
Global Intellectual Property,
Patents
Alderley Park
Macclesfield
Cheshire SK10 4TG
ROYAUME-UNI

RECEIVED

12 MAY 2000

ASTRAZENeca

GLOBAL INTELLECTUAL PROPERTY

IMPORTANT NOTIFICATION

Date of mailing (day/month/year)

05 May 2000 (05.05.00)

Applicant's or agent's file reference

PHM70495/WO

International application No.

PCT/GB00/00373

International filing date (day/month/year)

08 February 2000 (08.02.00)

1. The following indications appeared on record concerning:



the applicant



the inventor



the agent



the common representative

Name and Address

ZENECA LIMITED
15 Stanhope Gate
London W1Y 6LN
United Kingdom

State of Nationality

GB

State of Residence

GB

Telephone No.

Facsimile No.

Teleprinter No.

2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:



the person



the name



the address



the nationality



the residence

Name and Address

ASTRAZENeca UK LIMITED
15 Stanhope Gate
London W1Y 6LN
United Kingdom

State of Nationality

GB

State of Residence

GB

Telephone No.

Facsimile No.

Teleprinter No.

3. Further observations, if necessary:

4. A copy of this notification has been sent to:



the receiving Office



the designated Offices concerned



the International Searching Authority



the elected Offices concerned



the International Preliminary Examining Authority



other:

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland


Authorized officer

Lazar Joseph Panakal

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference PHM70495/WO		FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/GB00/00373	International filing date (day/month/year) 08/02/2000	Priority date (day/month/year) 10/02/1999	
International Patent Classification (IPC) or national classification and IPC A61K31/505			
Applicant ASTRAZENECA UK LIMITED			
<p>1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.</p> <p>2. This REPORT consists of a total of 8 sheets, including this cover sheet.</p> <p><input checked="" type="checkbox"/> This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).</p> <p>These annexes consist of a total of 1 sheets.</p>			
<p>3. This report contains indications relating to the following items:</p> <ul style="list-style-type: none">I <input checked="" type="checkbox"/> Basis of the reportII <input type="checkbox"/> PriorityIII <input checked="" type="checkbox"/> Non-establishment of opinion with regard to novelty, inventive step and industrial applicabilityIV <input type="checkbox"/> Lack of unity of inventionV <input checked="" type="checkbox"/> Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statementVI <input type="checkbox"/> Certain documents citedVII <input checked="" type="checkbox"/> Certain defects in the international applicationVIII <input type="checkbox"/> Certain observations on the international application			
Date of submission of the demand 22/08/2000		Date of completion of this report 30.04.2001	
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465		Authorized officer Scruton-Evans, I Telephone No. +49 89 2399 8272	



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/GB00/00373

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-314 as originally filed

Claims, No.:

1-22 as originally filed

23-30 as received on 09/04/2001 with letter of 09/04/2001

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/GB00/00373

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non-obvious), or to be industrially applicable have not been examined in respect of:

☐ the entire international application.

☒ claims Nos. 22,23-30.

because:

☒ the said international application, or the said claims Nos. 22 relate to the following subject matter which does not require an international preliminary examination (*specify*):
see separate sheet

☐ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. are so unclear that no meaningful opinion could be formed (*specify*):

☐ the claims, or said claims Nos. are so inadequately supported by the description that no meaningful opinion could be formed.

☒ no international search report has been established for the said claims Nos. 23-30.

2. A meaningful international preliminary examination cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

☐ the written form has not been furnished or does not comply with the standard.

☐ the computer readable form has not been furnished or does not comply with the standard.

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	4,5,12-13,15-18
	No:	Claims	1-3,6-8,9-11,14,19,20,21,22

Inventive step (IS)	Yes:	Claims
---------------------	------	--------

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/GB00/00373

	No:	Claims	1-22
Industrial applicability (IA)	Yes:	Claims	1-21
	No:	Claims	

2. Citations and explanations
see separate sheet

VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:
see separate sheet

Re Item III.

Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

Claim 22 relates to subject-matter considered by this Authority to be covered by the provisions of Rule 67.1(iv) PCT. Consequently, no opinion will be formulated with respect to the industrial applicability of the subject-matter of this claim (Article 34(4)(a)(i) PCT).

Claims 23-30 relate to intermediate compounds and processes, which were not claimed in the application as originally filed. Thus as no search report has been drawn up for these claims, no assessment of their novelty, inventive step or industrial applicability can be made. An additional search may be necessary in the national/regional phase. It is, however, pointed out that the claims 27-30 make reference to the description, which is not allowed under Rule 6.2(a) of the PCT.

Re Item V

Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

The following documents cited in the Search Report are referred to in this communication;

D1: WO-A-9629301

D2: WO-A-9515758

D3: J. Med. Chem. (jmcmr,00222623);1999; Vol.42 (26); Pp.5369-5389
(0000), , -

D4: Bioorg. Med. Chem. (bmecep,09680896);1995; Vol.3 (12); Pp.1651-6
(0000), , -

D5: Bioorg. Med. Chem. Lett. (bmcle8,0960894x);1997; Vol.7 (21); Pp.2723-

2728 (0000), , -

D6: EP-A-0602851

D7: WO-A-9742187

D8: WO-A-9910349

D9: WO-A-9722596

Documents D3 and D8 were published after the priority date of the present application, and thus their relevance cannot be assessed until such a time as the priority document of the present application has been studied.

With regard to the requirement for novelty (Article 33(2) of the PCT), the following comments are made;

Compound claims 9-19

D2 discloses certain compounds excluded by the disclaimer in claim 9, but the 6,7-diMe compound at the top of page 17 has not been excluded, and is thus prejudicial to the novelty of claims 9-11, 14, 19, 20 and 21, and an overlap exists between the general disclosure of D2 and the present application. .

Documents D4-D6 and D7 differ in the group Zb, and those of D9 in that the group corresponding to C is not bicyclic.

Use claims 1-8

For these claims, the definition of Z is broader than that for claims 9-19, and includes NH, CH₂ or a direct bond.

Document D7 discloses compounds which fall within the scope of the claimed compounds in claim 1 when Z is a direct bond, and the use for which they are disclosed in D7 is exactly the same as that for the present application, i.e. VEGF inhibitors, useful for diseases associated with angiogenesis and/or increased vascular permeability.

Thus this document is novelty destroying for the second medical use claim 1-3, 6-8 and the process claim 20.

D9 discloses compounds differing from those of claim 1 in that they do not have a bicyclic group attached to the quinazoline.

The compounds of D1, D2, D4, D5 and D6 all fall within the scope of the formula I of claim 1, but no mention is made in any of these documents specifically of angiogenesis and/or increased vascular permeability, and thus novelty can be formally acknowledged re these documents.

With regard to the requirement for inventive step, the following comments are made;

Compound claims 9-19

The compounds of these claims are described as being VEGF inhibitors, with use in the treatment of diseases associated with angiogenesis and/or increased vascular permeability. The closest prior art is considered to be the documents D7, D6 and D2. The man skilled in the art, faced with the problem of providing further novel compounds with this activity, would have been aware that compounds of the same general formula from D2 have a receptor Tyrosine kinase activity, as do those of D6 which differ only in the group Z as NH. Furthermore, it is known that similar compounds such as those of D7 have a VEGF inhibitory activity, as do those of D9 which are not bicyclic. There thus appears to be a direct relationship between the activities known for the compounds of these claims (Receptor tyrosine kinase) and their use in diseases associated with angiogenesis and/or increased vascular permeability (Compare D9 and D2) Thus it is considered that the man skilled in the art, faced with the problem of providing further novel compounds would have prepared these compounds, expecting them to have this activity, given the prior arts discussed above. Thus the problem must have been the provision of further novel compounds with unexpected advantages re the prior art, no solution to which has been shown. thus Article 33(3) of the PCT cannot be considered to have been satisfied for these claims and the use claim 21.

Claims 1-8

These claims are essentially second medical use type claims, and, as detailed above, are not novel due to D7. Furthermore, given that the scope is within that of D2, D4, D5 and D6, and that the activity described therein appears to be directly related to that claimed in the present application (compare D9 and D2, where the same compounds have the two activities), it is considered that the man skilled in the art, faced with the problem of providing compounds with VEGF activity, would have considered the use of compounds known to have receptor tyrosine kinase activity, and thus Article 33(3) of the PCT is not considered to have been satisfied.

The process claim 20, in that it refers to the preparation of known compounds is not considered to involve an inventive step

For the assessment of the present claim 22 on the question whether it is industrially

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/GB00/00373

applicable, no unified criteria exist in the PCT Contracting States. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject-matter of claims to the use of a compound in medical treatment, but may allow, however, claims to a known compound for first use in medical treatment and the use of such a compound for the manufacture of a medicament for a new medical treatment.

Re Item VII

Certain defects in the international application

The scope of the claims should be such that only those compounds which provide a solution to the problem underlying the present application should be included.

23. A compound 4-fluoro-5-hydroxy-2-methylindole or a salt thereof.
24. A compound 4-fluoro-5-hydroxyindole or a salt thereof.
- 5 25. A compound 6-fluoro-5-hydroxy-2-methylindole or a salt thereof.
26. A compound 6-fluoro-5-hydroxyindole or a salt thereof.
- 10 27. A process for the preparation of 4-fluoro-5-hydroxy-2-methylindole according to any one of those described in Example 237.
28. A process for the preparation of 4-fluoro-5-hydroxyindole as described in Example 242.
- 15 29. A process for the preparation of 6-fluoro-5-hydroxyindole as described in Example 242.
- 20 30. A process for the preparation of 6-fluoro-5-hydroxy-2-methylindole as described in Example 250.

AMENDED SHEET

09/913020

PCT

REQUEST

The undersigned requests that the present international application be processed according to the Patent Cooperation Treaty.

For receiving Office use only

International Application No.

International Filing Date

Name of receiving Office and "PCT International Application"

Applicant's or agent's file reference
(if desired) (12 characters maximum) PHM70495/WO

Box No. I TITLE OF INVENTION

CHEMICAL COMPOUNDS

Box No. II APPLICANT

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

ZENECA Limited
15 Stanhope Gate
London. W1Y 6LN
United Kingdom

☐ This person is also inventor.

Telephone No.

(01625) 516173

Facsimile No.

(01625) 583358

Teleprinter No.

State (that is, country) of nationality:

GB

State (that is, country) of residence:

GB

This person is applicant
for the purposes of:

all designated
Statesall designated States except
the United States of Americathe United States
of America onlythe States indicated in
the Supplemental Box

Box No. III FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S)

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

ZENECA-Pharma S.A.
'Le Galien', 1 rue des Chauffours
BP 127, 95022 Cergy Cedex
France.

This person is:



applicant only



applicant and inventor

inventor only (If this check-box
is marked, do not fill in below.)

State (that is, country) of nationality:

FR

State (that is, country) of residence:

FR

This person is applicant
for the purposes of:

all designated
Statesall designated States except
the United States of Americathe United States
of America onlythe States indicated in
the Supplemental Box

☒ Further applicants and/or (further) inventors are indicated on a continuation sheet.

Box No. IV AGENT OR COMMON REPRESENTATIVE; OR ADDRESS FOR CORRESPONDENCE

The person identified below is hereby/has been appointed to act on behalf
of the applicant(s) before the competent International Authorities as:



agent



common representative

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country.)

BRYANT, Tracey
Global Intellectual Property, Patents.
ASTRAZENECA
Alderley Park,
Macclesfield, Cheshire.
SK10 4TG. GB.

Telephone No.

(01625) 513228

Facsimile No.

(01625) 583358

Teleprinter No.

☐ Address for correspondence: Mark this check-box where no agent or common representative is/has been appointed and the space above is used instead to indicate a special address to which correspondence should be sent.

Continuation of Box No. III FURTHER APPLICANTS AND/OR (FURTHER) INVENTORS

If none of the following sub-boxes is used, this sheet should not be included in the request.

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

HENNEQUIN, Laurent Francois Andre
Z.I. La Pompelle,
BP 1050, 51689 Reims Cedex 2
France

This person is:

☐ applicant only

☒ applicant and inventor

☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
FR

State (that is, country) of residence:
FR

This person is applicant for the purposes of:

☐ all designated States

☐ all designated States except the United States of America

☒ the United States of America only

☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

PLE, Patrick
Z.I. La Pompelle,
BP 1050, 51689 Reims Cedex 2
France

This person is:

☐ applicant only

☒ applicant and inventor

☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
FR

State (that is, country) of residence:
FR

This person is applicant for the purposes of:

☐ all designated States

☐ all designated States except the United States of America

☒ the United States of America only

☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

STOKES, Elaine Sophie Elizabeth
Alderley Park,
Macclesfield, Cheshire.
SK10 4TG. GB.

This person is:

☐ applicant only

☒ applicant and inventor

☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
GB

State (that is, country) of residence:
GB

This person is applicant for the purposes of:

☐ all designated States

☐ all designated States except the United States of America

☒ the United States of America only

☐ the States indicated in the Supplemental Box

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

MCKERRECHER, Darren
Alderley Park,
Macclesfield, Cheshire.
SK10 4TG. GB.

This person is:

☐ applicant only

☒ applicant and inventor

☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
GB

State (that is, country) of residence:
GB

This person is applicant for the purposes of:

☐ all designated States

☐ all designated States except the United States of America

☒ the United States of America only

☐ the States indicated in the Supplemental Box

☐ Further applicants and/or (further) inventors are indicated on another continuation sheet.

Box No.V DESIGNATION OF STATES

The following designations are made under Rule 4.9(a) (mark the applicable check-boxes; at least one must be marked):

Regional Patent

- ☒ **AP ARIPO Patent:** GH Ghana, GM Gambia, KE Kenya, LS Lesotho, MW Malawi, SD Sudan, SZ Swaziland, UG Uganda, ZW Zimbabwe, and any other State which is a Contracting State of the Harare Protocol and of the PCT
- ☒ **EA Eurasian Patent:** AM Armenia, AZ Azerbaijan, BY Belarus, KG Kyrgyzstan, KZ Kazakhstan, MD Republic of Moldova, RU Russian Federation, TJ Tajikistan, TM Turkmenistan, and any other State which is a Contracting State of the Eurasian Patent Convention and of the PCT
- ☒ **EP European Patent:** AT Austria, BE Belgium, CH and LI Switzerland and Liechtenstein, CY Cyprus, DE Germany, DK Denmark, ES Spain, FI Finland, FR France, GB United Kingdom, GR Greece, IE Ireland, IT Italy, LU Luxembourg, MC Monaco, NL Netherlands, PT Portugal, SE Sweden, and any other State which is a Contracting State of the European Patent Convention and of the PCT
- ☒ **OA OAPI Patent:** BF Burkina Faso, BJ Benin, CF Central African Republic, CG Congo, CI Côte d'Ivoire, CM Cameroon, GA Gabon, GN Guinea, GW Guinea-Bissau, ML Mali, MR Mauritania, NE Niger, SN Senegal, TD Chad, TG Togo, and any other State which is a member State of OAPI and a Contracting State of the PCT (if other kind of protection or treatment desired, specify on dotted line)

National Patent (If other kind of protection or treatment desired, specify on dotted line):

- | | |
|---|---|
| <input checked="" type="checkbox"/> AL Albania | <input checked="" type="checkbox"/> LS Lesotho |
| <input checked="" type="checkbox"/> AM Armenia | <input checked="" type="checkbox"/> LT Lithuania |
| <input checked="" type="checkbox"/> AT Austria | <input checked="" type="checkbox"/> LU Luxembourg |
| <input checked="" type="checkbox"/> AU Australia | <input checked="" type="checkbox"/> LV Latvia |
| <input checked="" type="checkbox"/> AZ Azerbaijan | <input checked="" type="checkbox"/> MD Republic of Moldova |
| <input checked="" type="checkbox"/> BA Bosnia and Herzegovina | <input checked="" type="checkbox"/> MG Madagascar |
| <input checked="" type="checkbox"/> BB Barbados | <input checked="" type="checkbox"/> MK The former Yugoslav Republic of Macedonia |
| <input checked="" type="checkbox"/> BG Bulgaria | <input checked="" type="checkbox"/> MN Mongolia |
| <input checked="" type="checkbox"/> BR Brazil | <input checked="" type="checkbox"/> MW Malawi |
| <input checked="" type="checkbox"/> BY Belarus | <input checked="" type="checkbox"/> MX Mexico |
| <input checked="" type="checkbox"/> CA Canada | <input checked="" type="checkbox"/> NO Norway |
| <input checked="" type="checkbox"/> CH and LI Switzerland and Liechtenstein | <input checked="" type="checkbox"/> NZ New Zealand |
| <input checked="" type="checkbox"/> CN China | <input checked="" type="checkbox"/> PL Poland |
| <input checked="" type="checkbox"/> CU Cuba | <input checked="" type="checkbox"/> PT Portugal |
| <input checked="" type="checkbox"/> CZ Czech Republic | <input checked="" type="checkbox"/> RO Romania |
| <input checked="" type="checkbox"/> DE Germany | <input checked="" type="checkbox"/> RU Russian Federation |
| <input checked="" type="checkbox"/> DK Denmark | <input checked="" type="checkbox"/> SD Sudan |
| <input checked="" type="checkbox"/> EE Estonia | <input checked="" type="checkbox"/> SE Sweden |
| <input checked="" type="checkbox"/> ES Spain | <input checked="" type="checkbox"/> SG Singapore |
| <input checked="" type="checkbox"/> FI Finland | <input checked="" type="checkbox"/> SI Slovenia |
| <input checked="" type="checkbox"/> GB United Kingdom | <input checked="" type="checkbox"/> SK Slovakia |
| <input checked="" type="checkbox"/> GD Grenada | <input checked="" type="checkbox"/> SL Sierra Leone |
| <input checked="" type="checkbox"/> GE Georgia | <input checked="" type="checkbox"/> TJ Tajikistan |
| <input checked="" type="checkbox"/> GH Ghana | <input checked="" type="checkbox"/> TM Turkmenistan |
| <input checked="" type="checkbox"/> GM Gambia | <input checked="" type="checkbox"/> TR Turkey |
| <input checked="" type="checkbox"/> HR Croatia | <input checked="" type="checkbox"/> TT Trinidad and Tobago |
| <input checked="" type="checkbox"/> HU Hungary | <input checked="" type="checkbox"/> UA Ukraine |
| <input checked="" type="checkbox"/> ID Indonesia | <input checked="" type="checkbox"/> UG Uganda |
| <input checked="" type="checkbox"/> IL Israel | <input checked="" type="checkbox"/> US United States of America |
| <input checked="" type="checkbox"/> IN India | <input checked="" type="checkbox"/> UZ Uzbekistan |
| <input checked="" type="checkbox"/> IS Iceland | <input checked="" type="checkbox"/> VN Viet Nam |
| <input checked="" type="checkbox"/> JP Japan | <input checked="" type="checkbox"/> YU Yugoslavia |
| <input checked="" type="checkbox"/> KE Kenya | <input checked="" type="checkbox"/> ZW Zimbabwe |
| <input checked="" type="checkbox"/> KG Kyrgyzstan | |
| <input checked="" type="checkbox"/> KP Democratic People's Republic of Korea | |
| <input checked="" type="checkbox"/> KR Republic of Korea | |
| <input checked="" type="checkbox"/> KZ Kazakhstan | |
| <input checked="" type="checkbox"/> LC Saint Lucia | |
| <input checked="" type="checkbox"/> LK Sri Lanka | |
| <input checked="" type="checkbox"/> LR Liberia | |

Check-boxes reserved for designating States (for the purposes of a national patent) which have become party to the PCT after issuance of this sheet:

- ☒ **DM** DOMINICA ☒ **ZA** SOUTH AFRICA
☒ **AE** UNITED ARAB EMIRATES ☒ **TZ** TANZANIA
☒ **CR** COSTA RICA ☒ **MA** MOROCCO

Precautionary Designation Statement: In addition to the designations made above, the applicant also makes under Rule 4.9(b) all other designations which would be permitted under the PCT except any designation(s) indicated in the Supplemental Box as being excluded from the scope of this statement. The applicant declares that those additional designations are subject to confirmation and that any designation which is not confirmed before the expiration of 15 months from the priority date is to be regarded as withdrawn by the applicant at the expiration of that time limit. (Confirmation of a designation consists of the filing of a notice specifying that designation and the payment of the designation and confirmation fees. Confirmation must reach the receiving Office

Box No. VI PRIORITY CLAIM☐ Further priority claims are indicated in the Supplemental Box.

Filing date of earlier application (day/month/year)	Number of earlier application	Where earlier application is:		
		national application: country	regional application: regional Office	international application: receiving Office
item (1) 10/02/1999 (10FEB99)	99400305.1		EP (FR)	
item (2)				
item (3)				

☐ The receiving Office is requested to prepare and transmit to the International Bureau a certified copy of the earlier application(s) (only if the earlier application was filed with the Office which for the purposes of the present international application is the receiving Office) identified above as item(s):

* Where the earlier application is an ARIPO application, it is mandatory to indicate in the Supplemental Box at least one country party to the Paris Convention for the Protection of Industrial Property for which that earlier application was filed (Rule 4.10(b)(ii)). See Supplemental Box.

Box No. VII INTERNATIONAL SEARCHING AUTHORITY

Choice of International Searching Authority (ISA)
(if two or more International Searching Authorities are competent to carry out the international search, indicate the Authority chosen; the two-letter code may be used):

ISA /

Request to use results of earlier search; reference to that search (if an earlier search has been carried out by or requested from the International Searching Authority):

Date (day/month/year)

Number

Country (or regional Office)

Box No. VIII CHECK LIST; LANGUAGE OF FILING

This international application contains the following number of sheets:

request : 4

description (excluding sequence listing part) : 314

claims : 26

abstract : 1

drawings :

sequence listing part of description :

Total number of sheets : 345

This international application is accompanied by the item(s) marked below:

1. ☒ fee calculation sheet
2. ☐ separate signed power of attorney
3. ☐ copy of general power of attorney; reference number, if any:
4. ☐ statement explaining lack of signature
5. ☐ priority document(s) identified in Box No. VI as item(s):
6. ☐ translation of international application into (language):
7. ☐ separate indications concerning deposited microorganism or other biological material
8. ☐ nucleotide and/or amino acid sequence listing in computer readable form
9. ☐ other (specify):

Figure of the drawings which should accompany the abstract:

Language of filing of the international application: ENGLISH

Box No. IX SIGNATURE OF APPLICANT OR AGENT

Next to each signature, indicate the name of the person signing and the capacity in which the person signs (if such capacity is not obvious from reading the request).



BRYANT, Tracey
AGENT FOR APPLICANT

For receiving Office use only:

1. Date of actual receipt of the purported international application:	2. Drawings: <input type="checkbox"/> received: <input type="checkbox"/> not received:
3. Corrected date of actual receipt due to later but timely received papers or drawings completing the purported international application:	
4. Date of timely receipt of the required corrections under PCT Article 11(2):	
5. International Searching Authority (if two or more are competent): ISA /	6. <input type="checkbox"/> Transmittal of search copy delayed until search fee is paid.

For International Bureau use only:

Date of receipt of the record copy by the International Bureau:

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference PHM70495/WO	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/GB 00/ 00373	International filing date (day/month/year) 08/02/2000	(Earliest) Priority Date (day/month/year) 10/02/1999
Applicant ZENECA LIMITED et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 4 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :

☐ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☐ furnished subsequently to this Authority in computer readable form.

☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

☐ the text is approved as submitted by the applicant.

☒ the text has been established by this Authority to read as follows:

QUINAZOLINE DERIVATIVES AS ANGIOGENESIS INHIBITORS

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 00/00373

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 22 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/00373

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/505 C07D4 4 C07D413/14 C07D417/12 D405/12
C07D401/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 29301 A (AGREVO UK LTD ; CORNELL CLIVE LEONARD (GB); RICHARDS IAN CHRISTOPHE) 26 September 1996 (1996-09-26) see compounds 67 and 83 ---	9-11, 14
X	WO 95 15758 A (HSU CHIN YI JENNY ; ZILBERSTEIN ASHER (US); JOHNSON SUSAN E (US); M) 15 June 1995 (1995-06-15) see compound on top of page 17 the whole document ---	9-11, 14, 19-21
Y	---	1-22
	- / - -	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

7 April 2000

Date of mailing of the international search report

18.05.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Scruton-Evans, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/00373

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	HENNEQUIN L F ET AL: "Design and Structure-Activity Relationship of a New Class of Potent VEGF Receptor Tyrosine Kinase Inhibitors" J. MED. CHEM. (JMCMAR,00222623);1999; VOL.42 (26); PP.5369-5389, XP000197180 AstraZeneca Zeneca Pharma Centre de Recherches Z.I. La Pompelle;Reims; 51689; Fr. (FR) the whole document	1-22
Y	BRIDGES A J ET AL: "Enantioselective inhibition of the epidermal growth factor receptor tyrosine kinase by 4-(.alpha.-phenethylamino)quinazolines" BIOORG. MED. CHEM. (BMECEP,09680896);1995; VOL.3 (12); PP.1651-6, XP002134974 Parke-Davis Pharmaceutical Research Division;Ann Arbor; 48105; MI; USA (US) the whole document	1-22
Y	GIBSON K H ET AL: "Epidermal growth factor receptor tyrosine kinase: structure-activity relationships and antitumor activity of novel quinazolines" BIOORG. MED. CHEM. LETT. (BMCLE8,0960894X);1997; VOL.7 (21); PP.2723-2728, XP002134975 Zeneca Pharmaceuticals;Research Dep. Cancer, Metabolism and Endocrine; Alderley Park, Macclesfield, Cheshire; SK10 4TG; UK (GB) the whole document	1-22
Y	EP 0 602 851 A (ZENECA LTD) 22 June 1994 (1994-06-22) the whole document	1-22
X	WO 97 42187 A (LOHMANN JEAN JACQUES MARCEL ;PLE PATRICK (FR); HENNEQUIN LAURENT F) 13 November 1997 (1997-11-13) see the general formula I and pages 1-3	1-3,6-8, 20
P,X	WO 99 10349 A (LOHMANN JEAN JACQUES MARCEL ;PLE PATRICK (FR); HENNEQUIN LAURENT F) 4 March 1999 (1999-03-04) the whole document	1-3,6-8, 20
Y	WO 97 22596 A (ZENECA LTD ;ZENECA PHARMA SA (FR); LOHMANN JEAN JACQUES MARCEL (FR) 26 June 1997 (1997-06-26) the whole document	1-22

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No


PCT/GB 00/00373

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9629301 A	26-09-1996	AU 5116096 A EP 0819111 A JP 11503114 T ZA 9602319 A	08-10-1996 21-01-1998 23-03-1999 30-10-1996
WO 9515758 A	15-06-1995	US 5480883 A US 5710158 A AU 1305095 A EP 0871448 A SG 54172 A US 5795889 A US 5646153 A US 5721237 A US 5714493 A US RE36256 E	02-01-1996 20-01-1998 27-06-1995 21-10-1998 16-11-1998 18-08-1998 08-07-1997 24-02-1998 03-02-1998 20-07-1999
EP 0602851 A	22-06-1994	AT 143956 T AU 664496 B AU 5072893 A CA 2103383 A CN 1094043 A CZ 9302651 A DE 69305310 D DE 69305310 T DK 602851 T ES 2093367 T FI 935431 A GR 3021326 T HU 65622 A IL 107678 A JP 6336481 A MX 9307778 A NO 934504 A NZ 250218 A US 5580870 A ZA 9308594 A	15-10-1996 16-11-1995 23-06-1994 11-06-1994 26-10-1994 15-06-1994 14-11-1996 13-02-1997 24-03-1997 16-12-1996 11-06-1994 31-01-1997 28-07-1994 12-03-1999 06-12-1994 29-07-1994 13-06-1994 25-06-1996 03-12-1996 10-06-1994
WO 9742187 A	13-11-1997	AU 2647597 A EP 0912557 A	26-11-1997 06-05-1999
WO 9910349 A	04-03-1999	AU 8816298 A	16-03-1999
WO 9722596 A	26-06-1997	AU 712370 B AU 1106197 A CZ 9801882 A EP 0873319 A HU 9901243 A NO 982784 A PL 327310 A SK 82898 A US 5962458 A	04-11-1999 14-07-1997 16-09-1998 28-10-1998 30-08-1999 17-08-1998 07-12-1998 04-11-1998 05-10-1999



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/505, C07D 401/14, 413/14, 417/12, 405/12, 401/12	A1	(11) International Publication Number: WO 00/47212 (43) International Publication Date: 17 August 2000 (17.08.00)
(21) International Application Number: PCT/GB00/00373 (22) International Filing Date: 8 February 2000 (08.02.00) (30) Priority Data: 99400305.1 10 February 1999 (10.02.99) EP (71) Applicants (for all designated States except US): AS-TRAZENECA UK LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). ZENECA-PHARMA S.A. [FR/FR]; Le Galien, 1, rue des Chauffours, Boîte postale 127, F-95022 Cergy Cedex (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): HENNEQUIN, Laurent, François, Andre [FR/FR]; Z.I. La Pompelle, Boîte postale 1050, F-51689 Reims Cedex 2 (FR). PLE, Patrick [FR/FR]; Z.I. La Pompelle, Boîte postale 1050, F-51689 Reims Cedex 2 (FR). STOKES, Elaine, Sophie, Elizabeth [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). MCKERRECHER, Darren [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (74) Agent: BRYANT, Tracey; AstraZeneca, Global Intellectual Property, Patents, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: QUINAZOLINE DERIVATIVES AS ANGIOGENESIS INHIBITORS

(I)

(57) Abstract

The invention relates to the use of compounds of formula (I), wherein ring C is an 8, 9, 10, 12 or 13-membered bicyclic or tricyclic moiety which optionally may contain 1-3 heteroatoms selected independently from O, N and S; Z is -O-, -NH-, -S-, -CH₂- or a direct bond; n is 0-5; m is 0-3; R² represents hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C₁-alkyl, C₁-alkoxy, C₁-alkylsulphanyl, -NR³R⁴ (wherein R³ and R⁴, which may be the same or different, each represents hydrogen or C₁-alkyl), or R⁵X¹- (wherein X¹ and R⁵ are as defined herein; R¹ represents hydrogen, oxo, halogeno, hydroxy, C₁-alkoxy, C₁-alkyl, C₁-alkoxymethyl, C₁-alkanoyl, C₁-haloalkyl, cyano, amino, C₂-alkenyl, C₂-alkynyl, C₁-3alkanoyloxy, nitro, C₁-4alkanoylamino, C₁-4alkoxycarbonyl, C₁-4alkylsulphanyl, C₁-4alkylsulphinyl, C₁-4alkylsulphonyl, carbamoyl, N-C₁-4alkylcarbamoyl, N,N-di(C₁-4alkyl)carbamoyl, aminosulphonyl, N-C₁-4alkylaminosulphonyl, N,N-di(C₁-4alkyl)aminosulphonyl, N-(C₁-4alkylsulphonyl)amino, N-(C₁-4alkylsulphonyl)-N-(C₁-4alkyl)amino, N,N-di(C₁-4alkylsulphonyl)amino, a C₃-7alkylene chain joined to two ring C carbon atoms, C₁-4alkanoylaminoC₁-4alkyl, carboxy or a group R⁵⁶X¹⁰ (wherein X¹⁰ and R⁵⁶ are as defined herein); and salts thereof, in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals, processes for the preparation of such compounds, pharmaceutical compositions containing a compound of formula (I) or a pharmaceutically acceptable salt thereof as active ingredient and compounds of formula (I). The compounds of formula (I) and the pharmaceutically acceptable salts thereof inhibit the effects of VEGF, a property of value in the treatment of a number of disease states including cancer and rheumatoid arthritis.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

QUINAZOLINE DERIVATIVES AS ANGIOGENESIS INHIBITORS

The present invention relates to quinazoline derivatives, processes for their preparation, pharmaceutical compositions containing them as active ingredient, methods for the treatment of disease states associated with angiogenesis and/or increased vascular permeability, to their use as medicaments and to their use in the manufacture of medicaments for use in the production of antiangiogenic and/or vascular permeability reducing effects in warm-blooded animals such as humans.


Normal angiogenesis plays an important role in a variety of processes including embryonic development, wound healing and several components of female reproductive function. Undesirable or pathological angiogenesis has been associated with disease states including diabetic retinopathy, psoriasis, cancer, rheumatoid arthritis, atheroma, Kaposi's sarcoma and haemangioma (Fan et al, 1995, Trends Pharmacol. Sci. 16: 57-66; Folkman, 1995, Nature Medicine 1: 27-31). Alteration of vascular permeability is thought to play a role in both normal and pathological physiological processes (Cullinan-Bove et al, 1993, Endocrinology 133: 829-837; Senger et al, 1993, Cancer and Metastasis Reviews, 12: 303-324). Several polypeptides with in vitro endothelial cell growth promoting activity have been identified including, acidic and basic fibroblast growth factors (aFGF & bFGF) and vascular endothelial growth factor (VEGF). By virtue of the restricted expression of its receptors, the growth factor activity of VEGF, in contrast to that of the FGFs, is relatively specific towards endothelial cells. Recent evidence indicates that VEGF is an important stimulator of both normal and pathological angiogenesis (Jakeman et al, 1993, Endocrinology, 133: 848-859; Kolch et al, 1995, Breast Cancer Research and Treatment, 36:139-155) and vascular permeability (Connolly et al, 1989, J. Biol. Chem. 264: 20017-20024). Antagonism of VEGF action by sequestration of VEGF with antibody can result in inhibition of tumour growth (Kim et al, 1993, Nature 362: 841-844). Basic FGF (bFGF) is a potent stimulator of angiogenesis (e.g. Hayek et al, 1987, Biochem. Biophys. Res. Commun. 147: 876-880) and raised levels of FGFs have been found in the serum (Fujimoto et al, 1991, Biochem. Biophys. Res. Commun. 180: 386-392) and urine (Nguyen et al, 1993, J. Natl. Cancer. Inst. 85: 241-242) of patients with cancer.

Receptor tyrosine kinases (RTKs) are important in the transmission of biochemical signals across the plasma membrane of cells. These transmembrane molecules

characteristically consist of an extracellular ligand-binding domain connected through a segment in the plasma membrane to an intracellular tyrosine kinase domain. Binding of ligand to the receptor results in stimulation of the receptor-associated tyrosine kinase activity which leads to phosphorylation of tyrosine residues on both the receptor and other intracellular molecules. These changes in tyrosine phosphorylation initiate a signalling cascade leading to a variety of cellular responses. To date, at least nineteen distinct RTK subfamilies, defined by amino acid sequence homology, have been identified. One of these subfamilies is presently comprised by the fms-like tyrosine kinase receptor, Flt or Flt1, the kinase insert domain-containing receptor, KDR (also referred to as Flk-1), and another fms-like tyrosine kinase receptor, Flt4. Two of these related RTKs, Flt and KDR, have been shown to bind VEGF with high affinity (De Vries et al, 1992; Science 255: 989-991; Terman et al, 1992, Biochem. Biophys. Res. Comm. 1992, 187: 1579-1586). Binding of VEGF to these receptors expressed in heterologous cells has been associated with changes in the tyrosine phosphorylation status of cellular proteins and calcium fluxes.

The present invention is based on the discovery of compounds that surprisingly inhibit the effects of VEGF, a property of value in the treatment of disease states associated with angiogenesis and/or increased vascular permeability such as cancer, diabetes, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, arterial restenosis, autoimmune diseases, acute inflammation, excessive scar formation and adhesions, endometriosis, dysfunctional uterine bleeding and ocular diseases with retinal vessel proliferation. Compounds of the present invention generally possess higher potency against VEGF receptor tyrosine kinase than against epidermal growth factor (EGF) receptor tyrosine kinase. Compounds of the invention which have been tested possess activity against VEGF receptor tyrosine kinase such that they may be used in an amount sufficient to inhibit VEGF receptor tyrosine kinase whilst demonstrating no significant activity against EGF receptor tyrosine kinase. Compounds of the present invention generally possess higher potency against VEGF receptor tyrosine kinase than against FGF R1 receptor tyrosine kinase. Compounds of the invention which have been tested possess activity against VEGF receptor tyrosine kinase such that they may be used in an amount sufficient to inhibit VEGF receptor tyrosine kinase whilst demonstrating no significant activity against FGF R1 receptor tyrosine kinase.

According to one aspect of the present invention there is provided the use of a compound of the formula I:



(I)

wherein:

ring C is an 8, 9, 10, 12 or 13-membered bicyclic or tricyclic moiety which moiety may be saturated or unsaturated, which may be aromatic or non-aromatic, and which optionally may

contain 1-3 heteroatoms selected independently from O, N and S;

Z is -O-, -NH-, -S-, -CH₂- or a direct bond;

n is an integer from 0 to 5;

m is an integer from 0 to 3;

R² represents hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C₁₋₃alkyl, C₁₋

₃alkoxy, C₁₋₃alkylsulphanyl, -NR³R⁴ (wherein R³ and R⁴, which may be the same or different, each represents hydrogen or C₁₋₃alkyl), or R⁵X¹- (wherein X¹ represents a direct bond, -O-, -CH₂-, -OC(O)-, -C(O)-, -S-, -SO-, -SO₂-, -NR⁶C(O)-, -C(O)NR⁷-, -SO₂NR⁸-, -NR⁹SO₂- or -NR¹⁰- (wherein R⁶, R⁷, R⁸, R⁹ and R¹⁰ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), and R⁵ is selected from one of the following twenty-two groups:

1) hydrogen, oxiranylC₁₋₄alkyl or C₁₋₅alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, chloro, bromo and amino;

2) C₁₋₅alkylX²C(O)R¹¹ (wherein X² represents -O- or -NR¹²- (in which R¹² represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹¹ represents C₁₋₃alkyl, -NR¹³R¹⁴ or -OR¹⁵ (wherein R¹³, R¹⁴ and R¹⁵ which may be the same or different each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl));

3) C₁₋₅alkylX³R¹⁶ (wherein X³ represents -O-, -S-, -SO-, -SO₂-, -OC(O)-, -NR¹⁷C(O)-, -C(O)NR¹⁸-, -SO₂NR¹⁹-, -NR²⁰SO₂- or -NR²¹- (wherein R¹⁷, R¹⁸, R¹⁹, R²⁰ and R²¹ each

independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹⁶ represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group $-(O-)_f(C_{1-4}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));

4) C₁₋₅alkylX⁴C₁₋₅alkylX⁵R²² (wherein X⁴ and X⁵ which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR²³C(O)-, -C(O)NR²⁴-, -SO₂NR²⁵-, -NR²⁶SO₂- or -NR²⁷- (wherein R²³, R²⁴, R²⁵, R²⁶ and R²⁷ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²² represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl);

5) R²⁸ (wherein R²⁸ is a 5-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group $-(O-)_f(C_{1-4}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));

6) C₁₋₅alkylR²⁸ (wherein R²⁸ is as defined hereinbefore);

7) C₂₋₅alkenylR²⁸ (wherein R²⁸ is as defined hereinbefore);

8) C₂₋₅alkynylR²⁸ (wherein R²⁸ is as defined hereinbefore);

9) R²⁹ (wherein R²⁹ represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents

- selected from hydroxy, halogeno, amino, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄hydroxyalkyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, C₁₋₄hydroxyalkoxy, carboxy, trifluoromethyl, cyano, -C(O)NR³⁰R³¹, -NR³²C(O)R³³ (wherein R³⁰, R³¹, R³² and R³³, which may be the same or different, each represents hydrogen, C₁₋₄alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and a group -(O)_f(C₁₋₄alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));
- 5 10) C₁₋₅alkylR²⁹ (wherein R²⁹ is as defined hereinbefore);
- 11) C₂₋₅alkenylR²⁹ (wherein R²⁹ is as defined hereinbefore);
- 10 12) C₂₋₅alkynylR²⁹ (wherein R²⁹ is as defined hereinbefore);
- 13) C₁₋₅alkylX⁶R²⁹ (wherein X⁶ represents -O-, -S-, -SO-, -SO₂-, -NR³⁴C(O)-, -C(O)NR³⁵-, -SO₂NR³⁶-, -NR³⁷SO₂- or -NR³⁸- (wherein R³⁴, R³⁵, R³⁶, R³⁷ and R³⁸ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);
- 14) C₂₋₅alkenylX⁷R²⁹ (wherein X⁷ represents -O-, -S-, -SO-, -SO₂-, -NR³⁹C(O)-, -C(O)NR⁴⁰-, -SO₂NR⁴¹-, -NR⁴²SO₂- or -NR⁴³- (wherein R³⁹, R⁴⁰, R⁴¹, R⁴² and R⁴³ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);
- 15 15) C₂₋₅alkynylX⁸R²⁹ (wherein X⁸ represents -O-, -S-, -SO-, -SO₂-, -NR⁴⁴C(O)-, -C(O)NR⁴⁵-, -SO₂NR⁴⁶-, -NR⁴⁷SO₂- or -NR⁴⁸- (wherein R⁴⁴, R⁴⁵, R⁴⁶, R⁴⁷ and R⁴⁸ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);
- 20 16) C₁₋₄alkylX⁹C₁₋₄alkylR²⁹ (wherein X⁹ represents -O-, -S-, -SO-, -SO₂-, -NR⁴⁹C(O)-, -C(O)NR⁵⁰-, -SO₂NR⁵¹-, -NR⁵²SO₂- or -NR⁵³- (wherein R⁴⁹, R⁵⁰, R⁵¹, R⁵² and R⁵³ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);
- 17) C₁₋₄alkylX⁹C₁₋₄alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);
- 25 18) C₂₋₅alkenyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;
- 19) C₂₋₅alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;
- 30 20) C₂₋₅alkenylX⁹C₁₋₄alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);
- 21) C₂₋₅alkynylX⁹C₁₋₄alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore); and

- 22) $C_{1-4}alkylR^{54}(C_{1-4}alkyl)_q(X^9)_rR^{55}$ (wherein X^9 is as defined hereinbefore, q is 0 or 1, r is 0 or 1, and R^{54} and R^{55} are each independently selected from hydrogen, $C_{1-3}alkyl$, cyclopentyl, cyclohexyl and a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which $C_{1-3}alkyl$ group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and $C_{1-4}alkoxy$ and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, $C_{1-4}cyanoalkyl$, $C_{1-4}alkyl$, $C_{1-4}hydroxyalkyl$, $C_{1-4}alkoxy$, $C_{1-4}alkoxyC_{1-4}alkyl$, $C_{1-4}alkylsulphonylC_{1-4}alkyl$, $C_{1-4}alkoxycarbonyl$, $C_{1-4}aminoalkyl$, $C_{1-4}alkylamino$, $di(C_{1-4}alkyl)amino$, $C_{1-4}alkylaminoC_{1-4}alkyl$, $di(C_{1-4}alkyl)aminoC_{1-4}alkyl$, $C_{1-4}alkylaminoC_{1-4}alkoxy$, $di(C_{1-4}alkyl)aminoC_{1-4}alkoxy$ and a group $-(O-)_f(C_{1-4}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from $C_{1-4}alkyl$), with the proviso that R^{54} cannot be hydrogen);
- and additionally wherein any $C_{1-3}alkyl$, $C_{2-5}alkenyl$ or $C_{2-5}alkynyl$ group in R^5X^1 may bear one or more substituents selected from hydroxy, halogeno and amino);
- R^1 represents hydrogen, oxo, halogeno, hydroxy, $C_{1-4}alkoxy$, $C_{1-4}alkyl$, $C_{1-4}alkoxymethyl$, $C_{1-4}alkanoyl$, $C_{1-4}haloalkyl$, cyano, amino, $C_{2-5}alkenyl$, $C_{2-5}alkynyl$, $C_{1-3}alkanoyloxy$, nitro, $C_{1-4}alkanoylamino$, $C_{1-4}alkoxycarbonyl$, $C_{1-4}alkylsulphanyl$, $C_{1-4}alkylsulphinyl$, $C_{1-4}alkylsulphonyl$, carbamoyl, \underline{N} - $C_{1-4}alkylcarbamoyl$, $\underline{N,N}$ - $di(C_{1-4}alkyl)carbamoyl$, aminosulphonyl, \underline{N} - $C_{1-4}alkylaminosulphonyl$, $\underline{N,N}$ - $di(C_{1-4}alkyl)aminosulphonyl$, \underline{N} -($C_{1-4}alkylsulphonyl$)amino, \underline{N} -($C_{1-4}alkylsulphonyl$)- \underline{N} -($C_{1-4}alkyl$)amino, $\underline{N,N}$ - $di(C_{1-4}alkylsulphonyl)amino$, a $C_{3-7}alkylene$ chain joined to two ring C carbon atoms, $C_{1-4}alkanoylaminoC_{1-4}alkyl$, carboxy or a group $R^{56}X^{10}$ (wherein X^{10} represents a direct bond, -O-, -CH₂-, -OC(O)-, -C(O)-, -S-, -SO-, -SO₂-, -NR⁵⁷C(O)-, -C(O)NR⁵⁸-, -SO₂NR⁵⁹-, -NR⁶⁰SO₂- or -NR⁶¹- (wherein R^{57} , R^{58} , R^{59} , R^{60} and R^{61} each independently represents hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$), and R^{56} is selected from one of the following twenty-two groups:
- 1) hydrogen, oxiranyl $C_{1-4}alkyl$ or $C_{1-3}alkyl$ which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, chloro, bromo and amino;
 - 2) $C_{1-3}alkylX^{11}C(O)R^{62}$ (wherein X^{11} represents -O- or -NR⁶³- (in which R^{63} represents hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$) and R^{62} represents $C_{1-3}alkyl$, -NR⁶⁴R⁶⁵ or -OR⁶⁶ (wherein R^{64} , R^{65} and R^{66} which may be the same or different each represents hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$));

- 3) $C_{1-3}alkylX^{12}R^{67}$ (wherein X^{12} represents -O-, -S-, -SO-, -SO₂-, -OC(O)-, -NR⁶⁸C(O)-, -C(O)NR⁶⁹-, -SO₂NR⁷⁰-, -NR⁷¹SO₂- or -NR⁷²- (wherein R⁶⁸, R⁶⁹, R⁷⁰, R⁷¹ and R⁷² each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁶⁷ represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group $-(O-)_f(C_{1-4}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));
- 4) $C_{1-5}alkylX^{13}C_{1-5}alkylX^{14}R^{73}$ (wherein X^{13} and X^{14} which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR⁷⁴C(O)-, -C(O)NR⁷⁵-, -SO₂NR⁷⁶-, -NR⁷⁷SO₂- or -NR⁷⁸- (wherein R⁷⁴, R⁷⁵, R⁷⁶, R⁷⁷ and R⁷⁸ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁷³ represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl);
- 5) R⁷⁹ (wherein R⁷⁹ is a 5-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group $-(O-)_f(C_{1-4}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));
- 6) C₁₋₅alkylR⁷⁹ (wherein R⁷⁹ is as defined hereinbefore);
- 7) C₂₋₅alkenylR⁷⁹ (wherein R⁷⁹ is as defined hereinbefore);
- 8) C₂₋₅alkynylR⁷⁹ (wherein R⁷⁹ is as defined hereinbefore);

- 9) R^{80} (wherein R^{80} represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} hydroxyalkyl, C_{1-4} aminoalkyl, C_{1-4} alkylamino, C_{1-4} hydroxyalkoxy, carboxy, trifluoromethyl, cyano, -
 5 $C(O)NR^{81}R^{82}$, $-NR^{83}C(O)R^{84}$ (wherein R^{81} , R^{82} , R^{83} and R^{84} , which may be the same or different, each represents hydrogen, C_{1-4} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and a group $-(O-)_f(C_{1-4}$ alkyl) $_g$ ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which
 10 cyclic group may bear one or more substituents selected from C_{1-4} alkyl));
- 10) C_{1-5} alkyl R^{80} (wherein R^{80} is as defined hereinbefore);
- 11) C_{2-5} alkenyl R^{80} (wherein R^{80} is as defined hereinbefore);
- 12) C_{2-5} alkynyl R^{80} (wherein R^{80} is as defined hereinbefore);
- 13) C_{1-5} alkyl $X^{15}R^{80}$ (wherein X^{15} represents -O-, -S-, -SO-, -SO₂-, $-NR^{85}C(O)-$, $-C(O)NR^{86}-$, -
 15 SO₂NR⁸⁷-, $-NR^{88}SO_2-$ or $-NR^{89}-$ (wherein R^{85} , R^{86} , R^{87} , R^{88} and R^{89} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{80} is as defined hereinbefore);
- 14) C_{2-5} alkenyl $X^{16}R^{80}$ (wherein X^{16} represents -O-, -S-, -SO-, -SO₂-, $-NR^{90}C(O)-$, $-C(O)NR^{91}-$,
 -SO₂NR⁹²-, $-NR^{93}SO_2-$ or $-NR^{94}-$ (wherein R^{90} , R^{91} , R^{92} , R^{93} and R^{94} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{80} is as defined hereinbefore);
- 20 15) C_{2-5} alkynyl $X^{17}R^{80}$ (wherein X^{17} represents -O-, -S-, -SO-, -SO₂-, $-NR^{95}C(O)-$, $-C(O)NR^{96}-$,
 -SO₂NR⁹⁷-, $-NR^{98}SO_2-$ or $-NR^{99}-$ (wherein R^{95} , R^{96} , R^{97} , R^{98} and R^{99} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{80} is as defined hereinbefore);
- 25 16) C_{1-4} alkyl $X^{18}C_{1-4}$ alkyl R^{80} (wherein X^{18} represents -O-, -S-, -SO-, -SO₂-, $-NR^{100}C(O)-$,
 $-C(O)NR^{101}-$, $-SO_2NR^{102}-$, $-NR^{103}SO_2-$ or $-NR^{104}-$ (wherein R^{100} , R^{101} , R^{102} , R^{103} and R^{104} each
 30 independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{80} is as defined hereinbefore);
- 17) C_{1-4} alkyl $X^{18}C_{1-4}$ alkyl R^{79} (wherein X^{18} and R^{79} are as defined hereinbefore);
- 18) C_{2-5} alkenyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C_{1-4} alkylamino, N,N-di(C_{1-4} alkyl)amino, aminosulphonyl, N- C_{1-4} alkylaminosulphonyl and N,N-di(C_{1-4} alkyl)aminosulphonyl;

19) C₂₋₃alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;

20) C₂₋₃alkenylX¹⁸C₁₋₄alkylR⁷⁹ (wherein X¹⁸ and R⁷⁹ are as defined hereinbefore);


5 21) C₂₋₃alkynylX¹⁸C₁₋₄alkylR⁷⁹ (wherein X¹⁸ and R⁷⁹ are as defined hereinbefore); and

22) C₁₋₄alkylR¹⁰⁵(C₁₋₄alkyl)_x(X¹⁸)_yR¹⁰⁶ (wherein X¹⁸ is as defined hereinbefore, x is 0 or 1, y is 0 or 1, and R¹⁰⁵ and R¹⁰⁶ are each independently selected from hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl and a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected

10 from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group -(O-)_f(C₁₋₄alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl) with the proviso that R¹⁰⁵ cannot be hydrogen);

and additionally wherein any C₁₋₃alkyl, C₂₋₃alkenyl or C₂₋₃alkynyl group in R⁵⁶X¹⁰- may bear
 20 one or more substituents selected from hydroxy, halogeno and amino);
 or a salt thereof, or a prodrug thereof for example an ester or an amide, in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals such as humans.

According to another aspect of the present invention there is provided the use of
 25 compounds of the formula I:



(I)

wherein:

- 5 ring C is a 9-10-membered bicyclic moiety which may be saturated or unsaturated, which may be aromatic or non-aromatic, and which optionally may contain 1-3 heteroatoms selected independently from O, N and S;
- Z is -O-, -NH-, -S-, -CH₂- or a direct bond;
- R¹ represents hydrogen, oxo, halogeno, hydroxy, C₁₋₄alkoxy, C₁₋₄alkyl, C₁₋₄alkoxymethyl, C₁₋₄alkanoyl, C₁₋₄haloalkyl, cyano, amino, C₂₋₅alkenyl, C₂₋₅alkynyl, C₁₋₃alkanoyloxy, nitro, C₁₋₄alkanoylamino, C₁₋₄alkoxycarbonyl, C₁₋₄alkylsulphanyl, C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, carbamoyl, N-C₁₋₄alkylcarbamoyl, N,N-di(C₁₋₄alkyl)carbamoyl, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl, N,N-di(C₁₋₄alkyl)aminosulphonyl, N-(C₁₋₄alkylsulphonyl)amino, N-(C₁₋₄alkylsulphonyl)-N-(C₁₋₄alkyl)amino, N,N-di(C₁₋₄alkylsulphonyl)amino or a C₃₋₇alkylene chain joined to two ring C carbon atoms;
- 10 n is an integer from 0 to 5;
- m is an integer from 0 to 3;
- R² represents hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C₁₋₃alkyl, C₁₋₃alkoxy, C₁₋₃alkylsulphanyl, -NR³R⁴ (wherein R³ and R⁴, which may be the same or different, each represents hydrogen or C₁₋₃alkyl), or R⁵X¹- (wherein X¹ represents a direct bond, -O-, -CH₂-, -OC(O)-, -C(O)-, -S-, -SO-, -SO₂-, -NR⁶C(O)-, -C(O)NR⁷-, -SO₂NR⁸-, -NR⁹SO₂- or -NR¹⁰- (wherein R⁶, R⁷, R⁸, R⁹ and R¹⁰ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), and R⁵ is selected from one of the following twenty-one groups:
- 1) hydrogen or C₁₋₃alkyl which may be unsubstituted or which may be substituted with one or
- 25 more groups selected from hydroxy, fluoro and amino;
- 2) C₁₋₅alkylX²C(O)R¹¹ (wherein X² represents -O- or -NR¹²- (in which R¹² represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹¹ represents C₁₋₃alkyl, -NR¹³R¹⁴ or -OR¹⁵

(wherein R^{13} , R^{14} and R^{15} which may be the same or different each represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));

3) C_{1-5} alkyl X^3R^{16} (wherein X^3 represents -O-, -S-, -SO-, -SO₂-, -OC(O)-, -NR¹⁷C(O)-, -C(O)NR¹⁸-, -SO₂NR¹⁹-, -NR²⁰SO₂- or -NR²¹- (wherein R^{17} , R^{18} , R^{19} , R^{20} and R^{21} each

5 independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{16} represents hydrogen, C_{1-3} alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C_{1-3} alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C_{1-4} alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);

4) C_{1-5} alkyl X^4C_{1-5} alkyl X^5R^{22} (wherein X^4 and X^5 which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR²³C(O)-, -C(O)NR²⁴-, -SO₂NR²⁵-, -NR²⁶SO₂- or -NR²⁷- (wherein R^{23} , R^{24} , R^{25} , R^{26} and R^{27} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{22} represents hydrogen or C_{1-3} alkyl);

5) R^{28} (wherein R^{28} is a 5-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-4} cyanoalkyl, C_{1-4} alkyl, C_{1-4} hydroxyalkyl, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl and C_{1-4} alkylsulphonyl C_{1-4} alkyl);

6) C_{1-5} alkyl R^{28} (wherein R^{28} is as defined hereinbefore);

7) C_{2-5} alkenyl R^{28} (wherein R^{28} is as defined hereinbefore);

8) C_{2-5} alkynyl R^{28} (wherein R^{28} is as defined hereinbefore);

9) R^{29} (wherein R^{29} represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N

and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents on an available carbon atom selected from hydroxy, halogeno, amino, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} hydroxyalkyl, C_{1-4} aminoalkyl, C_{1-4} alkylamino, C_{1-4} hydroxyalkoxy, carboxy, trifluoromethyl, cyano, -C(O)NR³⁰R³¹ and -NR³²C(O)R³³ (wherein R^{30} , R^{31} , R^{32} and R^{33} , which may be the same or different, each represents hydrogen, C_{1-4} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));

10) C_{1-5} alkyl R^{29} (wherein R^{29} is as defined hereinbefore);

11) C_{2-5} alkenyl R^{29} (wherein R^{29} is as defined hereinbefore);

12) C_{2-5} alkynyl R^{29} (wherein R^{29} is as defined hereinbefore);

- 13) $C_{1-5}alkylX^6R^{29}$ (wherein X^6 represents -O-, -S-, -SO-, -SO₂-, -NR³⁴C(O)-, -C(O)NR³⁵-, -SO₂NR³⁶-, -NR³⁷SO₂- or -NR³⁸- (wherein R³⁴, R³⁵, R³⁶, R³⁷ and R³⁸ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);
- 14) $C_{2-5}alkenylX^7R^{29}$ (wherein X^7 represents -O-, -S-, -SO-, -SO₂-, -NR³⁹C(O)-, -C(O)NR⁴⁰-, -SO₂NR⁴¹-, -NR⁴²SO₂- or -NR⁴³- (wherein R³⁹, R⁴⁰, R⁴¹, R⁴² and R⁴³ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);
- 15) $C_{2-5}alkynylX^8R^{29}$ (wherein X^8 represents -O-, -S-, -SO-, -SO₂-, -NR⁴⁴C(O)-, -C(O)NR⁴⁵-, -SO₂NR⁴⁶-, -NR⁴⁷SO₂- or -NR⁴⁸- (wherein R⁴⁴, R⁴⁵, R⁴⁶, R⁴⁷ and R⁴⁸ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);
- 16) $C_{1-3}alkylX^9C_{1-3}alkylR^{29}$ (wherein X^9 represents -O-, -S-, -SO-, -SO₂-, -NR⁴⁹C(O)-, -C(O)NR⁵⁰-, -SO₂NR⁵¹-, -NR⁵²SO₂- or -NR⁵³- (wherein R⁴⁹, R⁵⁰, R⁵¹, R⁵² and R⁵³ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);
- 17) $C_{1-3}alkylX^9C_{1-3}alkylR^{28}$ (wherein X^9 and R²⁸ are as defined hereinbefore);
- 18) $C_{2-5}alkenyl$ which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;
- 19) $C_{2-5}alkynyl$ which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;
- 20) $C_{2-5}alkenylX^9C_{1-4}alkylR^{28}$ (wherein X^9 and R²⁸ are as defined hereinbefore); and
- 21) $C_{2-5}alkynylX^9C_{1-4}alkylR^{28}$ (wherein X^9 and R²⁸ are as defined hereinbefore);
- and salts thereof, and prodrugs thereof for example esters, amides and sulphides, in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals such as humans.

Preferably ring C is a 9-10-membered aromatic bicyclic moiety which may optionally contain 1-3 heteroatoms selected independently from O, N and S.

More preferably ring C is a 9-10-membered heteroaromatic bicyclic moiety which contains 1-3 heteroatoms selected independently from O, N and S.

Particularly ring C is a 9-10-membered heteroaromatic bicyclic moiety which contains 1 or 2 nitrogen atoms.

According to one aspect of the present invention ring C is a 9-membered heteroaromatic bicyclic moiety which contains 1 or 2 nitrogen atoms, for example indolyl.

According to another aspect of the present invention ring C is a 10-membered heteroaromatic bicyclic moiety which contains 1 or 2 nitrogen atoms, for example quinoliny.

5 Especially ring C is indolyl or quinoliny.

Preferably Z is -O-, -NH-, -S- or a direct bond.

More preferably Z is -O-, -NH- or -S-.

Particularly Z is -O- or -S-, especially -O-.

Advantageously X^{10} represents a direct bond, -O-, -S-, $-NR^{57}C(O)-$, $-NR^{60}SO_2-$ or -
10 $NR^{61}-$ (wherein R^{57} , R^{60} and R^{61} each independently represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl).

Preferably X^{10} represents a direct bond, -O-, -S-, $-NR^{57}C(O)-$, $-NR^{60}SO_2-$ (wherein R^{57} and R^{60} each independently represents hydrogen or C_{1-2} alkyl) or NH.

More preferably X^{10} represents -O-, -S-, $-NR^{57}C(O)-$ (wherein R^{57} represents hydrogen
15 or C_{1-2} alkyl) or NH.

Particularly X^{10} represents -O- or $-NR^{57}C(O)-$ (wherein R^{57} represents hydrogen or C_{1-2} alkyl), more particularly -O- or $-NHC(O)-$, especially -O-.

According to another aspect of the present invention X^{10} represents -O- or a direct bond.

20 Advantageously X^{12} represents -O-, -S-, -SO-, $-SO_2-$, $-NR^{68}C(O)-$, $-NR^{71}SO_2-$ or $-NR^{72}-$ (wherein R^{68} , R^{71} and R^{72} each independently represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl).

Preferably X^{12} represents -O-, -S-, -SO-, $-SO_2-$ or $-NR^{72}-$ (wherein R^{72} represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl).

25 More preferably X^{12} represents -O- or $-NR^{72}-$ (wherein R^{72} represents hydrogen or C_{1-2} alkyl).

According to another aspect of the present invention X^{12} represents -O-, $-SO_2-$, $-NR^{71}SO_2-$ or $-NR^{72}-$ (wherein R^{71} and R^{72} each independently represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl).

30 Advantageously X^{18} represents -O-, -S- or $-NR^{104}-$ (wherein R^{104} represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl).

Preferably X^{18} represents -O- or $-NR^{104}$ - (wherein R^{104} represents hydrogen or C_{1-3} alkyl).

According to another aspect of the present invention X^{18} represents -O-, $-CONR^{101}$ - or $-NR^{104}$ - (wherein R^{101} and R^{104} each independently represents hydrogen or C_{1-2} alkyl).

5 Advantageously R^{67} represents a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C_{1-3} alkyl).

15 Preferably R^{67} is pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C_{1-3} alkyl).

25 More preferably R^{67} is pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

30 Particularly R^{67} is pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from a group $-(O-)_f(C_{1-3}$

,alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

Preferably R⁷⁹ is pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl).

More preferably R⁷⁹ is pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

Particularly R⁷⁹ is pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

Advantageously R¹⁰⁵ and R¹⁰⁶ are each independently a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl).

Preferably R^{105} and R^{106} are each independently selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(O)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C_{1-3} alkyl).

More preferably R^{105} and R^{106} are each independently selected from pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(O)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

Particularly R^{105} and R^{106} are each independently selected from pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino which group may bear 1 or 2 substituents selected from a group $-(O)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

Advantageously R^1 represents oxo, halogeno, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, C_{1-4} alkoxymethyl, C_{1-4} alkanoyl, C_{1-4} haloalkyl, cyano, amino, C_{2-3} alkenyl, C_{2-3} alkynyl, C_{1-3} alkanoyloxy, nitro, C_{1-4} alkanoylamino, C_{1-4} alkoxycarbonyl, C_{1-4} alkylsulphanyl, C_{1-4} alkylsulphinyl, C_{1-4} alkylsulphonyl, carbamoyl, \underline{N} - C_{1-4} alkylcarbamoyl, $\underline{N,N}$ -di(C_{1-4} alkyl)carbamoyl, aminosulphonyl, \underline{N} - C_{1-4} alkylaminosulphonyl, $\underline{N,N}$ -di(C_{1-4} alkyl)aminosulphonyl, \underline{N} -(C_{1-4} alkylsulphonyl)amino, \underline{N} -(C_{1-4} alkylsulphonyl)- \underline{N} -(C_{1-4} alkyl)amino, $\underline{N,N}$ -di(C_{1-4} alkylsulphonyl)amino, a $C_{3,7}$ alkylene chain joined to two ring C

carbon atoms, C₁₋₄alkanoylaminoC₁₋₄alkyl, carboxy or a group R⁵⁶X¹⁰ (wherein X¹⁰ is as defined hereinbefore and R⁵⁶ is selected from one of the following nine groups:

- 1) C₁₋₅alkylX¹²R⁶⁷ (wherein X¹² and R⁶⁷ are as defined hereinbefore);
- 2) R⁷⁹ (wherein R⁷⁹ is as defined hereinbefore);
- 5 3) C₁₋₅alkylR⁷⁹ (wherein R⁷⁹ is as defined hereinbefore);
- 4) C₂₋₅alkenylR⁷⁹ (wherein R⁷⁹ is as defined hereinbefore);
- 5) C₂₋₅alkynylR⁷⁹ (wherein R⁷⁹ is as defined hereinbefore);
- 6) C₁₋₃alkylX¹⁸C₁₋₃alkylR⁷⁹ (wherein X¹⁸ and R⁷⁹ are as defined hereinbefore);
- 7) C₂₋₅alkenylX¹⁸C₁₋₄alkylR⁷⁹ (wherein X¹⁸ and R⁷⁹ are as defined hereinbefore);
- 10 8) C₂₋₅alkynylX¹⁸C₁₋₄alkylR⁷⁹ (wherein X¹⁸ and R⁷⁹ are as defined hereinbefore); and
- 9) C₁₋₃alkylR¹⁰⁵(C₁₋₃alkyl)_x(X¹⁸)_yR¹⁰⁶ (wherein X¹⁸, x, y, R¹⁰⁵ and R¹⁰⁶ are as defined hereinbefore;

and additionally wherein any C₁₋₅alkyl, C₂₋₅alkenyl or C₂₋₅alkynyl group in R⁵⁶X¹⁰ - may bear one or more substituents selected from hydroxy, halogeno and amino,

- 15 with the proviso that when X¹⁰ is a direct bond R⁵⁶ is not R⁷⁹).

Preferably R¹ represents oxo, halogeno, hydroxy, C₁₋₂alkoxy, C₁₋₂alkyl, C₁₋₂alkoxymethyl, C₂₋₃alkanoyl, C₁₋₂haloalkyl, cyano, amino, C₂₋₄alkenyl, C₂₋₄alkynyl, C₂₋₃alkanoyloxy, nitro, C₂₋₃alkanoylamino, C₁₋₂alkoxycarbonyl, C₁₋₂alkylsulphanyl, C₁₋₂alkylsulphinyl, C₁₋₂alkylsulphonyl, carbamoyl, N-C₁₋₂alkylcarbamoyl, N,N-di(C₁₋₂alkyl)carbamoyl, aminosulphonyl, N-C₁₋₂alkylaminosulphonyl, N,N-di(C₁₋₂alkyl)aminosulphonyl, N-(C₁₋₂alkylsulphonyl)amino, N-(C₁₋₂alkylsulphonyl)-N-(C₁₋₂alkyl)amino or a C₃₋₇alkylene chain joined to two ring C carbon atoms.

More preferably R¹ represents oxo, hydroxy, C₁₋₂alkoxymethyl, amino, halogeno, C₁₋₂alkyl, C₁₋₂alkoxy, trifluoromethyl, cyano, nitro, C₂₋₃alkanoyl.

- 25 Particularly R¹ represents methyl, ethyl, trifluoromethyl or halogeno.

Especially R¹ represents methyl, fluoro, chloro or bromo, more especially methyl or fluoro.

Preferably n is an integer from 0 to 3.

More preferably n is 0, 1 or 2.

- 30 Preferably m is an integer from 0 to 2, more preferably 1 or 2, most preferably 2.

Advantageously X¹ represents a direct bond, -O-, -S-, -NR⁶C(O)-, -NR⁹SO₂- or -NR¹⁰- (wherein R⁶, R⁹ and R¹⁰ each independently represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^1 represents a direct bond, -O-, -S-, -NR⁶C(O)-, -NR⁹SO₂- (wherein R⁶ and R⁹ each independently represents hydrogen or C₁₋₂alkyl) or NH.

More preferably X^1 represents -O-, -S-, -NR⁶C(O)- (wherein R⁶ represents hydrogen or C₁₋₂alkyl) or NH.

5 Particularly X^1 represents -O- or -NR⁶C(O)- (wherein R⁶ represents hydrogen or C₁₋₂alkyl), more particularly -O- or -NHC(O)-, especially -O-.

According to another aspect of the present invention X^1 represents -O- or a direct bond.

10 Advantageously X^2 represents -O- or NR¹² (wherein R¹² represents hydrogen, C₁₋₃alkyl or C₁₋₂alkoxyethyl).

Advantageously X^3 represents -O-, -S-, -SO-, -SO₂-, -NR¹⁷C(O)-, -NR²⁰SO₂- or -NR²¹- (wherein R¹⁷, R²⁰ and R²¹ each independently represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

15 Preferably X^3 represents -O-, -S-, -SO-, -SO₂- or -NR²¹- (wherein R²¹ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

More preferably X^3 represents -O- or -NR²¹- (wherein R²¹ represents hydrogen or C₁₋₂alkyl).

20 According to another aspect of the present invention X^3 represents -O-, -SO₂-, -NR²⁰SO₂- or -NR²¹- (wherein R²⁰ and R²¹ each independently represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Advantageously X^4 and X^5 which may be the same or different each represents -O-, -S-, -SO-, -SO₂- or -NR²⁷- (wherein R²⁷ represents hydrogen, C₁₋₃alkyl or C₁₋₂alkoxyethyl).

Preferably X^4 and X^5 which may be the same or different each represents -O-, -S- or -NR²⁷- (wherein R²⁷ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

25 More preferably X^4 and X^5 which may be the same or different each represents -O- or -NH-.

Advantageously X^6 represents -O-, -S- or -NR³⁸- (wherein R³⁸ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^6 represents -O- or -NR³⁸- (wherein R³⁸ represents hydrogen or C₁₋₂alkyl).

30 Advantageously X^7 represents -O-, -S- or -NR⁴³- (wherein R⁴³ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^7 represents -O- or -NR⁴³- (wherein R⁴³ represents hydrogen or C₁₋₂alkyl).

Advantageously X^8 represents -O-, -S- or -NR⁴⁸- (wherein R⁴⁸ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^8 represents -O- or -NR⁴⁸- (wherein R⁴⁸ represents hydrogen or C₁₋₂alkyl).

Advantageously X^9 represents -O-, -S- or -NR⁵³- (wherein R⁵³ represents hydrogen, C₁₋₂alkyl or C₁₋₂alkoxyethyl).

Preferably X^9 represents -O- or -NR⁵³- (wherein R⁵³ represents hydrogen or C₁₋₂alkyl).

According to another aspect of the present invention X^9 represents -O-, -CONR⁵⁰- or -NR⁵³- (wherein R⁵⁰ and R⁵³ each independently represents hydrogen or C₁₋₂alkyl).

Conveniently R²⁸ is pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl).

Advantageously R²⁸ is pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

In one embodiment of the present invention R²⁸ is pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

Particularly R²⁸ is pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy,

halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl and C₁₋₂alkylsulphonylC₁₋₃alkyl.

According to another aspect of the present invention, preferably R²⁸ is pyrrolidinyl, piperazinyl, piperidinyl, morpholino or thiomorpholino which group may bear 1 or 2
5 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl and C₁₋₂alkylsulphonylC₁₋₃alkyl.

Where R²⁹ is a 5-6-membered aromatic heterocyclic group, it preferably has 1 or 2 heteroatoms, selected from O, N and S, of which more preferably one is N, and may be substituted as hereinbefore defined.

10 R²⁹ is particularly a pyridone, phenyl, pyridyl, imidazolyl, thiazolyl, thienyl, triazolyl or pyridazinyl group which group may be substituted as hereinbefore defined, more particularly a pyridone, pyridyl, imidazolyl, thiazolyl or triazolyl group, especially a pyridone, pyridyl, imidazolyl or triazolyl group which group may be substituted as hereinbefore defined.

In one embodiment of the invention R²⁹ represents a pyridone, phenyl or 5-
15 6-membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which group may preferably carry up to 2 substituents, more preferably up to one substituent, selected from the group of substituents as hereinbefore defined.

In the definition of R²⁹, conveniently substituents are selected from halogeno, C₁₋₄alkyl, C₁₋₄alkoxy, cyano and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1
20 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl).

In the definition of R²⁹, more conveniently substituents are selected from chloro, fluoro, methyl, ethyl and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and
25 ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

According to another embodiment of the present invention in the definition of R²⁹, conveniently substituents are selected from halogeno, C₁₋₄alkyl, C₁₋₄alkoxy and cyano, more conveniently substituents are selected from chloro, fluoro, methyl and ethyl.

30 Advantageously R⁵⁴ and R⁵⁵ are each independently a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋

₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents
5 selected from C₁₋₃alkyl).

Preferably R⁵⁴ and R⁵⁵ are each selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring
10 D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl).

More preferably R⁵⁴ and R⁵⁵ are each selected from pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino which group may bear 1 or 2
15 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl and a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).
20

Particularly R⁵⁴ and R⁵⁵ are each selected from pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino which group may bear 1 or 2 substituents selected from a group $-(O-)_f(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).
25

More particularly R⁵⁴ and R⁵⁵ are each selected from pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino which group is unsubstituted.

Conveniently R² represents hydroxy, halogeno, cyano, nitro, trifluoromethyl, C₁₋₃alkyl, amino or R⁵X¹ - [wherein X¹ is as hereinbefore defined and R⁵ is selected from one of the
30 following twenty-two groups:

- 1) oxiranylC₁₋₄alkyl or C₁₋₅alkyl which may be unsubstituted or which may be substituted with one or more groups selected from fluoro, chloro and bromo, or C₂₋₅alkyl which may be unsubstituted or substituted with one or more groups selected from hydroxy and amino;
- 2) C₂₋₃alkylX²C(O)R¹¹ (wherein X² is as hereinbefore defined and R¹¹ represents C₁₋₃alkyl, -NR¹³R¹⁴ or -OR¹⁵ (wherein R¹³, R¹⁴ and R¹⁵ which may be the same or different are each C₁₋₄alkyl or C₁₋₂alkoxyethyl));
- 3) C₂₋₄alkylX³R¹⁶ (wherein X³ is as hereinbefore defined and R¹⁶ represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C₁₋₃alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group -(-O-)_f(C₁₋₄alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));
- 4) C₂₋₃alkylX⁴C₂₋₃alkylX⁵R²² (wherein X⁴ and X⁵ are as hereinbefore defined and R²² represents hydrogen or C₁₋₃alkyl);
- 5) R²⁸ (wherein R²⁸ is as defined hereinbefore);
- 6) C₁₋₅alkylR¹⁰⁷ (wherein R¹⁰⁷ is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group is linked to C₁₋₅alkyl through a carbon atom and which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group -(-O-)_f(C₁₋₄alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl)) or C₂₋₅alkylR¹⁰⁸ (wherein R¹⁰⁸ is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, of which one is N and the other may be selected independently from O, S and N, which heterocyclic group is linked to C₂₋₅alkyl

through a nitrogen atom and which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group $-(O-)_f(C_{1-4}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));

7) C₃₋₄alkenylR¹⁰⁹ (wherein R¹⁰⁹ represents R¹⁰⁷ or R¹⁰⁸ as defined hereinbefore);

10 8) C₃₋₄alkynylR¹⁰⁹ (wherein R¹⁰⁹ represents R¹⁰⁷ or R¹⁰⁸ as defined hereinbefore);

9) R²⁹ (wherein R²⁹ is as defined hereinbefore);

10) C₁₋₅alkylR²⁹ (wherein R²⁹ is as defined hereinbefore);

11) C₃₋₅alkenylR²⁹ (wherein R²⁹ is as defined hereinbefore);

12) C₃₋₅alkynylR²⁹ (wherein R²⁹ is as defined hereinbefore);

15 13) C₁₋₅alkylX⁶R²⁹ (wherein X⁶ and R²⁹ are as defined hereinbefore);

14) C₄₋₅alkenylX⁷R²⁹ (wherein X⁷ and R²⁹ are as defined hereinbefore);

15) C₄₋₅alkynylX⁸R²⁹ (wherein X⁸ and R²⁹ are as defined hereinbefore);

16) C₂₋₃alkylX⁹C₁₋₃alkylR²⁹ (wherein X⁹ and R²⁹ are as defined hereinbefore);

17) C₂₋₃alkylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);

20 18) C₂₋₅alkenyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;

19) C₂₋₅alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino,

25 aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;

20) C₂₋₅alkenylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);

21) C₂₋₅alkynylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore); and

22) C₁₋₃alkylR⁵⁴(C₁₋₃alkyl)_q(X⁹)_rR⁵⁵ (wherein X⁹, q, r, R⁵⁴ and R⁵⁵ are as defined hereinbefore);

and additionally wherein any C₁₋₅alkyl, C₂₋₅alkenyl or C₂₋₅alkynyl group in R⁵X¹ - may bear one or more substituents selected from hydroxy, halogeno and amino].

Advantageously R^2 represents hydroxy, halogeno, cyano, nitro, trifluoromethyl, C_{1-3} alkyl, amino or R^5X^1 - [wherein X^1 is as hereinbefore defined and R^5 is selected from one of the following twenty-two groups:

- 1) C_{1-4} alkyl which may be unsubstituted or which may be substituted with one or more groups selected from fluoro, chloro and bromo, or C_{2-3} alkyl which may be unsubstituted or substituted with one or more groups selected from hydroxy and amino;
- 2) C_{2-3} alkyl $X^2C(O)R^{11}$ (wherein X^2 is as hereinbefore defined and R^{11} represents $-NR^{13}R^{14}$ or $-OR^{15}$ (wherein R^{13} , R^{14} and R^{15} which may be the same or different are each C_{1-4} alkyl or C_{1-2} alkoxyethyl));
- 3) C_{2-4} alkyl X^3R^{16} (wherein X^3 is as hereinbefore defined and R^{16} is a group selected from C_{1-3} alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny and tetrahydropyranyl, which C_{1-3} alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C_{1-2} alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny or tetrahydropyranyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(-O-)_f(C_{1-3}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C_{1-3} alkyl));
- 4) C_{2-3} alkyl X^4C_{2-3} alkyl X^5R^{22} (wherein X^4 and X^5 are as hereinbefore defined and R^{22} represents hydrogen or C_{1-3} alkyl);
- 5) R^{28} (wherein R^{28} is as defined hereinbefore);
- 6) C_{1-4} alkyl R^{110} (wherein R^{110} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidin-1-yl, azetidiny, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to C_{1-4} alkyl through a carbon atom and which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(-O-)_f(C_{1-3}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C_{1-3} alkyl));

- O-)_f(C₁₋₃alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl)) or C₂₋₄alkylR¹¹¹ (wherein R¹¹¹ is a group selected from morpholino, thiomorpholino, azetidin-1-yl, pyrrolidin-1-yl, piperazin-1-yl and piperidino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group -(-
- 10 O-)_f(C₁₋₃alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl));
- 7) C₃₋₄alkenylR¹¹² (wherein R¹¹² represents R¹¹⁰ or R¹¹¹ as defined hereinbefore);
- 15 8) C₃₋₄alkynylR¹¹² (wherein R¹¹² represents R¹¹⁰ or R¹¹¹ as defined hereinbefore);
- 9) R²⁹ (wherein R²⁹ is as defined hereinbefore);
- 10) C₁₋₄alkylR²⁹ (wherein R²⁹ is as defined hereinbefore);
- 11) 1-R²⁹prop-1-en-3-yl or 1-R²⁹but-2-en-4-yl (wherein R²⁹ is as defined hereinbefore with the proviso that when R⁵ is 1-R²⁹prop-1-en-3-yl, R²⁹ is linked to the alkenyl group via a carbon
- 20 atom);
- 12) 1-R²⁹prop-1-yn-3-yl or 1-R²⁹but-2-yn-4-yl (wherein R²⁹ is as defined hereinbefore with the proviso that when R⁵ is 1-R²⁹prop-1-yn-3-yl, R²⁹ is linked to the alkynyl group via a carbon atom);
- 13) C₁₋₅alkylX⁶R²⁹ (wherein X⁶ and R²⁹ are as defined hereinbefore);
- 25 14) 1-(R²⁹X⁷)but-2-en-4-yl (wherein X⁷ and R²⁹ are as defined hereinbefore);
- 15) 1-(R²⁹X⁸)but-2-yn-4-yl (wherein X⁸ and R²⁹ are as defined hereinbefore);
- 16) C₂₋₃alkylX⁹C₁₋₃alkylR²⁹ (wherein X⁹ and R²⁹ are as defined hereinbefore);
- 17) C₂₋₃alkylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);
- 18) C₂₋₅alkenyl which may be unsubstituted or which may be substituted with one or more
- 30 fluorine atoms or with one or two groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;

- 19) $C_{2,5}$ alkynyl which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, fluoro, amino, $C_{1,4}$ alkylamino, N,N -di($C_{1,4}$ alkyl)amino, aminosulphonyl, N - $C_{1,4}$ alkylaminosulphonyl and N,N -di($C_{1,4}$ alkyl)aminosulphonyl;
- 5 20) $C_{2,4}$ alkenyl $X^9C_{1,3}$ alkyl R^{28} (wherein X^9 and R^{28} are as defined hereinbefore);
- 21) $C_{2,4}$ alkynyl $X^9C_{1,3}$ alkyl R^{28} (wherein X^9 and R^{28} are as defined hereinbefore); and
- 22) $C_{1,3}$ alkyl $R^{54}(C_{1,3}$ alkyl) $_q(X^9)_rR^{55}$ (wherein X^9 , q , r , R^{54} and R^{55} are as defined hereinbefore); and additionally wherein any $C_{1,5}$ alkyl, $C_{2,5}$ alkenyl or $C_{2,5}$ alkynyl group in R^5X^1 - may bear one or more substituents selected from hydroxy, halogeno and amino].
- 10 Preferably R^2 represents hydroxy, halogeno, nitro, trifluoromethyl, $C_{1,3}$ alkyl, cyano, amino or R^5X^1 - [wherein X^1 is as hereinbefore defined and R^5 is selected from one of the following twenty groups:
- 1) $C_{1,3}$ alkyl which may be unsubstituted or which may be substituted with one or more groups selected from fluoro, chloro and bromo, or $C_{2,3}$ alkyl which may be unsubstituted or
- 15 substituted with one or more groups selected from hydroxy and amino;
- 2) 2-(3,3-dimethylureido)ethyl, 3-(3,3-dimethylureido)propyl, 2-(3-methylureido)ethyl, 3-(3-methylureido)propyl, 2-ureidoethyl, 3-ureidopropyl, 2-(N,N -dimethylcarbamoyloxy)ethyl, 3-(N,N -dimethylcarbamoyloxy)propyl, 2-(N -methylcarbamoyloxy)ethyl, 3-(N -methylcarbamoyloxy)propyl, 2-(carbamoyloxy)ethyl, 3-(carbamoyloxy)propyl, or 2-(N -methyl- N -(butoxycarbonyl)amino)ethyl;
- 20 3) $C_{2,3}$ alkyl X^3R^{16} (wherein X^3 is as hereinbefore defined and R^{16} is a group selected from $C_{1,3}$ alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl, piperidinyl, piperazinyl, azetidiny, imidazolidinyl and tetrahydropyranyl which group is linked to X^3 through a carbon atom and which $C_{1,3}$ alkyl group may bear 1 or 2 substituents selected from hydroxy, halogeno and $C_{1,2}$ alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl, piperidinyl, piperazinyl, azetidiny, imidazolidinyl or tetrahydropyranyl group may bear one substituent selected from oxo, hydroxy, halogeno, cyano, $C_{1,2}$ cyanoalkyl, $C_{1,2}$ alkyl, $C_{1,2}$ hydroxyalkyl, $C_{1,2}$ alkoxy, $C_{1,2}$ alkoxy $C_{1,3}$ alkyl, $C_{1,2}$ alkylsulphonyl $C_{1,3}$ alkyl, $C_{1,2}$ alkoxycarbonyl, $C_{1,3}$ alkylamino, di($C_{1,3}$ alkyl)amino, $C_{1,3}$ alkylamino $C_{1,3}$ alkyl, di($C_{1,3}$ alkyl)amino $C_{1,3}$ alkyl, $C_{1,3}$ alkylamino $C_{1,3}$ alkoxy, di($C_{1,3}$ alkyl)amino $C_{1,3}$ alkoxy and a group $-(O-)_f(C_{1,3}alkyl)_g$ ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino));
- 30

- 4) $C_{2-3}alkylX^4C_{2-3}alkylX^5R^{22}$ (wherein X^4 and X^5 are as hereinbefore defined and R^{22} represents hydrogen or $C_{1-2}alkyl$);
- 5) R^{28} (wherein R^{28} is as defined hereinbefore);
- 6) $C_{1-3}alkylR^{110}$ (wherein R^{110} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, imidazolidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to $C_{1-3}alkyl$ through a carbon atom and which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, $C_{1-2}cyanoalkyl$, $C_{1-2}alkyl$, $C_{1-2}hydroxyalkyl$, $C_{1-2}alkoxy$, $C_{1-2}alkoxyC_{1-3}alkyl$, $C_{1-2}alkylsulphonylC_{1-3}alkyl$, $C_{1-2}alkoxycarbonyl$, $C_{1-3}alkylamino$, $di(C_{1-3}alkyl)amino$, $C_{1-3}alkylaminoC_{1-3}alkyl$, $di(C_{1-3}alkyl)aminoC_{1-3}alkyl$, $C_{1-3}alkylaminoC_{1-3}alkoxy$, $di(C_{1-3}alkyl)aminoC_{1-3}alkoxy$ and a group $-(O-)_f(C_{1-3}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino)) or $C_{2-3}alkylR^{111}$ (wherein R^{111} is a group selected from morpholino, thiomorpholino, azetidin-1-yl, pyrrolidin-1-yl, piperazin-1-yl and piperidino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, $C_{1-2}cyanoalkyl$, $C_{1-2}alkyl$, $C_{1-2}hydroxyalkyl$, $C_{1-2}alkoxy$, $C_{1-2}alkoxyC_{1-3}alkyl$, $C_{1-2}alkylsulphonylC_{1-3}alkyl$, $C_{1-2}alkoxycarbonyl$, $C_{1-3}alkylamino$, $di(C_{1-3}alkyl)amino$, $C_{1-3}alkylaminoC_{1-3}alkyl$, $di(C_{1-3}alkyl)aminoC_{1-3}alkyl$, $C_{1-3}alkylaminoC_{1-3}alkoxy$, $di(C_{1-3}alkyl)aminoC_{1-3}alkoxy$ and a group $-(O-)_f(C_{1-3}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino));
- 7) R^{29} (wherein R^{29} is as defined hereinbefore);
- 8) $C_{1-4}alkylR^{29}$ (wherein R^{29} is as defined hereinbefore);
- 9) $1-R^{29}but-2-en-4-yl$ (wherein R^{29} is as defined hereinbefore);
- 10) $1-R^{29}but-2-yn-4-yl$ (wherein R^{29} is as defined hereinbefore);
- 11) $C_{1-3}alkylX^6R^{29}$ (wherein X^6 and R^{29} are as defined hereinbefore);
- 12) $1-(R^{29}X^7)but-2-en-4-yl$ (wherein X^7 and R^{29} are as defined hereinbefore);
- 13) $1-(R^{29}X^8)but-2-yn-4-yl$ (wherein X^8 and R^{29} are as defined hereinbefore);
- 14) $C_{2-3}alkylX^9C_{1-3}alkylR^{29}$ (wherein X^9 and R^{29} are as defined hereinbefore);
- 15) $C_{2-3}alkylX^9C_{1-3}alkylR^{28}$ (wherein X^9 and R^{28} are as defined hereinbefore);
- 16) $C_{2-3}alkenyl$ which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, fluoro, amino, $C_{1-2}alkyl$, $C_{1-2}hydroxyalkyl$, $C_{1-2}alkoxy$, $C_{1-2}alkoxyC_{1-3}alkyl$, $C_{1-2}alkylsulphonylC_{1-3}alkyl$, $C_{1-2}alkoxycarbonyl$, $C_{1-3}alkylamino$, $di(C_{1-3}alkyl)amino$, $C_{1-3}alkylaminoC_{1-3}alkyl$, $di(C_{1-3}alkyl)aminoC_{1-3}alkyl$, $C_{1-3}alkylaminoC_{1-3}alkoxy$, $di(C_{1-3}alkyl)aminoC_{1-3}alkoxy$ and a group $-(O-)_f(C_{1-3}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino));

4alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;

17) C₂₋₃alkynyl which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, fluoro, amino, C₁₋

4alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;

18) C₂₋₃alkenylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);

19) C₂₋₃alkynylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore); and

20) C₁₋₃alkylR⁵⁴(C₁₋₃alkyl)_q(X⁹)_rR⁵⁵ (wherein X⁹, q, r, R⁵⁴ and R⁵⁵ are as defined hereinbefore);

and additionally wherein any C₁₋₃alkyl, C₂₋₃alkenyl or C₂₋₃alkynyl group in R⁵X¹- may bear one or more substituents selected from hydroxy, halogeno and amino].

More preferably R² represents hydroxy, C₁₋₃alkyl, amino or R⁵X¹- [wherein X¹ is as hereinbefore defined and R⁵ represents methyl, ethyl, benzyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-

(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(ethylsulphinyl)ethyl, 2-(ethylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl, 2-(methylamino)ethyl, 3-(methylamino)propyl, 2-(ethylamino)ethyl, 3-(ethylamino)propyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-(N,N-diethylamino)ethyl, 3-(N,N-diethylamino)propyl, 2-(N-methyl-N-

methylsulphonylamino)ethyl, 3-(N-methyl-N-methylsulphonylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, 2-(piperidin-2-yl)ethyl, 3-(piperidin-2-yl)propyl, (1-methylpiperidin-3-yl)methyl, (1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-

(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonyl)ethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonyl)ethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonyl)ethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonyl)ethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-yl)methyl, 1-isopropylpiperidin-3-yl)methyl, 1-isopropylpiperidin-4-yl)methyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperidin-4-yloxy)ethyl, 3-(piperidin-4-yloxy)propyl, 2-(1-(cyanomethyl)piperidin-4-yloxy)ethyl, 3-(1-(cyanomethyl)piperidin-4-yloxy)propyl, 2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethyl, 3-(1-(2-cyanoethyl)piperidin-4-yloxy)propyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, 5(R)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (5S)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-(2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-yl)methyl, 2-acetamidothiazol-4-yl)methyl, 1-methylimidazol-2-yl)methyl, 2-(imidazol-1-yl)ethyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(2-ethylimidazol-1-yl)ethyl, 3-(2-methylimidazol-1-yl)propyl, 3-(2-ethylimidazol-1-yl)propyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridyl)methyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-(2-oxo-imidazolidin-1-yl)ethyl, 3-(2-oxo-imidazolidin-1-yl)propyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 3-(ethylsulphinyl)propyl, 3-(ethylsulphonyl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-(N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-(N-(3-

morpholinopropylsulphonyl)-N-methylamino)ethyl, 2-((N-methyl-N-4-pyridyl)amino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-(2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl, 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yl, 1-(2-pyrrolidinylethyl)piperidin-4-ylmethyl, 1-(3-pyrrolidinylpropyl)piperidin-4-ylmethyl, 1-(2-piperidinylethyl)piperidin-4-ylmethyl, 1-(3-piperidinylpropyl)piperidin-4-ylmethyl, 1-(2-morpholinoethyl)piperidin-4-ylmethyl, 1-(3-morpholinopropyl)piperidin-4-ylmethyl, 1-(2-thiomorpholinoethyl)piperidin-4-ylmethyl, 1-(3-thiomorpholinopropyl)piperidin-4-ylmethyl, 1-(2-azetidinyethyl)piperidin-4-ylmethyl or 1-(3-azetidinyethyl)piperidin-4-ylmethyl, 3-morpholino-2-hydroxypropyl, (2*R*)-3-morpholino-2-hydroxypropyl, (2*S*)-3-morpholino-2-hydroxypropyl, 3-piperidino-2-hydroxypropyl, (2*R*)-3-piperidino-2-hydroxypropyl, (2*S*)-3-piperidino-2-hydroxypropyl, 3-pyrrolidin-1-yl-2-hydroxypropyl, (2*R*)-3-pyrrolidin-1-yl-2-hydroxypropyl, (2*S*)-3-pyrrolidin-1-yl-2-hydroxypropyl, 3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*R*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*S*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, 3-(N,N-diethylamino)-2-hydroxypropyl, (2*R*)-3-(N,N-diethylamino)-2-hydroxypropyl, (2*S*)-3-(N,N-diethylamino)-2-hydroxypropyl, 3-(isopropylamino)-2-hydroxypropyl, (2*R*)-3-(isopropylamino)-2-hydroxypropyl, (2*S*)-3-(isopropylamino)-2-hydroxypropyl, 3-(N,N-diisopropylamino)-2-hydroxypropyl, (2*R*)-3-(N,N-diisopropylamino)-2-hydroxypropyl or (2*S*)-3-(N,N-diisopropylamino)-2-hydroxypropyl].

Particularly R² represents C₁₋₃alkyl, amino or R⁵X¹- [wherein X¹ is as hereinbefore defined and R⁵ represents ethyl, benzyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(ethylsulphinyl)ethyl, 2-(ethylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoylethyl, 2-(methylamino)ethyl, 3-(methylamino)propyl, 2-(ethylamino)ethyl, 3-(ethylamino)propyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-(N,N-diethylamino)ethyl, 3-(N,N-diethylamino)propyl, 2-(N-methyl-N-methylsulphonylamino)ethyl, 3-(N-methyl-N-methylsulphonylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-

methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, 2-(piperidin-2-yl)ethyl, 3-(piperidin-2-yl)propyl, (1-methylpiperidin-3-yl)methyl, (1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonyl)ethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonyl)ethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonyl)ethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonyl)ethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-ylmethyl, 1-isopropylpiperidin-3-ylmethyl, 1-isopropylpiperidin-4-ylmethyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperidin-4-yloxy)ethyl, 3-(piperidin-4-yloxy)propyl, 2-(1-(cyanomethyl)piperidin-4-yloxy)ethyl, 3-(1-(cyanomethyl)piperidin-4-yloxy)propyl, 2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethyl, 3-(1-(2-cyanoethyl)piperidin-4-yloxy)propyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, 5(R)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (5S)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-(2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(2-ethylimidazol-1-yl)ethyl, 3-(2-methylimidazol-1-yl)propyl, 3-(2-ethylimidazol-1-yl)propyl, 2-

(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-(2-oxo-imidazolidin-1-yl)ethyl, 3-(2-oxo-imidazolidin-1-yl)propyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 3-(ethylsulphinyl)propyl, 3-(ethylsulphonyl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-((N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-((N-methyl-N-4-pyridyl)amino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-(2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl, 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yl, 1-(2-pyrrolidinylethyl)piperidin-4-ylmethyl, 1-(3-pyrrolidinylpropyl)piperidin-4-ylmethyl, 1-(2-piperidinylethyl)piperidin-4-ylmethyl, 1-(3-piperidinylpropyl)piperidin-4-ylmethyl, 1-(2-morpholinoethyl)piperidin-4-ylmethyl, 1-(3-morpholinopropyl)piperidin-4-ylmethyl, 1-(2-thiomorpholinoethyl)piperidin-4-ylmethyl, 1-(3-thiomorpholinopropyl)piperidin-4-ylmethyl, 1-(2-azetidinyethyl)piperidin-4-ylmethyl or 1-(3-azetidylpropyl)piperidin-4-ylmethyl, 3-morpholino-2-hydroxypropyl, (2*R*)-3-morpholino-2-hydroxypropyl, (2*S*)-3-morpholino-2-hydroxypropyl, 3-piperidino-2-hydroxypropyl, (2*R*)-3-piperidino-2-hydroxypropyl, (2*S*)-3-piperidino-2-hydroxypropyl, 3-pyrrolidin-1-yl-2-hydroxypropyl, (2*R*)-3-pyrrolidin-1-yl-2-hydroxypropyl, (2*S*)-3-pyrrolidin-1-yl-2-hydroxypropyl, 3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*R*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*S*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, 3-((N,N-diethylamino)-2-hydroxypropyl, (2*R*)-3-((N,N-diethylamino)-2-hydroxypropyl, (2*S*)-3-((N,N-diethylamino)-2-hydroxypropyl, 3-(isopropylamino)-2-hydroxypropyl, (2*R*)-3-(isopropylamino)-2-hydroxypropyl, (2*S*)-3-(isopropylamino)-2-hydroxypropyl, 3-((N,N-diisopropylamino)-2-hydroxypropyl, (2*R*)-3-((N,N-diisopropylamino)-2-hydroxypropyl or (2*S*)-3-((N,N-diisopropylamino)-2-hydroxypropyl].

More particularly R^2 represents C_{1-3} alkyl, amino or R^5X^1 - [wherein X^1 is as hereinbefore defined and R^5 represents ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-

hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(ethylsulphinyl)ethyl, 2-(ethylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoylethyl, 2-(methylamino)ethyl, 3-(methylamino)propyl, 2-(ethylamino)ethyl, 3-(ethylamino)propyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-(N,N-diethylamino)ethyl, 3-(N,N-diethylamino)propyl, 2-(N-methyl-N-methylsulphonylamino)ethyl, 3-(N-methyl-N-methylsulphonylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, 2-(piperidin-2-yl)ethyl, 3-(piperidin-2-yl)propyl, (1-methylpiperidin-3-yl)methyl, (1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonylethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonylethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonylethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonylethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonylethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonylethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-ylmethyl, 1-isopropylpiperidin-3-ylmethyl, 1-isopropylpiperidin-4-ylmethyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperidin-4-yloxy)ethyl, 3-(piperidin-4-yloxy)propyl, 2-(1-(cyanomethyl)piperidin-4-yloxy)ethyl, 3-(1-(cyanomethyl)piperidin-4-yloxy)propyl, 2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethyl, 3-(1-(2-cyanoethyl)piperidin-4-yloxy)propyl, 2-

(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2*H*-pyrrolidin-5-yl)methyl, 5(*R*)-(2-oxo-tetrahydro-2*H*-pyrrolidin-5-yl)methyl, (5*S*)-(2-oxo-tetrahydro-2*H*-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-
5 (2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-(2-oxo-imidazolidin-1-yl)ethyl, 3-(2-oxo-imidazolidin-1-yl)propyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 3-(ethylsulphinyl)propyl, 3-(ethylsulphonyl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl,
15 morpholino, 2-(N-(3-morpholinopropylsulphonyl)-N-methylamino)ethyl, 2-(N-methyl-N-4-pyridylamino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-(2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl, 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yl, 1-(2-pyrrolidinylethyl)piperidin-4-ylmethyl, 1-(3-pyrrolidinylpropyl)piperidin-4-ylmethyl, 1-(2-piperidinylethyl)piperidin-4-ylmethyl, 1-(3-piperidinylpropyl)piperidin-4-ylmethyl, 1-(2-morpholinoethyl)piperidin-4-ylmethyl, 1-(3-morpholinopropyl)piperidin-4-ylmethyl, 1-(2-thiomorpholinoethyl)piperidin-4-ylmethyl, 1-(3-thiomorpholinopropyl)piperidin-4-ylmethyl, 1-(2-azetidinyethyl)piperidin-4-ylmethyl or
25 1-(3-azetidinypropyl)piperidin-4-ylmethyl, 3-morpholino-2-hydroxypropyl, (2*R*)-3-morpholino-2-hydroxypropyl, (2*S*)-3-morpholino-2-hydroxypropyl, 3-piperidino-2-hydroxypropyl, (2*R*)-3-piperidino-2-hydroxypropyl, (2*S*)-3-piperidino-2-hydroxypropyl, 3-pyrrolidin-1-yl-2-hydroxypropyl, (2*R*)-3-pyrrolidin-1-yl-2-hydroxypropyl, (2*S*)-3-pyrrolidin-1-yl-2-hydroxypropyl, 3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*R*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, (2*S*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropyl, 3-(N,N-diethylamino)-2-hydroxypropyl, (2*R*)-3-(N,N-diethylamino)-2-hydroxypropyl, (2*S*)-3-(N,N-diethylamino)-2-hydroxypropyl, 3-(isopropylamino)-2-hydroxypropyl, (2*R*)-3-

(isopropylamino)-2-hydroxypropyl, (2*S*)-3-(isopropylamino)-2-hydroxypropyl, 3-(N,N-diisopropylamino)-2-hydroxypropyl, (2*R*)-3-(N,N-diisopropylamino)-2-hydroxypropyl or (2*S*)-3-(N,N-diisopropylamino)-2-hydroxypropyl].

In another aspect R² represents ethoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy, 2-hydroxyethoxy, 3-hydroxypropoxy, 2-methoxyethoxy, 3-methoxypropoxy, 2-(methylsulphinyl)ethoxy, 2-(methylsulphonyl)ethoxy, 2-(ethylsulphinyl)ethoxy, 2-(ethylsulphonyl)ethoxy, 2-(N,N-dimethylsulphamoyl)ethoxy, 2-(N-methylsulphamoyl)ethoxy, 2-sulphamoylethoxy, 2-(methylamino)ethoxy, 3-(methylamino)propoxy, 2-(ethylamino)ethoxy, 3-(ethylamino)propoxy, 2-(N,N-dimethylamino)ethoxy, 3-(N,N-dimethylamino)propoxy, 2-(N,N-diethylamino)ethoxy, 3-(N,N-diethylamino)propoxy, 2-(N-methyl-N-methylsulphonylamino)ethoxy, 3-(N-methyl-N-methylsulphonylamino)propoxy, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-(methylpiperidino)ethoxy, 3-(methylpiperidino)propoxy, 2-(ethylpiperidino)ethoxy, 3-(ethylpiperidino)propoxy, 2-((2-methoxyethyl)piperidino)ethoxy, 3-((2-methoxyethyl)piperidino)propoxy, 2-((2-methylsulphonyl)ethylpiperidino)ethoxy, 3-((2-methylsulphonyl)ethylpiperidino)propoxy, piperidin-3-ylmethoxy, piperidin-4-ylmethoxy, 2-(piperidin-3-yl)ethoxy, 2-(piperidin-4-yl)ethoxy, 3-(piperidin-3-yl)propoxy, 3-(piperidin-4-yl)propoxy, 2-(piperidin-2-yl)ethoxy, 3-(piperidin-2-yl)propoxy, (1-methylpiperidin-3-yl)ethoxy, (1-methylpiperidin-4-yl)ethoxy, (1-cyanomethylpiperidin-3-yl)ethoxy, (1-cyanomethylpiperidin-4-yl)ethoxy, 2-(methylpiperidin-3-yl)ethoxy, 2-(methylpiperidin-4-yl)ethoxy, 2-(1-cyanomethylpiperidin-3-yl)ethoxy, 2-(1-cyanomethylpiperidin-4-yl)ethoxy, 3-(methylpiperidin-3-yl)propoxy, 3-(methylpiperidin-4-yl)propoxy, 3-(1-cyanomethylpiperidin-3-yl)propoxy, 3-(1-cyanomethylpiperidin-4-yl)propoxy, 2-(ethylpiperidin-3-yl)ethoxy, 2-(ethylpiperidin-4-yl)ethoxy, 3-(ethylpiperidin-3-yl)propoxy, 3-(ethylpiperidin-4-yl)propoxy, ((2-methoxyethyl)piperidin-3-yl)ethoxy, ((2-methoxyethyl)piperidin-4-yl)ethoxy, 2-((2-methoxyethyl)piperidin-3-yl)ethoxy, 2-((2-methoxyethyl)piperidin-4-yl)ethoxy, 3-((2-methoxyethyl)piperidin-3-yl)propoxy, 3-((2-methoxyethyl)piperidin-4-yl)propoxy, (1-(2-methylsulphonylethyl)piperidin-3-yl)ethoxy, (1-(2-methylsulphonylethyl)piperidin-4-yl)ethoxy, 2-((2-methylsulphonylethyl)piperidin-3-yl)ethoxy, 2-((2-methylsulphonylethyl)piperidin-4-yl)ethoxy, 3-((2-methylsulphonylethyl)piperidin-3-yl)propoxy, 3-((2-methylsulphonylethyl)piperidin-4-yl)propoxy, 1-isopropylpiperidin-2-ylmethoxy, 1-isopropylpiperidin-3-ylmethoxy, 1-isopropylpiperidin-4-ylmethoxy, 2-(1-

isopropylpiperidin-2-yl)ethoxy, 2-(1-isopropylpiperidin-3-yl)ethoxy, 2-(1-isopropylpiperidin-4-yl)ethoxy, 3-(1-isopropylpiperidin-2-yl)propoxy, 3-(1-isopropylpiperidin-3-yl)propoxy, 3-(1-isopropylpiperidin-4-yl)propoxy, 2-(piperidin-4-yloxy)ethoxy, 3-(piperidin-4-yloxy)propoxy, 2-(1-(cyanomethyl)piperidin-4-yloxy)ethoxy, 3-(1-(cyanomethyl)piperidin-4-yloxy)propoxy, 2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethoxy, 3-(1-(2-cyanoethyl)piperidin-4-yloxy)propoxy, 2-(piperazin-1-yl)ethoxy, 3-(piperazin-1-yl)propoxy, (pyrrolidin-2-yl)methoxy, 2-(pyrrolidin-1-yl)ethoxy, 3-(pyrrolidin-1-yl)propoxy, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methoxy, 5(R)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methoxy, (5S)-(2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methoxy, (1,3-dioxolan-2-yl)methoxy, 2-(1,3-dioxolan-2-yl)ethoxy, 2-(2-methoxyethylamino)ethoxy, 2-(N-(2-methoxyethyl)-N-methylamino)ethoxy, 2-(2-hydroxyethylamino)ethoxy, 3-(2-methoxyethylamino)propoxy, 3-(N-(2-methoxyethyl)-N-methylamino)propoxy, 3-(2-hydroxyethylamino)propoxy, 2-(1,2,3-triazol-1-yl)ethoxy, 2-(1,2,3-triazol-2-yl)ethoxy, 2-(1,2,4-triazol-1-yl)ethoxy, 2-(1,2,4-triazol-4-yl)ethoxy, 4-pyridylmethoxy, 2-(4-pyridyl)ethoxy, 3-(4-pyridyl)propoxy, 2-(4-pyridyloxy)ethoxy, 2-(4-pyridylamino)ethoxy, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethoxy, 2-(2-oxo-imidazolidin-1-yl)ethoxy, 3-(2-oxo-imidazolidin-1-yl)propoxy, 2-thiomorpholinoethoxy, 3-thiomorpholinopropoxy, 2-(1,1-dioxothiomorpholino)ethoxy, 3-(1,1-dioxothiomorpholino)propoxy, 2-(2-methoxyethoxy)ethoxy, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(4-methylpiperazin-1-yl)propoxy, 3-(methylsulphonyl)propoxy, 3-(methylsulphonyl)propoxy, 3-(ethylsulphonyl)propoxy, 2-(5-methyl-1,2,4-triazol-1-yl)ethoxy, 2-(N-(3-morpholinopropylsulphonyl)-N-methylamino)ethoxy, 2-(N-methyl-N-4-pyridylamino)ethoxy, 3-(4-oxidomorpholino)propoxy, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethoxy, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propoxy, 2-(2-morpholinoethoxy)ethoxy, 3-(2-morpholinoethoxy)propoxy, 2-(tetrahydropyran-4-yloxy)ethoxy, 3-(tetrahydropyran-4-yloxy)propoxy, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl, 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yloxy, 1-(2-pyrrolidinylethyl)piperidin-4-ylmethoxy, 1-(3-pyrrolidinylpropyl)piperidin-4-ylmethoxy, 1-(2-piperidinylethyl)piperidin-4-ylmethoxy, 1-(3-piperidinylpropyl)piperidin-4-ylmethoxy, 1-(2-morpholinoethyl)piperidin-4-ylmethoxy, 1-(3-morpholinopropyl)piperidin-4-ylmethoxy, 1-(2-thiomorpholinoethyl)piperidin-4-ylmethoxy, 1-(3-thiomorpholinopropyl)piperidin-4-ylmethoxy, 1-(2-azetidinyethyl)piperidin-4-ylmethoxy or 1-(3-azetidinyethyl)piperidin-4-ylmethoxy, 3-morpholino-2-hydroxypropoxy,

(2*R*)-3-morpholino-2-hydroxypropoxy, (2*S*)-3-morpholino-2-hydroxypropoxy, 3-piperidino-2-hydroxypropoxy, (2*R*)-3-piperidino-2-hydroxypropoxy, (2*S*)-3-piperidino-2-hydroxypropoxy, 3-pyrrolidin-1-yl-2-hydroxypropoxy, (2*R*)-3-pyrrolidin-1-yl-2-hydroxypropoxy, (2*S*)-3-pyrrolidin-1-yl-2-hydroxypropoxy, 3-(1-methylpiperazin-4-yl)-2-hydroxypropoxy, (2*R*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropoxy, (2*S*)-3-(1-methylpiperazin-4-yl)-2-hydroxypropoxy, 3-(N,N-diethylamino)-2-hydroxypropoxy, (2*R*)-3-(N,N-diethylamino)-2-hydroxypropoxy, (2*S*)-3-(N,N-diethylamino)-2-hydroxypropoxy, 3-(isopropylamino)-2-hydroxypropoxy, (2*R*)-3-(isopropylamino)-2-hydroxypropoxy, (2*S*)-3-(isopropylamino)-2-hydroxypropoxy, 3-(N,N-diisopropylamino)-2-hydroxypropoxy, (2*R*)-3-(N,N-diisopropylamino)-2-hydroxypropoxy or (2*S*)-3-(N,N-diisopropylamino)-2-hydroxypropoxy.

According to another aspect of the present invention conveniently R² represents hydroxy, halogeno, nitro, trifluoromethyl, C₁₋₃alkyl, cyano, amino or R⁵X¹- [wherein X¹ is as hereinbefore defined and R⁵ is selected from one of the following twenty-one groups:

- 1) C₁₋₅alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C₂₋₅alkyl which may be unsubstituted or substituted with one or more groups selected from hydroxy and amino;
- 2) C₂₋₃alkylX²C(O)R¹¹ (wherein X² is as hereinbefore defined and R¹¹ represents C₁₋₃alkyl, -NR¹³R¹⁴ or -OR¹⁵ (wherein R¹³, R¹⁴ and R¹⁵ which may be the same or different are each C₁₋₂alkyl or C₁₋₂alkoxyethyl));
- 3) C₂₋₄alkylX³R¹⁶ (wherein X³ is as hereinbefore defined and R¹⁶ represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C₁₋₃alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);
- 4) C₂₋₃alkylX⁴C₂₋₃alkylX⁵R²² (wherein X⁴ and X⁵ are as hereinbefore defined and R²² represents hydrogen or C₁₋₃alkyl);
- 5) C₁₋₅alkylR¹²⁹ (wherein R¹²⁹ is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group is linked to C₁₋₅alkyl through a carbon atom and which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋

- alkoxy, C₁₋₄alkoxyC₁₋₄alkyl and C₁₋₄alkylsulphonylC₁₋₄alkyl) or C₂₋₅alkylR¹³⁰ (wherein R¹³⁰ is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms of which one is N and the other is selected independently from O, S and N, which heterocyclic group is linked to C₂₋₅alkyl through a nitrogen atom and which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl and C₁₋₄alkylsulphonylC₁₋₄alkyl);
- 6) C₃₋₄alkenylR¹³¹ (wherein R¹³¹ represents R¹²⁹ or R¹³⁰ as defined hereinbefore);
- 7) C₃₋₄alkynylR¹³¹ (wherein R¹³¹ represents R¹²⁹ or R¹³⁰ as defined hereinbefore);
- 8) R²⁹ (wherein R²⁹ is as defined hereinbefore);
- 9) C₁₋₅alkylR²⁹ (wherein R²⁹ is as defined hereinbefore);
- 10) C₃₋₅alkenylR²⁹ (wherein R²⁹ is as defined hereinbefore);
- 11) C₃₋₅alkynylR²⁹ (wherein R²⁹ is as defined hereinbefore);
- 12) C₁₋₅alkylX⁶R²⁹ (wherein X⁶ and R²⁹ are as defined hereinbefore);
- 13) C₄₋₅alkenylX⁷R²⁹ (wherein X⁷ and R²⁹ are as defined hereinbefore);
- 14) C₄₋₅alkynylX⁸R²⁹ (wherein X⁸ and R²⁹ are as defined hereinbefore);
- 15) C₂₋₃alkylX⁹C₁₋₂alkylR²⁹ (wherein X⁹ and R²⁹ are as defined hereinbefore);
- 16) R²⁸ (wherein R²⁸ is as defined hereinbefore);
- 17) C₂₋₃alkylX⁹C₁₋₂alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);
- 18) C₂₋₅alkenyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;
- 19) C₂₋₅alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;
- 20) C₂₋₅alkenylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore); and
- 21) C₂₋₅alkynylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore)].

According to another aspect of the present invention advantageously R² represents hydroxy, halogeno, nitro, trifluoromethyl, C₁₋₃alkyl, cyano, amino or R⁵X¹ - [wherein X¹ is as hereinbefore defined and R⁵ is selected from one of the following twenty-one groups:

- 1) C₁₋₄alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C₂₋₄alkyl which may be unsubstituted or substituted with 1 or 2 groups selected from hydroxy and amino;

- 2) $C_{2-3}alkylX^2C(O)R^{11}$ (wherein X^2 is as hereinbefore defined and R^{11} represents $-NR^{13}R^{14}$ or $-OR^{15}$ (wherein R^{13} , R^{14} and R^{15} which may be the same or different are each $C_{1-2}alkyl$ or $C_{1-2}alkoxyethyl$));
- 3) $C_{2-4}alkylX^3R^{16}$ (wherein X^3 is as hereinbefore defined and R^{16} is a group selected from $C_{1-3}alkyl$, cyclopentyl, cyclohexyl, pyrrolidinyl, piperidinyl and tetrahydropyranyl which group is linked to X^3 through a carbon atom and which $C_{1-3}alkyl$ group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and $C_{1-2}alkoxy$ and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1-2}alkyl$, $C_{1-2}hydroxyalkyl$ and $C_{1-2}alkoxy$);
- 4) $C_{2-3}alkylX^4C_{2-3}alkylX^5R^{22}$ (wherein X^4 and X^5 are as hereinbefore defined and R^{22} represents hydrogen or $C_{1-3}alkyl$);
- 5) $C_{1-4}alkylR^{132}$ (wherein R^{132} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to $C_{1-4}alkyl$ through a carbon atom and which group may carry 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, $C_{1-3}cyanoalkyl$, $C_{1-3}alkyl$, $C_{1-3}hydroxyalkyl$, $C_{1-3}alkoxy$, $C_{1-2}alkoxyC_{1-3}alkyl$ and $C_{1-2}alkylsulphonylC_{1-3}alkyl$) or $C_{2-4}alkylR^{133}$ (wherein R^{133} is a group selected from morpholino, thiomorpholino, pyrrolidin-1-yl, piperazin-1-yl and piperidino which group may carry 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, $C_{1-3}cyanoalkyl$, $C_{1-3}alkyl$, $C_{1-3}hydroxyalkyl$, $C_{1-3}alkoxy$, $C_{1-2}alkoxyC_{1-3}alkyl$ and $C_{1-2}alkylsulphonylC_{1-3}alkyl$);
- 6) $C_{3-4}alkenylR^{134}$ (wherein R^{134} represents R^{132} or R^{133} as defined hereinbefore);
- 7) $C_{3-4}alkynylR^{134}$ (wherein R^{134} represents R^{132} or R^{133} as defined hereinbefore);
- 8) R^{29} (wherein R^{29} is as defined hereinbefore);
- 9) $C_{1-4}alkylR^{29}$ (wherein R^{29} is as defined hereinbefore);
- 10) 1- R^{29} prop-1-en-3-yl or 1- R^{29} but-2-en-4-yl (wherein R^{29} is as defined hereinbefore with the proviso that when R^5 is 1- R^{29} prop-1-en-3-yl, R^{29} is linked to the alkenyl group via a carbon atom);
- 11) 1- R^{29} prop-1-yn-3-yl or 1- R^{29} but-2-yn-4-yl (wherein R^{29} is as defined hereinbefore with the proviso that when R^5 is 1- R^{29} prop-1-yn-3-yl, R^{29} is linked to the alkynyl group via a carbon atom);
- 12) $C_{1-3}alkylX^6R^{29}$ (wherein X^6 and R^{29} are as defined hereinbefore);
- 13) 1-($R^{29}X^7$)but-2-en-4-yl (wherein X^7 and R^{29} are as defined hereinbefore);

14) 1-(R²⁹X⁸)but-2-yn-4-yl (wherein X⁸ and R²⁹ are as defined hereinbefore);

15) C_{2,3}alkylX⁹C_{1,2}alkylR²⁹ (wherein X⁹ and R²⁹ are as defined hereinbefore);

16) R²⁸ (wherein R²⁸ is as defined hereinbefore);

17) C_{2,3}alkylX⁹C_{1,2}alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);

5 18) C_{2,5}alkenyl which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, amino, C_{1,4}alkylamino, N,N-di(C_{1,4}alkyl)amino, aminosulphonyl, N-C_{1,4}alkylaminosulphonyl and N,N-di(C_{1,4}alkyl)aminosulphonyl;

19) C_{2,5}alkynyl which may be unsubstituted or which may be substituted with one or more
10 fluorine atoms or with one or two groups selected from hydroxy, amino, C_{1,4}alkylamino, N,N-di(C_{1,4}alkyl)amino, aminosulphonyl, N-C_{1,4}alkylaminosulphonyl and N,N-di(C_{1,4}alkyl)aminosulphonyl;

20) C_{2,4}alkenylX⁹C_{1,3}alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore); and

21) C_{2,4}alkynylX⁹C_{1,3}alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore)].

15 According to another aspect of the present invention preferably R² represents hydroxy, halogeno, nitro, trifluoromethyl, C_{1,3}alkyl, cyano, amino or R⁵X¹- [wherein X¹ is as hereinbefore defined and R⁵ is selected from one of the following nineteen groups:

1) C_{1,3}alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or C_{2,3}alkyl which may be unsubstituted or substituted with 1 or 2 groups selected from hydroxy
20 and amino;

2) 2-(3,3-dimethylureido)ethyl, 3-(3,3-dimethylureido)propyl, 2-(3-methylureido)ethyl, 3-(3-methylureido)propyl, 2-ureidoethyl, 3-ureidopropyl, 2-(N,N-dimethylcarbamoyloxy)ethyl, 3-(N,N-dimethylcarbamoyloxy)propyl, 2-(N-methylcarbamoyloxy)ethyl, 3-(N-methylcarbamoyloxy)propyl, 2-(carbamoyloxy)ethyl, 3-(carbamoyloxy)propyl;

25 3) C_{2,3}alkylX³R¹⁶ (wherein X³ is as defined hereinbefore and R¹⁶ is a group selected from C_{1,2}alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl, piperidinyl and tetrahydropyranyl which group is linked to X³ through a carbon atom and which C_{1,2}alkyl group may bear 1 or 2 substituents selected from hydroxy, halogeno and C_{1,2}alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy,
30 halogeno, C_{1,2}alkyl, C_{1,2}hydroxyalkyl and C_{1,2}alkoxy);

4) C_{2,3}alkylX⁴C_{2,3}alkylX⁵R²² (wherein X⁴ and X⁵ are as hereinbefore defined and R²² represents hydrogen or C_{1,2}alkyl);

- 5) $C_{1,2}$ alkylR¹³² (wherein R¹³² is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to $C_{1,2}$ alkyl through a carbon atom and which group may carry one substituent selected from oxo, hydroxy, halogeno, cyano, $C_{1,3}$ cyanoalkyl, $C_{1,3}$ alkyl, $C_{1,3}$ hydroxyalkyl, $C_{1,3}$ alkoxy, $C_{1,2}$ alkoxy $C_{1,3}$ alkyl and $C_{1,2}$ alkylsulphonyl $C_{1,3}$ alkyl) or $C_{2,3}$ alkylR¹³³ (wherein R¹³³ is a group selected from morpholino, thiomorpholino, piperidino, piperazin-1-yl and pyrrolidin-1-yl which group may carry one or two substituents selected from oxo, hydroxy, halogeno, cyano, $C_{1,3}$ cyanoalkyl, $C_{1,3}$ alkyl, $C_{1,3}$ hydroxyalkyl, $C_{1,3}$ alkoxy, $C_{1,2}$ alkoxy $C_{1,3}$ alkyl and $C_{1,2}$ alkylsulphonyl $C_{1,3}$ alkyl);
- 6) R²⁹ (wherein R²⁹ is as defined hereinbefore);
- 7) $C_{1,4}$ alkylR²⁹ (wherein R²⁹ is as defined hereinbefore);
- 8) 1-R²⁹but-2-en-4-yl (wherein R²⁹ is as defined hereinbefore);
- 9) 1-R²⁹but-2-yn-4-yl (wherein R²⁹ is as defined hereinbefore);
- 10) $C_{1,3}$ alkylX⁶R²⁹ (wherein X⁶ and R²⁹ are as defined hereinbefore);
- 11) 1-(R²⁹X⁷)but-2-en-4-yl (wherein X⁷ and R²⁹ are as defined hereinbefore);
- 12) 1-(R²⁹X⁸)but-2-yn-4-yl (wherein X⁸ and R²⁹ are as defined hereinbefore);
- 13) ethylX⁹methylR²⁹ (wherein X⁹ and R²⁹ are as defined hereinbefore);
- 14) R²⁸ (wherein R²⁸ is as defined hereinbefore);
- 15) ethylX⁹ $C_{1,2}$ alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore);
- 16) $C_{2,5}$ alkenyl which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, amino, $C_{1,4}$ alkylamino, N,N-di($C_{1,4}$ alkyl)amino, aminosulphonyl, N- $C_{1,4}$ alkylaminosulphonyl and N,N-di($C_{1,4}$ alkyl)aminosulphonyl;
- 17) $C_{2,5}$ alkynyl which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, amino, $C_{1,4}$ alkylamino, N,N-di($C_{1,4}$ alkyl)amino, aminosulphonyl, N- $C_{1,4}$ alkylaminosulphonyl and N,N-di($C_{1,4}$ alkyl)aminosulphonyl;
- 18) $C_{2,3}$ alkenylX⁹ $C_{1,3}$ alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore); and
- 19) $C_{2,3}$ alkynylX⁹ $C_{1,3}$ alkylR²⁸ (wherein X⁹ and R²⁸ are as defined hereinbefore)].

According to another aspect of the present invention more preferably R² represents hydroxy, $C_{1,3}$ alkyl, amino or R⁵X¹ - [wherein X¹ is as hereinbefore defined and R⁵ represents methyl, ethyl, benzyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl,

2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphanyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl ethyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, (1-methylpiperidin-3-yl)methyl, (1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonyl)ethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonyl)ethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonyl)ethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonyl)ethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-ylmethyl, 1-isopropylpiperidin-3-ylmethyl, 1-isopropylpiperidin-4-ylmethyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-(2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(2-ethylimidazol-1-yl)ethyl, 3-(2-methylimidazol-1-yl)propyl, 3-

(2-ethylimidazol-1-yl)propyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-((N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-((N-methyl-N-4-pyridyl)amino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-(2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl or 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yl].

According to another aspect of the present invention particularly R^2 represents C_{1-3} alkyl, amino or R^5X^1 - [wherein X^1 is as hereinbefore defined and R^5 represents ethyl, benzyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, (1-methylpiperidin-3-yl)methyl, (1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-

yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonyl)ethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonyl)ethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonyl)ethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonyl)ethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-yl)methyl, 1-isopropylpiperidin-3-yl)methyl, 1-isopropylpiperidin-4-yl)methyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-(2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-yl)methyl, 2-acetamidothiazol-4-yl)methyl, 1-methylimidazol-2-yl)methyl, 2-(imidazol-1-yl)ethyl, 2-(2-methylimidazol-1-yl)ethyl, 2-(2-ethylimidazol-1-yl)ethyl, 3-(2-methylimidazol-1-yl)propyl, 3-(2-ethylimidazol-1-yl)propyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridyl)methyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, morpholino, 2-(N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-(N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-(N-methyl-N-4-pyridyl)amino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-(2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)vinyl or 3-((2-(pyrrolidin-1-yl)ethyl)carbamoyl)prop-2-en-1-yl].

According to another aspect of the present invention more particularly R^2 represents C_{1-3} alkyl, amino or R^5X^1 - [wherein X^1 is as hereinbefore defined and R^5 represents ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl ethyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(methylpiperidino)ethyl, 3-(methylpiperidino)propyl, 2-(ethylpiperidino)ethyl, 3-(ethylpiperidino)propyl, 2-((2-methoxyethyl)piperidino)ethyl, 3-((2-methoxyethyl)piperidino)propyl, 2-((2-methylsulphonyl)ethylpiperidino)ethyl, 3-((2-methylsulphonyl)ethylpiperidino)propyl, piperidin-3-ylmethyl, piperidin-4-ylmethyl, 2-(piperidin-3-yl)ethyl, 2-(piperidin-4-yl)ethyl, 3-(piperidin-3-yl)propyl, 3-(piperidin-4-yl)propyl, (1-methylpiperidin-3-yl)methyl, (1-methylpiperidin-4-yl)methyl, (1-cyanomethylpiperidin-3-yl)methyl, (1-cyanomethylpiperidin-4-yl)methyl, 2-(methylpiperidin-3-yl)ethyl, 2-(methylpiperidin-4-yl)ethyl, 2-(1-cyanomethylpiperidin-3-yl)ethyl, 2-(1-cyanomethylpiperidin-4-yl)ethyl, 3-(methylpiperidin-3-yl)propyl, 3-(methylpiperidin-4-yl)propyl, 3-(1-cyanomethylpiperidin-3-yl)propyl, 3-(1-cyanomethylpiperidin-4-yl)propyl, 2-(ethylpiperidin-3-yl)ethyl, 2-(ethylpiperidin-4-yl)ethyl, 3-(ethylpiperidin-3-yl)propyl, 3-(ethylpiperidin-4-yl)propyl, ((2-methoxyethyl)piperidin-3-yl)methyl, ((2-methoxyethyl)piperidin-4-yl)methyl, 2-((2-methoxyethyl)piperidin-3-yl)ethyl, 2-((2-methoxyethyl)piperidin-4-yl)ethyl, 3-((2-methoxyethyl)piperidin-3-yl)propyl, 3-((2-methoxyethyl)piperidin-4-yl)propyl, (1-(2-methylsulphonyl)ethyl)piperidin-3-yl)methyl, (1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methyl, 2-((2-methylsulphonyl)ethyl)piperidin-3-yl)ethyl, 2-((2-methylsulphonyl)ethyl)piperidin-4-yl)ethyl, 3-((2-methylsulphonyl)ethyl)piperidin-3-yl)propyl, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propyl, 1-isopropylpiperidin-2-ylmethyl, 1-isopropylpiperidin-3-ylmethyl, 1-isopropylpiperidin-4-ylmethyl, 2-(1-isopropylpiperidin-2-yl)ethyl, 2-(1-isopropylpiperidin-3-yl)ethyl, 2-(1-isopropylpiperidin-4-yl)ethyl, 3-(1-isopropylpiperidin-2-yl)propyl, 3-(1-isopropylpiperidin-3-yl)propyl, 3-(1-isopropylpiperidin-4-yl)propyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, (pyrrolidin-2-yl)methyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(N-(2-methoxyethyl)-N-methylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(N-(2-methoxyethyl)-N-

methylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-thiomorpholinoethyl, 3-
 thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-
 dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-
 (4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl,
 5 morpholino, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-((N-methyl-N-4-
 pyridyl)amino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(2-(4-methylpiperazin-1-
 yl)ethoxy)ethyl, 3-(2-(4-methylpiperazin-1-yl)ethoxy)propyl, 2-(2-morpholinoethoxy)ethyl, 3-
 (2-morpholinoethoxy)propyl, 2-(tetrahydropyran-4-yloxy)ethyl, 3-(tetrahydropyran-4-
 yloxy)propyl, 2-((2-(pyrrolidin-1-yl)ethyl)carbonyl)vinyl or 3-((2-(pyrrolidin-1-
 10 yl)ethyl)carbonyl)prop-2-en-1-yl].

According to another embodiment of the present invention in another aspect R^2
 represents methoxy, 2-methoxyethoxy, 2-(2-methoxyethoxy)ethoxy, 3-methoxypropoxy, 2-
 methylsulphonylethoxy, 3-methylsulphonylpropoxy, benzyloxy, 2-(tetrahydropyran-4-
 yloxy)ethoxy, 3-(tetrahydropyran-4-yloxy)propoxy, 2-(4-methylpiperazin-1-yl)ethoxy, 3-(4-
 15 methylpiperazin-1-yl)propoxy, 2-morpholinoethoxy, 3-morpholinopropoxy, 2-(imidazol-1-
 yl)ethoxy, 3-(imidazol-1-yl)propoxy 2-(1,1-dioxothiomorpholino)ethoxy, 3-(1,1-
 dioxothiomorpholino)propoxy, 2-(1,2,3-triazol-1-yl)ethoxy, 3-(1,2,3-triazol-1-yl)propoxy, 2-
 (1,2,4-triazol-1-yl)ethoxy, 2-((N-methyl-N-4-pyridyl)amino)ethoxy, 2-(N,N-
 dimethylamino)ethoxy, 3-(N,N-dimethylamino)propoxy, 2-(N-methoxyacetyl-N-
 20 methylamino)ethoxy, 3-(N-methoxyacetyl-N-methylamino)propoxy, 1-methylpiperidin-3-
 ylmethoxy, 1-methylpiperidin-4-ylmethoxy, (1-cyanomethylpiperidin-3-yl)methoxy, (1-
 cyanomethylpiperidin-4-yl)methoxy, 2-(1-cyanomethylpiperidin-3-yl)ethoxy, 2-(1-
 cyanomethylpiperidin-4-yl)ethoxy, 3-(1-cyanomethylpiperidin-3-yl)propoxy, 3-(1-
 cyanomethylpiperidin-4-yl)propoxy, ((2-methoxyethyl)piperidin-3-yl)methoxy, ((2-
 25 methoxyethyl)piperidin-4-yl)methoxy, 2-(N-(2-methoxyethyl)-N-methylamino)ethoxy, 4-
 (pyrrolidin-1-yl)but-2-en-yloxy, 2-(2-oxopyrrolidin-1-yl)ethoxy, 3-(2-oxopyrrolidin-1-
 yl)propoxy, (pyrrolidin-2-yl)methoxy, 2-(pyrrolidin-1-yl)ethoxy, 3-(pyrrolidin-1-yl)propoxy,
 2-(2-(pyrrolidin-1-yl)ethoxy)ethoxy, (2-oxo-tetrahydro-2*H*-pyrrolidin-5-yl)methoxy, 2-(2-(4-
 methylpiperazin-1-yl)ethoxy)ethoxy, 2-piperidinoethoxy, 3-piperidinopropoxy, 2-
 30 (methylpiperidino)ethoxy, 3-(methylpiperidino)propoxy, 2-(ethylpiperidino)ethoxy, 3-
 (ethylpiperidino)propoxy, 2-((2-methoxyethyl)piperidino)ethoxy, 3-((2-
 methoxyethyl)piperidino)propoxy, 1-(2-methylsulphonylethyl)piperidin-3-ylmethoxy, 1-(2-


methylsulphonyl)ethyl)piperidin-4-ylmethoxy, 2-((2-methylsulphonyl)ethylpiperidino)ethoxy,
 3-((2-methylsulphonyl)ethylpiperidino)propoxy, piperidin-3-ylmethoxy, piperidin-4-
 ylmethoxy, 2-(piperidin-3-yl)ethoxy, 2-(piperidin-4-yl)ethoxy, 3-(piperidin-3-yl)propoxy, 3-
 (piperidin-4-yl)propoxy, 2-(methylpiperidin-3-yl)ethoxy, 2-(methylpiperidin-4-yl)ethoxy, 3-
 5 (methylpiperidin-3-yl)propoxy, 3-(methylpiperidin-4-yl)propoxy, 2-(ethylpiperidin-3-
 yl)ethoxy, 2-(ethylpiperidin-4-yl)ethoxy, 3-(ethylpiperidin-3-yl)propoxy, 3-(ethylpiperidin-4-
 yl)propoxy, 2-((2-methoxyethyl)piperidin-3-yl)ethoxy, 2-((2-methoxyethyl)piperidin-4-
 yl)ethoxy, 3-((2-methoxyethyl)piperidin-3-yl)propoxy, 3-((2-methoxyethyl)piperidin-4-
 yl)propoxy, 2-((2-methylsulphonyl)ethyl)piperidin-3-yl)ethoxy, 2-((2-
 10 methylsulphonyl)ethyl)piperidin-4-yl)ethoxy, 3-((2-methylsulphonyl)ethyl)piperidin-3-
 yl)propoxy, 3-((2-methylsulphonyl)ethyl)piperidin-4-yl)propoxy, 1-isopropylpiperidin-2-
 ylmethoxy, 1-isopropylpiperidin-3-ylmethoxy, 1-isopropylpiperidin-4-ylmethoxy, 2-(1-
 isopropylpiperidin-2-yl)ethoxy, 2-(1-isopropylpiperidin-3-yl)ethoxy, 2-(1-isopropylpiperidin-
 4-yl)ethoxy, 3-(1-isopropylpiperidin-2-yl)propoxy, 3-(1-isopropylpiperidin-3-yl)propoxy, 3-
 15 (1-isopropylpiperidin-4-yl)propoxy, 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethoxy, 3-(2-(4-
 methylpiperazin-1-yl)ethoxy)propoxy, 2-(2-morpholinoethoxy)ethoxy, 3-(2-
 morpholinoethoxy)propoxy, 2-((2-(pyrrolidin-1-yl)ethyl)carbonyl)vinyl or 3-((2-(pyrrolidin-
 1-yl)ethyl)carbonyl)prop-2-en-1-yl.

Where one of the R^2 substituents is R^5X^1 - the substituent R^5X^1 - is preferably at the 6-
 20 or 7-position of the quinazoline ring, more preferably at the 7-position of the quinazoline ring.

When one of the R^2 substituents is at the 6-position of the quinazoline ring it is
 preferably hydrogen, halogeno, C_{1-3} alkyl, trifluoromethyl, C_{1-3} alkoxy, C_{1-3} alkylsulphanyl or -
 NR^3R^4 (wherein R^3 and R^4 are as defined hereinbefore).

When one of the R^2 substituents is at the 6-position of the quinazoline ring it is more
 25 preferably C_{1-3} alkoxy, especially methoxy.

In another aspect of the present invention there is provided the use of compounds of
 the formula Ia:




(Ia)

[wherein:

- 10 ring C, R^1 , R^2 , n and Z are as defined hereinbefore with the provisos that R^2 is not hydrogen and that Z is not CH_2 or a direct bond; and
- R^{2a} represents hydrogen, halogeno, C_{1-3} alkyl, trifluoromethyl, C_{1-3} alkoxy, C_{1-3} alkylsulphonyl, -
 $NR^{3a}R^{4a}$ (wherein R^{3a} and R^{4a} , which may be the same or different, each represents hydrogen
 or C_{1-3} alkyl), or $R^{5a}(CH_2)_{za}X^{1a}$ (wherein R^{5a} is a 5- or 6-membered saturated heterocyclic group
 15 with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may
 bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-4} cyanoalkyl, C_{1-4}
 alkyl, C_{1-4} hydroxyalkyl, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, C_{1-4} alkylsulphonyl C_{1-4} alkyl, C_{1-4}
 alkoxycarbonyl, C_{1-4} aminoalkyl, C_{1-4} alkylamino, di(C_{1-4} alkyl)amino, C_{1-4} alkylamino C_{1-4} alkyl,
 di(C_{1-4} alkyl)amino C_{1-4} alkyl, C_{1-4} alkylamino C_{1-4} alkoxy, di(C_{1-4} alkyl)amino C_{1-4} alkoxy and a
 20 group $-(O)_f(C_{1-4}alkyl)_g$ ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered
 saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N,
 which cyclic group may bear one or more substituents selected from C_{1-4} alkyl), za is an
 integer from 0 to 4 and X^{1a} represents a direct bond, -O-, $-CH_2-$, -S-, -SO-, $-SO_2-$, $-NR^{6a}C(O)-$,
 $-C(O)NR^{7a}-$, $-SO_2NR^{8a}-$, $-NR^{9a}SO_2-$ or $-NR^{10a}-$ (wherein R^{6a} , R^{7a} , R^{8a} , R^{9a} and R^{10a} each
 25 independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));
 and salts thereof, and prodrugs thereof for example esters and amides, in the manufacture of a
 medicament for use in the production of an antiangiogenic and/or vascular permeability
 reducing effect in warm-blooded animals such as humans.

In another aspect of the present invention there is provided the use of compounds of
 30 the formula Ia:



(Ia)

[wherein:

- 10 ring C , R^1 , R^2 , n and Z are as defined hereinbefore with the provisos that R^2 is not hydrogen and that Z is not CH_2 or a direct bond; and
- R^{2a} represents hydrogen, halogeno, C_{1-3} alkyl, trifluoromethyl, C_{1-3} alkoxy, C_{1-3} alkylsulphanyl, -
 $NR^{3a}R^{4a}$ (wherein R^{3a} and R^{4a} , which may be the same or different, each represents hydrogen
 or C_{1-3} alkyl), or $R^{5a}(CH_2)_{za}X^{1a}$ (wherein R^{5a} is a 5- or 6-membered saturated heterocyclic group
 15 with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may
 bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and
 C_{1-4} alkoxy, za is an integer from 0 to 4 and X^{1a} represents a direct bond, -O-, - CH_2 -, -S-, -SO-,
 -SO₂-, -NR^{6a}C(O)-, -C(O)NR^{7a}-, -SO₂NR^{8a}-, -NR^{9a}SO₂- or -NR^{10a}- (wherein R^{6a} , R^{7a} , R^{8a} , R^{9a}
 and R^{10a} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));
- 20 and salts thereof, and prodrugs thereof for example esters, amides and sulphides, in the
 manufacture of a medicament for use in the production of an antiangiogenic and/or vascular
 permeability reducing effect in warm-blooded animals such as humans.

Advantageously X^{1a} represents -O-, -S-, -NR^{6a}C(O)-, -NR^{9a}SO₂- or -NR^{10a}- (wherein
 R^{6a} , R^{9a} and R^{10a} each independently represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl).

- 25 Preferably X^{1a} represents -O-, -S-, -NR^{6a}CO-, -NR^{9a}SO₂- (wherein R^{6a} and R^{9a} each
 independently represents hydrogen or C_{1-2} alkyl) or NH.

More preferably X^{1a} represents -O-, -S-, -NR^{6a}CO- (wherein R^{6a} represents hydrogen or
 C_{1-2} alkyl) or NH.

- Particularly X^{1a} represents -O- or -NR^{6a}CO- (wherein R^{6a} represents hydrogen or C_{1-2}
 30 C_{1-2} alkyl), more particularly -O- or -NHCO-, especially -O-.

Preferably za is an integer from 1 to 3.

Preferably R^{5a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(O-)(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C_{1-3} alkyl).

More preferably R^{5a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-3} cyanoalkyl, C_{1-3} alkyl, C_{1-3} hydroxyalkyl, C_{1-3} alkoxy, C_{1-2} alkoxy C_{1-3} alkyl, C_{1-2} alkylsulphonyl C_{1-3} alkyl, C_{1-3} alkoxycarbonyl, C_{1-3} alkylamino, di(C_{1-3} alkyl)amino, C_{1-3} alkylamino C_{1-3} alkyl, di(C_{1-3} alkyl)amino C_{1-3} alkyl, C_{1-3} alkylamino C_{1-3} alkoxy, di(C_{1-3} alkyl)amino C_{1-3} alkoxy and a group $-(O-)(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

Particularly R^{5a} is pyrrolidinyl, piperazinyl, piperidinyl, azetidiny, morpholino or thiomorpholino which group may bear 1 or 2 substituents selected from a group $-(O-)(C_{1-3}alkyl)_g ringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, methylpiperazinyl, piperidinyl, azetidiny, morpholino and thiomorpholino).

According to another aspect of the present invention preferably R^{5a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, morpholino and thiomorpholino which group may carry 1 or 2 substituents selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy.

Advantageously R^{2a} represents C_{1-3} alkyl, C_{1-3} alkoxy, amino or $R^{5a}(CH_2)_{za}X^{1a}$ (wherein R^{5a} , X^{1a} and za are as defined hereinbefore). Another advantageous value of R^{2a} is hydrogen.

Preferably R^{2a} is methyl, ethyl, methoxy, ethoxy or $R^{5a}(CH_2)_{za}X^{1a}$ (wherein R^{5a} , X^{1a} and za are as defined hereinbefore). Another preferred value of R^{2a} is hydrogen.


More preferably R^{2a} is methyl, ethyl, methoxy, ethoxy or $R^{5a}(CH_2)_{za}X^{1a}$ (wherein R^{5a} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, morpholino and thiomorpholino

which group may carry 1 or 2 substituents selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy, X^{1a} is -O-, -S-, $-NR^{6a}C(O)-$, $-NR^{9a}SO_2-$ (wherein R^{6a} and R^{9a} each independently represents hydrogen or C_{1-2} alkyl) or NH, and za is an integer from 1 to 3).

Particularly R^{2a} represents methyl, methoxy or $R^{5a}(CH_2)_{za}X^{1a}$ (wherein R^{5a} , X^{1a} and za are as defined hereinbefore).

More particularly R^{2a} represents methoxy.

In a further aspect of the present invention there is provided the use of compounds of the formula Ib:



(Ib)

[wherein:


ring C, R^1 , R^2 , R^{2a} and n are as defined hereinbefore with the proviso that R^2 is not hydrogen; and

Zb is -O- or -S-;

and salts thereof, and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides, in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals such as humans.

Preferably Zb is -O-.

According to another aspect of the present invention there are provided compounds of the formula II:



(II)


10 [wherein:

ring C, R^1 , R^2 , R^{2a} , Zb and n are as defined hereinbefore with the proviso that R^2 is not hydrogen and excluding the compounds:

6,7-dimethoxy-4-(1-naphthylsulphonyl)quinazoline, 6,7-dimethoxy-4-(2-naphthylsulphonyl)quinazoline, 6,7-dimethoxy-4-(1-naphthyloxy)quinazoline and 6,7-dimethoxy-4-(2-naphthyloxy)quinazoline;

and salts thereof, and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides.

According to another aspect of the present invention there are provided compounds of the formula IIa:



(IIa)

[wherein:


ring C, R^1 , R^2 , R^{2a} , Zb and n are as defined hereinbefore with the proviso that R^2 does not have any of the following values:

hydrogen, substituted or unsubstituted C_{1-5} alkyl, halogeno or phenoxy and excluding the compounds:

6,7-dimethoxy-4-(1-naphthylsulphanyl)quinazoline, 6,7-dimethoxy-4-(2-naphthylsulphanyl)quinazoline, 6,7-dimethoxy-4-(1-naphthyloxy)quinazoline and 6,7-dimethoxy-4-(2-naphthyloxy)quinazoline;

and salts thereof, and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides.

According to another aspect of the present invention there are provided compounds of the formula IIb:



(IIb)

[wherein:

ring C, R^1 , R^2 , R^{2a} , Zb and n are as defined hereinbefore with the proviso that R^2 does not have any of the following values:

hydrogen, substituted or unsubstituted C_{1-5} alkyl, halogeno, C_{1-5} alkoxy, C_{2-5} alkenyl, phenoxy or phenyl C_{1-5} alkoxy;

and salts thereof, and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides.

Preferred compounds of the present invention include

6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(2-naphthyloxy)quinazoline,
6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,
7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(quinolin-7-yloxy)quinazoline,

- 6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)-4-(quinolin-7-yloxy)quinazoline,
6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,
4-(4-chloroquinolin-7-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(4-methylquinolin-7-yloxy)quinazoline,
5 6-methoxy-4-(4-methylquinolin-7-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(quinolin-7-yloxy)quinazoline,
6-methoxy-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,
4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline,
10 4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,
6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline,
6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline,
(*R,S*)-4-(3-fluoroquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline,
15 4-(indol-5-yloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline,
7-(3-*N,N*-dimethylaminopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(2-morpholinoethoxy)ethoxy)quinazoline,
7-(2-(*N,N*-diethylamino)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
20 6-methoxy-7-(3-piperidinopropoxy)-4-(quinolin-7-yloxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(2-(piperidin-1-yl)ethoxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(2-(1*H*-1,2,4-triazol-1-yl)ethoxy)quinazoline,
6-methoxy-7-(3-piperidinopropoxy)-4-(6-trifluoromethylindol-5-yloxy)quinazoline,
25 7-(3-(methylsulphonyl)propoxy)-4-(2-methylindol-5-yloxy)quinazoline,
7-(3-(*N,N*-dimethylamino)propoxy)-4-(2,3-dimethylindol-5-yloxy)-6-methoxyquinazoline,
4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-3-ylmethoxy)quinazoline,
7-(2-(*N,N*-diethylamino)ethoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(2-(piperidin-2-yl)ethoxy)quinazoline,
30 4-(indol-5-yloxy)-6-methoxy-7-(2-(piperidin-1-yl)ethoxy)quinazoline,
4-(indol-6-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
7-(3-(ethylsulphonyl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,

6-methoxy-4-(3-methylindol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline,
7-(2-hydroxy-3-piperidinopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
7-(2-hydroxy-3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,

- 5 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methylamino)ethoxy)quinazoline, and
7-(2-hydroxy-3-(isopropylamino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example
esters and amides.

Especially preferred compounds of the present invention include

- 10 6-methoxy-7-(3-morpholinopropoxy)-4-(quinolin-7-yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-methylsulphonylpropoxy)quinazoline,
15 7-((1-cyanomethyl)piperidin-4-ylmethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-morpholinoethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-pyrrolidin-1-ylethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-methylpiperidin-3-ylmethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-piperidinoethoxy)quinazoline,
20 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-(4-
pyridyl)amino)ethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(1*H*-1,2,4-triazol-1-yl)ethoxy)quinazoline,
25 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(2-(4-methylpiperazin-1-
yl)ethoxy)ethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline
6-methoxy-7-(1-(2-methoxyethyl)piperidin-4-ylmethoxy)-4-(2-methylindol-5-
30 yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-((2-(2-pyrrolidin-1-
ylethyl)carbamoyl)vinyl)quinazoline,

- 6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(piperidin-4-yloxy)ethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-
5 methylsulphonylamino)ethoxy)quinazoline,
7-(2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethoxy)-6-methoxy-4-(2-methylindol-5-
yloxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(3-(pyrrolidin-yl)propoxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(3-(1,1-dioxothiomorpholino)propoxy)quinazoline,
10 4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline,
7-(3-(N,N-dimethylamino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
7-(3-(N,N-diethylamino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
7-(3-(1,1-dioxothiomorpholino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
15 4-(indol-5-yloxy)-6-methoxy-7-(2-(4-pyridyloxy)ethoxy)quinazoline,
4-(indol-6-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
7-(1-(2-methoxyethyl)piperidin-4-ylmethoxy)-4-(2-methylindol-5-yloxy)quinazoline,
7-(2-hydroxy-3-morpholinopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
7-(2-(1-(2-methoxyethyl)piperidin-4-yl)ethoxy)-6-methoxy-4-(2-methylindol-5-
20 yloxy)quinazoline,
7-(2-hydroxy-3-pyrrolidin-1-ylpropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
7-(3-(N,N-diethylamino)-2-hydroxypropoxy)-6-methoxy-4-(2-methylindol-5-
yloxy)quinazoline,
7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
25 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(4-pyridyloxy)ethoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
(2*R*)-6-methoxy-(2-methyl-1*H*-indol-5-yloxy)-7-(2-hydroxy-3-
piperidinopropoxy)quinazoline,
(5*R*)-6-methoxy-4-(2-methyl-1*H*-indol-5-yloxy)-7-(2-oxopyrrolidin-5-ylmethoxy)quinazoline,
30 4-(4-bromoindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-(2-(pyrrolidin-1-yl)ethyl)-piperidin-4-
ylmethoxy)quinazoline,

(2R)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
(2R)-7-(2-hydroxy-3-morpholinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
(2R)-7-(2-hydroxy-3-piperidinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
(2S)-7-(2-hydroxy-3-((N,N)-diisopropyl)amino)propoxy)-4-(indol-5-yloxy)-6-
5 methoxyquinazoline,
(2S)-7-(2-hydroxy-3-piperidinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
(2R)-7-(2-hydroxy-3-piperidinopropoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline,
(2R)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-4-(3-methylindol-5-
yloxy)quinazoline,
10 (2R)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-
yloxy)quinazoline,
(2R)-7-(2-hydroxy-3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-
yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-(2-morpholinoethyl)piperidin-4-
15 ylmethoxy)quinazoline,
4-(3-fluoro-quinolin-7-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
4-(3-fluoro-quinolin-7-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)-4-(1H-pyrrolo[2,3-b]pyridin-5-yloxy)quinazoline,
(2S)-6-methoxy-(2-methyl-1H-indol-5-yloxy)-7-(2-hydroxy-3-piperidinopropoxy)quinazoline,
20 and
4-(6-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example
esters and amides.
More especially preferred compounds of the present invention include
25 6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline,
4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline,
4-(6-fluoroindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
30 4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,

4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline,

5 4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline,
(2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxyquinazoline, and

4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline,

10 and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters and amides.

Thus preferred compounds of the present invention include those, the preparation of which is described in Examples 23, 10, 5, 176, 7, 22, 13, 15, 177, 12, 35, 47, 44, 45, 157, 52, 62, 66, 75, 159, 87, 88, 89, 167, 83, 97, 101, 108, 113, 114, 121, 124, 178, 162, 165, 150 and 166,

15 and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters and amides.

Thus especially preferred compounds of the present invention include those, the preparation of which is described in Examples 2, 11, 34, 36, 186, 151, 57, 54, 55, 58, 56, 60, 61, 64, 65, 67, 68, 71, 72, 74, 70, 77, 79, 80, 82, 86, 122, 107, 110, 112, 117, 118, 119, 123, 161, 147, 20 163, 164, 63, 78, 115, 320, 318, 290, 252, 292, 293, 294, 301, 299, 279, 280, 305, 269, 246, 266, 267, 182, 321 and 250,

and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters and amides.

Thus more especially preferred compounds of the present invention include those, the preparation of which is described in Examples 9, 243, 251, 245, 247, 249, 240, 238, 237, 239, 25 241, 258 and 322,

and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters and amides.

In another embodiment, preferred compounds of the present invention include

30 6-methoxy-7-(3-morpholinopropoxy)-4-(quinolin-6-yloxy)quinazoline,
(*S*)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,
6-methoxy-7-(3-morpholinopropoxy)-4-(1-naphthyloxy)quinazoline,

- 4-(1*H*-indazol-5-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6,7-dimethoxy-4-(quinolin-7-yloxy)quinazoline,
6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(2,2,4-trimethyl-1,2-dihydroquinolin-6-yloxy)quinazoline,
5 6-methoxy-7-((2-piperidin-1-yl)ethoxy)-4-(quinolin-7-yloxy)quinazoline,
6-methoxy-4-(2-methylquinolin-7-yloxy)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,
6-methoxy-4-(2-methylquinolin-7-yloxy)-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)quinazoline,
6-methoxy-4-(2-methylquinolin-7-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,
10 4-(2-chloro-1*H*-benzimidazol-5-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,
4-(2,4-dimethylquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,
4-(1*H*-indazol-6-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
15 4-(1,3-benzothiazol-6-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(3-oxo-2*H*-4*H*-1,4-benzoxazin-6-yloxy)quinazoline,
7-hydroxy-6-methoxy-4-(quinolin-7-yloxy)quinazoline,
6-methoxy-4-(2-methyl-1,3-benzothiazol-5-yloxy)-7-(3-methylsulphonylpropoxy)quinazoline,
20 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(tetrahydropyran-4-yloxy)ethoxy)quinazoline,
6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(1,2-cycloheptanebenzimidazol-5-yloxy)quinazoline,
6-methoxy-7-(3-morpholinopropoxy)-4-(quinolin-2-yloxy)quinazoline,
25 6-methoxy-7-(3-morpholinopropoxy)-4-(3-oxo-1,2-dihydro-3*H*-indazol-1-yl)quinazoline,
4-(2,3-dihydro-1*H*-indan-5-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-4-(2-methyl-4-oxo-4*H*-chromen-7-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,
6-methoxy-4-(4-methyl-4*H*-1,4-benzoxazin-6-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,
30 6-methoxy-4-(2-methyl-4-oxo-4*H*-chromen-7-yloxy)-7-((3-pyrrolidin-1-yl)propoxy)quinazoline,

6-methoxy-4-(4-methyl-3,4-dihydro-2*H*-1,4-benzoxazin-6-yloxy)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,

7-benzyloxy-6-methoxy-4-(quinolin-7-yloxy)quinazoline,

4-(2,4-dimethylquinolin-7-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,

5 6-methoxy-7-(3-methylsulphonylpropoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline,

6-methoxy-4-(2-methylquinolin-7-yloxy)-7-(3-methylsulphonylpropoxy)quinazoline,

6-methoxy-7-(3-morpholinopropoxy)-4-(quinazolin-7-yloxy)quinazoline,

6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)-4-(3-oxo-2*H*-4*H*-1,4-benzoxazin-6-yloxy)quinazoline,

10 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,

6,7-dimethoxy-4-(2-methyl-1*H*-benzimidazol-5-yloxy)quinazoline,

and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides.

In another embodiment more preferred compounds of the present invention include

15 6-methoxy-4-(4-methylquinolin-7-yloxy)-7-(3-morpholinopropoxy)quinazoline,

6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(quinolin-6-yloxy)quinazoline,

6-methoxy-4-(2-methyl-1,3-benzothiazol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline,

(*R*)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,

20 6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)-4-(2,2,4-trimethyl-1,2-dihydroquinolin-6-yloxy)quinazoline,

6-methoxy-7-(2-morpholinoethoxy)-4-(quinolin-7-yloxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-ylamino)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

25 6-methoxy-4-(2-methylindol-5-ylamino)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,

4-(4-chloroquinolin-7-yloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline,

4-(7-hydroxy-2-naphthyloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline,

6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline,

7-(2-(*N,N*-dimethylamino)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,

30 6-methoxy-7-(2-(*N*-(2-methoxyethyl)-*N*-methylamino)ethoxy)-4-(2-methylindol-5-yloxy)quinazoline,

4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,
(S)-6-methoxy-7-((2-oxo-tetrahydro-2H-pyrrolidin-5-yl)methoxy)-4-(quinolin-7-

5 yloxy)quinazoline,

and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides.

In another embodiment especially preferred compounds of the present invention include
6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,

10 4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-methylsulphonylpropoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-((1-methylpiperidin-3-yl)methoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(piperidin-1-yl)ethoxy)quinazoline,

6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,

15 6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

4-(indol-5-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

4-(indol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,

6-methoxy-7-(3-methylsulphonylpropoxy)-4-(quinolin-7-yloxy)quinazoline,

20 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-(4-pyridyl)amino)ethoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline,

6-methoxy-7-(3-morpholinopropoxy)-4-(quinolin-7-yloxy)quinazoline,

25 6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(2-naphthyloxy)quinazoline,

7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(quinolin-7-yloxy)quinazoline,

6-methoxy-7-(3-(1-methylpiperazin-4-yl)propoxy)-4-(quinolin-7-yloxy)quinazoline,

4-(4-chloroquinolin-7-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,

6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(4-methylquinolin-7-yloxy)quinazoline,

30 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(quinolin-7-yloxy)quinazoline,

6-methoxy-7-((1-(2-methylsulphonylethyl)piperidin-4-yl)methoxy)-4-(quinolin-7-yloxy)-quinazoline,

7-((1-cyanomethylpiperidin-4-yl)methoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline,

4-(3-fluoroquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline,
5 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-morpholinoethoxy)quinazoline,

6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(2-methylindol-5-yloxy)quinazoline,

7-(3-(N,N-dimethylamino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,

7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(1-methylpiperazin-4-

10 yl)ethoxy)ethoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(2-morpholinoethoxy)ethoxy)quinazoline,

6-methoxy-4-(4-methylquinolin-7-yloxy)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(1,2,4-triazol-1-yl)ethoxy)quinazoline,

4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

15 4-(indol-5-yloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline,

and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides.

In another aspect of the present invention preferred compounds include

6-methoxy-7-((1-(2-methoxyethyl)piperidin-4-yl)methoxy)-4-(2-methylindol-5-

20 yloxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(2-(pyrrolidin-1-yl)ethylcarbamoyl)vinyl)quinazoline,

4-(3-cyanoquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(4-trifluoromethylquinolin-7-

25 yloxy)quinazoline,

6-methoxy-4-(2-methyl-1*H*-benzimidazol-5-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

4-(3-carbamoylquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,

30 6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-(1-methylpiperazin-4-yl)propoxy)quinazoline,

6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline,

and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides.

An especially preferred compound of the present invention is

6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline

5 and salts thereof especially hydrochloride salts thereof and prodrugs thereof for example esters, amides and sulphides, preferably esters and amides.

For the avoidance of doubt it is to be understood that where in this specification a group is qualified by 'hereinbefore defined' or 'defined hereinbefore' the said group encompasses the first occurring and broadest definition as well as each and all of the preferred
10 definitions for that group.

In this specification unless stated otherwise the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. An analogous convention applies to other generic terms. Unless otherwise stated the term "alkyl" advantageously refers to chains with 1-6
15 carbon atoms, preferably 1-4 carbon atoms. The term "alkoxy" as used herein, unless stated otherwise includes "alkyl"-O- groups in which "alkyl" is as hereinbefore defined. The term "aryl" as used herein unless stated otherwise includes reference to a C₆₋₁₀ aryl group which may, if desired, carry one or more substituents selected from halogeno, alkyl, alkoxy, nitro, trifluoromethyl and cyano, (wherein alkyl and alkoxy are as hereinbefore defined). The term
20 "aryloxy" as used herein unless otherwise stated includes "aryl"-O-groups in which "aryl" is as hereinbefore defined. The term "sulphonyloxy" as used herein refers to alkylsulphonyloxy and arylsulphonyloxy groups in which "alkyl" and "aryl" are as hereinbefore defined. The term "alkanoyl" as used herein unless otherwise stated includes formyl and alkylC=O groups in which "alkyl" is as defined hereinbefore, for example C₂alkanoyl is ethanoyl and refers to
25 CH₃C=O, C₁alkanoyl is formyl and refers to CHO. In this specification unless stated otherwise the term "alkenyl" includes both straight and branched chain alkenyl groups but references to individual alkenyl groups such as 2-butenyl are specific for the straight chain version only. Unless otherwise stated the term "alkenyl" advantageously refers to chains with 2-5 carbon atoms, preferably 3-4 carbon atoms. In this specification unless stated otherwise
30 the term "alkynyl" includes both straight and branched chain alkynyl groups but references to individual alkynyl groups such as 2-butylnyl are specific for the straight chain version only. Unless otherwise stated the term "alkynyl" advantageously refers to chains with 2-5 carbon

atoms, preferably 3-4 carbon atoms. Unless stated otherwise the term "haloalkyl" refers to an alkyl group as defined hereinbefore which bears one or more halogeno groups, such as for example trifluoromethyl.

For the avoidance of any doubt, where R^2 has a value of substituted or unsubstituted C_{1-5} alkyl, R^2 has been selected from C_{1-3} alkyl or from a group R^5X^1 wherein X^1 is a direct bond or $-CH_2-$ and R^5 is C_{1-5} alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, chloro, bromo and amino.

Within the present invention it is to be understood that a compound of the formula I or a salt thereof may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be understood that the invention encompasses any tautomeric form which inhibits VEGF receptor tyrosine kinase activity and is not to be limited merely to any one tautomeric form utilised within the formulae drawings. The formulae drawings within this specification can represent only one of the possible tautomeric forms and it is to be understood that the specification encompasses all possible tautomeric forms of the compounds drawn not just those forms which it has been possible to show graphically herein.

It will be appreciated that compounds of the formula I or a salt thereof may possess an asymmetric carbon atom. Such an asymmetric carbon atom is also involved in the tautomerism described above, and it is to be understood that the present invention encompasses any chiral form (including both pure enantiomers, scalemic and racemic mixtures) as well as any tautomeric form which inhibits VEGF receptor tyrosine kinase activity, and is not to be limited merely to any one tautomeric form or chiral form utilised within the formulae drawings. It is to be understood that the invention encompasses all optical and diastereomers which inhibit VEGF receptor tyrosine kinase activity. It is further to be understood that in the names of chiral compounds (R,S) denotes any scalemic or racemic mixture while (R) and (S) denote the enantiomers. In the absence of (R,S), (R) or (S) in the name it is to be understood that the name refers to any scalemic or racemic mixture, wherein a scalemic mixture contains R and S enantiomers in any relative proportions and a racemic mixture contains R and S enantiomers in the ratio 50:50.

It is also to be understood that certain compounds of the formula I and salts thereof can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to

be understood that the invention encompasses all such solvated forms which inhibit VEGF receptor tyrosine kinase activity.

For the avoidance of any doubt, it is to be understood that when X^1 is, for example, a group of formula $-NR^6C(O)-$, it is the nitrogen atom bearing the R^6 group which is attached to the quinazoline ring and the carbonyl ($C(O)$) group is attached to R^5 , whereas when X^1 is, for example, a group of formula $-C(O)NR^7-$, it is the carbonyl group which is attached to the quinazoline ring and the nitrogen atom bearing the R^7 group is attached to R^5 . A similar convention applies to the other two atom X^1 linking groups such as $-NR^9SO_2-$ and $-SO_2NR^8-$. When X^1 is $-NR^{10}-$ it is the nitrogen atom bearing the R^{10} group which is linked to the quinazoline ring and to R^5 . An analogous convention applies to other groups. It is further to be understood that when X^1 represents $-NR^{10}-$ and R^{10} is $C_{1-3}alkoxyC_{2-3}alkyl$ it is the $C_{2-3}alkyl$ moiety which is linked to the nitrogen atom of X^1 and an analogous convention applies to other groups.

For the avoidance of any doubt, it is to be understood that in a compound of the formula I when R^5 is, for example, a group of formula $C_{1-3}alkylX^9C_{1-3}alkylR^{29}$, it is the terminal $C_{1-3}alkyl$ moiety which is linked to X^1 , similarly when R^5 is, for example, a group of formula $C_{2-5}alkenylR^{28}$ it is the $C_{2-5}alkenyl$ moiety which is linked to X^1 and an analogous convention applies to other groups. When R^5 is a group $1-R^{29}prop-1-en-3-yl$ it is the first carbon to which the group R^{29} is attached and it is the third carbon which is linked to X^1 and an analogous convention applies to other groups.

For the avoidance of any doubt, it is to be understood that in a compound of the formula I when R^5 is, for example, R^{28} and R^{28} is a pyrrolidinyl ring which bears a group $-(O-)(C_{1-4}alkyl)_gringD$, it is the $-O-$ or $C_{1-4}alkyl$ which is linked to the pyrrolidinyl ring, unless f and g are both 0 when it is ring D which is linked to the pyrrolidinyl ring and an analogous convention applies to other groups.

For the avoidance of any doubt, it is to be understood that when R^{29} carries a $C_{1-4}aminoalkyl$ substituent it is the $C_{1-4}alkyl$ moiety which is attached to R^{29} whereas when R^{29} carries a $C_{1-4}alkylamino$ substituent it is the amino moiety which is attached to R^{29} and an analogous convention applies to other groups.

For the avoidance of any doubt, it is to be understood that when R^{28} carries a $C_{1-4}alkoxyC_{1-4}alkyl$ substituent it is the $C_{1-4}alkyl$ moiety which is attached to R^{28} and an analogous convention applies to other groups.

The present invention relates to the compounds of formula I as hereinbefore defined as well as to the salts thereof. Salts for use in pharmaceutical compositions will be pharmaceutically acceptable salts, but other salts may be useful in the production of the compounds of formula I and their pharmaceutically acceptable salts. Pharmaceutically acceptable salts of the invention may, for example, include acid addition salts of the compounds of formula I as hereinbefore defined which are sufficiently basic to form such salts. Such acid addition salts include for example salts with inorganic or organic acids affording pharmaceutically acceptable anions such as with hydrogen halides (especially hydrochloric or hydrobromic acid of which hydrochloric acid is particularly preferred) or with sulphuric or phosphoric acid, or with trifluoroacetic, citric or maleic acid. In addition where the compounds of formula I are sufficiently acidic, pharmaceutically acceptable salts may be formed with an inorganic or organic base which affords a pharmaceutically acceptable cation. Such salts with inorganic or organic bases include for example an alkali metal salt, such as a sodium or potassium salt, an alkaline earth metal salt such as a calcium or magnesium salt, an ammonium salt or for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.


A compound of the formula I, or salt thereof, and other compounds of the invention (as hereinafter defined) may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Such processes include, for example, those illustrated in European Patent Applications Publication Nos. 0520722, 0566226, 0602851 and 0635498. Such processes also include, for example, solid phase synthesis. Such processes, are provided as a further feature of the invention and are as described hereinafter. Necessary starting materials may be obtained by standard procedures of organic chemistry. The preparation of such starting materials is described within the accompanying non-limiting Examples. Alternatively necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

Thus, the following processes (a) to (f) and (i) to (vi) constitute further features of the present invention.

Synthesis of Compounds of Formula I


(a) Compounds of the formula I and salts thereof may be prepared by the reaction of a compound of the formula III:

- 67 -



(III)

(wherein R^2 and m are as defined hereinbefore and L^1 is a displaceable moiety), with a compound of the formula IV:



(IV)

(wherein ring C, R^1 , Z and n are as defined hereinbefore) to obtain compounds of the formula I and salts thereof. A convenient displaceable moiety L^1 is, for example, a halogeno, alkoxy (preferably C_{1-4} alkoxy), aryloxy, alkylsulphanyl, arylsulphanyl, alkoxyalkylsulphanyl or sulphonyloxy group, for example a chloro, bromo, methoxy, phenoxy, methylsulphanyl, 2-methoxyethylsulphanyl, methanesulphonyloxy or toluene-4-sulphonyloxy group.


The reaction is advantageously effected in the presence of a base. Such a base is, for example, an organic amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, morpholine, N-methylemorpholine or diazabicyclo[5.4.0]undec-7-ene, tetramethylguanidine or for example, an alkali metal or alkaline earth metal carbonate or hydroxide, for example sodium carbonate, potassium carbonate, calcium carbonate, sodium hydroxide or potassium hydroxide. Alternatively such a base is, for example, an alkali metal hydride, for example sodium hydride, or an alkali metal or alkaline earth metal amide, for example sodium amide, sodium bis(trimethylsilyl)amide, potassium amide or potassium bis(trimethylsilyl)amide. The reaction is preferably effected in the presence of an inert solvent or diluent, for example an ether such as tetrahydrofuran or 1,4-dioxan, an aromatic hydrocarbon solvent such as toluene, or a dipolar aprotic solvent such

as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one or dimethyl sulphoxide. The reaction is conveniently effected at a temperature in the range, for example, 10 to 150°C, preferably in the range 20 to 90°C.

When it is desired to obtain the acid salt, the free base may be treated with an acid such as a hydrogen halide, for example hydrogen chloride, sulphuric acid, a sulphonic acid, for example methane sulphonic acid, or a carboxylic acid, for example acetic or citric acid, using a conventional procedure.

(b) Production of those compounds of formula I and salts thereof wherein at least one R^2 is R^5X^1 wherein R^5 is as defined hereinbefore and X^1 is -O-, -S-, -OC(O)- or -NR¹⁰-

(wherein R^{10} independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) can be achieved by the reaction, conveniently in the presence of a base (as defined hereinbefore in process (a)) of a compound of the formula V:



(V)


(wherein ring C, Z, R^1 , R^2 and n are as hereinbefore defined and X^1 is as hereinbefore defined in this section and s is an integer from 0 to 2) with a compound of formula VI:



(wherein R^5 and L^1 are as hereinbefore defined), L^1 is a displaceable moiety for example a halogeno or sulphonyloxy group such as a bromo, methanesulphonyloxy or toluene-4-sulphonyloxy group, or L^1 may be generated in situ from an alcohol under standard Mitsunobu conditions ("Organic Reactions", John Wiley & Sons Inc, 1992, vol 42, chapter 2, David L Hughes). The reaction is preferably effected in the presence of a base (as defined hereinbefore in process (a)) and advantageously in the presence of an inert solvent or diluent


(as defined hereinbefore in process (a)), advantageously at a temperature in the range, for example 10 to 150°C, conveniently at about 50°C.

(c) Compounds of the formula I and salts thereof wherein at least one R^2 is R^5X^1 wherein R^5 is as defined hereinbefore and X^1 is -O-, -S-, -OC(O)- or -NR¹⁰- (wherein R^{10} represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) may be prepared by the reaction of a compound of the formula VII:



(VII)

with a compound of the formula VIII:



(VIII)

(wherein L^1 , R^1 , R^2 , R^5 , ring C, Z, n and s are all as hereinbefore defined and X^1 is as hereinbefore defined in this section). The reaction may conveniently be effected in the presence of a base (as defined hereinbefore in process (a)) and advantageously in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)), advantageously at a temperature in the range, for example 10 to 150°C, conveniently at about 100°C.

(d) Compounds of the formula I and salts thereof wherein at least one R^2 is R^5X^1 wherein X^1 is as defined hereinbefore and R^5 is C₁₋₅alkylR¹¹³, wherein R¹¹³ is selected from one of the following six groups:

1) X¹⁹C₁₋₃alkyl (wherein X¹⁹ represents -O-, -S-, -SO₂-, -NR¹¹⁴C(O)- or -NR¹¹⁵SO₂- (wherein R¹¹⁴ and R¹¹⁵ which may be the same or different are each hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl);

2) NR¹¹⁶R¹¹⁷ (wherein R¹¹⁶ and R¹¹⁷ which may be the same or different are each hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl);

3) $X^{20}C_{1-5}alkylX^5R^{22}$ (wherein X^{20} represents -O-, -S-, -SO₂-, -NR¹¹⁸C(O)-, -NR¹¹⁹SO₂- or -NR¹²⁰- (wherein R¹¹⁸, R¹¹⁹, and R¹²⁰ which may be the same or different are each hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and X⁵ and R²² are as defined hereinbefore);

4) R²⁸ (wherein R²⁸ is as defined hereinbefore);

5) $X^{21}R^{29}$ (wherein X^{21} represents -O-, -S-, -SO₂-, -NR¹²¹C(O)-, -NR¹²²SO₂-, or -NR¹²³- (wherein R¹²¹, R¹²², and R¹²³ which may be the same or different are each hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore); and

6) $X^{22}C_{1-3}alkylR^{29}$ (wherein X^{22} represents -O-, -S-, -SO₂-, -NR¹²⁴C(O)-, -NR¹²⁵SO₂- or -NR¹²⁶- (wherein R¹²⁴, R¹²⁵ and R¹²⁶ each independently represents hydrogen, C₁₋₃alkyl or

C₁₋₃alkoxyC₂₋₃alkyl) and R²⁹ is as defined hereinbefore);


and additionally R¹¹³ may be selected from the following three groups:

7) R²⁹ (wherein R²⁹ is as defined hereinbefore);

8) $X^{22}C_{1-4}alkylR^{28}$ (wherein X^{22} and R²⁸ are as defined hereinbefore); and

9) $R^{54}(C_{1-4}alkyl)_q(X^9)_rR^{55}$ (wherein q, r, X⁹, R⁵⁴ and R⁵⁵ are as defined hereinbefore);

may be prepared by reacting a compound of the formula IX:



(IX)

(wherein L¹, X¹, R¹, R², ring C, Z, n and s are as hereinbefore defined) with a compound of the formula X:



(wherein R¹¹³ is as defined hereinbefore) to give a compound of the formula I or salt thereof.

The reaction may conveniently be effected in the presence of a base (as defined hereinbefore

in process (a)) and advantageously in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)), and at a temperature in the range, for example 0 to 150°C, conveniently at about 50°C.

Processes (a) and (b) are preferred over processes (c) and (d).

5 Process (a) is preferred over processes (b), (c) and (d).

(e) The production of those compounds of the formula I and salts thereof wherein one or more of the substituents $(R^2)_m$ is represented by $-NR^{127}R^{128}$, where one (and the other is hydrogen) or both of R^{127} and R^{128} are C_{1-3} alkyl, may be effected by the reaction of compounds of formula I wherein the substituent $(R^2)_m$ is an amino group and an alkylating agent,


10 preferably in the presence of a base as defined hereinbefore. Such alkylating agents are C_{1-3} alkyl moieties bearing a displaceable moiety as defined hereinbefore such as C_{1-3} alkyl halides for example C_{1-3} alkyl chloride, bromide or iodide. The reaction is preferably effected in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)) and at a temperature in the range, for example, 10 to 100°C, conveniently at about ambient

15 temperature. The production of compounds of formula I and salts thereof wherein one or more of the substituents R^2 is an amino group may be effected by the reduction of a corresponding compound of formula I wherein the substituent(s) at the corresponding position(s) of the quinazoline group is/are a nitro group(s). The reduction may conveniently be effected as described in process (i) hereinafter. The production of a compound of formula I and salts thereof wherein the substituent(s) at the corresponding position(s) of the quinazoline group is/are a nitro group(s) may be effected by the processes described hereinbefore and hereinafter in processes (a-d) and (i-v) using a compound selected from the compounds of the formulae (I-XXII) in which the substituent(s) at the corresponding position(s) of the quinazoline group is/are a nitro group(s).

25 (f) Compounds of the formula I and salts thereof wherein X^1 is $-SO-$ or $-SO_2-$ may be prepared by oxidation from the corresponding compound in which X^1 is $-S-$ or $-SO-$ (when X^1 is $-SO_2-$ is required in the final product). Conventional oxidation conditions and reagents for such reactions are well known to the skilled chemist.

Synthesis of Intermediates

30 (i) The compounds of formula III and salts thereof in which L^1 is halogeno may for example be prepared by halogenating a compound of the formula XI:




(XI)

wherein R^2 and m are as hereinbefore defined).

Convenient halogenating agents include inorganic acid halides, for example thionyl chloride, phosphorus(III)chloride, phosphorus(V)oxychloride and phosphorus(V)chloride.

The halogenation reaction may be effected in the presence of an inert solvent or diluent such as for example a halogenated solvent such as methylene chloride, trichloromethane or carbon tetrachloride, or an aromatic hydrocarbon solvent such as benzene or toluene, or the reaction may be effected without the presence of a solvent. The reaction is conveniently effected at a temperature in the range, for example 10 to 150°C, preferably in the range 40 to 100°C.

The compounds of formula XI and salts thereof may, for example, be prepared by reacting a compound of the formula XII:




(XII)

(wherein R^2 , s and L^1 are as hereinbefore defined) with a compound of the formula VIII as hereinbefore defined. The reaction may conveniently be effected in the presence of a base (as defined hereinbefore in process (a)) and advantageously in the presence of an inert solvent or diluent (as defined hereinbefore in process (a)), advantageously at a temperature in the range, for example 10 to 150°C, conveniently at about 100°C.

Compounds of formula XI and salts thereof wherein at least one R^2 is R^5X^1 and wherein X^1 is -O-, -S-, -SO-, -SO₂-, -C(O)-, -C(O)NR⁷-, -SO₂NR⁸- or -NR¹⁰- (wherein R^7 , R^8


and R^{10} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl), may for example also be prepared by the reaction of a compound of the formula XIII:



(XIII)

(wherein R^2 and s are as hereinbefore defined and X^1 is as hereinbefore defined in this section) with a compound of the formula VI as hereinbefore defined. The reaction may for example be effected as described for process (b) hereinbefore. The pivaloyloxymethyl group can then be cleaved by reacting the product with a base such as, for example, aqueous ammonia, triethylamine in water, an alkali metal or alkaline earth metal hydroxide or alkoxide, preferably aqueous ammonia, aqueous sodium hydroxide or aqueous potassium hydroxide, in a polar protic solvent such as an alcohol, for example methanol or ethanol. The reaction is conveniently effected at a temperature in the range 20 to 100°C, preferably in the range 20 to 50°C.

The compounds of formula XI and salts thereof may also be prepared by cyclising a compound of the formula XIV:




(XIV)

(wherein R^2 and m , are as hereinbefore defined, and A^1 is an hydroxy, alkoxy (preferably C_{1-3} alkoxy) or amino group) whereby to form a compound of formula XI or salt thereof. The cyclisation may be effected by reacting a compound of the formula XIV, where A^1 is an hydroxy or alkoxy group, with formamide or an equivalent thereof effective to cause cyclisation whereby a compound of formula XI or salt thereof is obtained, such as [3-

(dimethylamino)-2-azaprop-2-enylidene]dimethylammonium chloride. The cyclisation is conveniently effected in the presence of formamide as solvent or in the presence of an inert solvent or diluent such as an ether for example 1,4-dioxan. The cyclisation is conveniently effected at an elevated temperature, preferably in the range 80 to 200°C. The compounds of formula XI may also be prepared by cyclising a compound of the formula XIV, where A¹ is an amino group, with formic acid or an equivalent thereof effective to cause cyclisation whereby a compound of formula XI or salt thereof is obtained. Equivalents of formic acid effective to cause cyclisation include for example a tri-C₁₋₄alkoxymethane, for example triethoxymethane and trimethoxymethane. The cyclisation is conveniently effected in the presence of a catalytic amount of an anhydrous acid, such as a sulphonic acid for example p-toluenesulphonic acid, and in the presence of an inert solvent or diluent such as for example a halogenated solvent such as methylene chloride, trichloromethane or carbon tetrachloride, an ether such as diethyl ether or tetrahydrofuran, or an aromatic hydrocarbon solvent such as toluene. The cyclisation is conveniently effected at a temperature in the range, for example 10 to 100°C, preferably in the range 20 to 50°C.

Compounds of formula XIV and salts thereof may for example be prepared by the reduction of the nitro group in a compound of the formula XV:




(XV)

(wherein R², m and A¹ are as hereinbefore defined) to yield a compound of formula XIV as hereinbefore defined. The reduction of the nitro group may conveniently be effected by any of the procedures known for such a transformation. The reduction may be carried out, for example, by stirring a solution of the nitro compound under hydrogen at 1 to 4 atmospheres pressure in the presence of an inert solvent or diluent as defined hereinbefore in the presence of a metal effective to catalyse hydrogenation reactions such as palladium or platinum. A

further reducing agent is, for example, an activated metal such as activated iron (produced for example by washing iron powder with a dilute solution of an acid such as hydrochloric acid). Thus, for example, the reduction may be effected by heating the nitro compound under hydrogen at 2 atmospheres pressure in the presence of the activated metal and a solvent or diluent such as a mixture of water and alcohol, for example methanol or ethanol, at a temperature in the range, for example 50 to 150°C, conveniently at about 70°C.


Compounds of the formula XV and salts thereof may for example be prepared by the reaction of a compound of the formula XVI:



(XVI)

(wherein R^2 , s , L^1 and A^1 are as hereinbefore defined) with a compound of the formula VIII as hereinbefore defined to give a compound of the formula XV. The reaction of the compounds of formulae XVI and VIII is conveniently effected under conditions as described for process (c) hereinbefore.

Compounds of formula XV and salts thereof wherein at least one R^2 is R^5X^1 and wherein X^1 is $-\text{O}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^7-$, $-\text{SO}_2\text{NR}^8-$ or $-\text{NR}^{10}-$ (wherein R^7 , R^8 and R^{10} each independently represents hydrogen, $\text{C}_{1-3}\text{alkyl}$ or $\text{C}_{1-3}\text{alkoxyC}_{2-3}\text{alkyl}$), may for example also be prepared by the reaction of a compound of the formula XVII:



(XVII)


(wherein R^2 , s and A^1 are as hereinbefore defined and X^1 is as hereinbefore defined in this section) with a compound of the formula VI as hereinbefore defined to yield a compound of formula XV as hereinbefore defined. The reaction of the compounds of formulae XVII and VI is conveniently effected under conditions as described for process (b) hereinbefore.

5 The compounds of formula III and salts thereof wherein at least one R^2 is R^5X^1 and wherein X^1 is $-CH_2-$ may be prepared for example as described above from a compound of the formula XV (in which R^2 is $-CH_3$) or XIII (in which HX^1- is $-CH_3$), by radical bromination or chlorination to give a $-CH_2Br$ or $-CH_2Cl$ group which may then be reacted with a compound of the formula R^5-H under standard conditions for such substitution reactions.

10 The compounds of formula III and salts thereof wherein at least one R^2 is R^5X^1 and wherein X^1 is a direct bond may be prepared for example as described above from a compound of the formula XI, wherein the R^5 group is already present in the intermediate compounds (for example in a compound of the formula XV) used to prepare the compound of formula XI.

15 The compounds of formula III and salts thereof wherein at least one R^2 is R^5X^1 and wherein X^1 is $-NR^6C(O)-$ or $-NR^9SO_2-$ may be prepared for example from a compound of the formula XIII in which HX^1- is an $-NHR^6-$ or $-NHR^9-$ group (prepared for example from an amino group (later functionalised if necessary) by reduction of a nitro group) which is reacted with an acid chloride or sulfonyl chloride compound of the formula R^5COCl or R^5SO_2Cl .

20 The compounds of formula III and salts thereof wherein at least one R^2 is R^5X^1 and wherein X^1 is $-O-$, $-S-$, $-SO_2-$, $-OC(O)-$, $-C(O)NR^7-$, $-SO_2NR^8-$ or $-NR^{10}-$ (wherein R^7 , R^8 and R^{10} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl), may also be prepared for example by reacting a compound of the formula XVIII:




(XVIII)

(wherein R^2 and s are as hereinbefore defined, X^1 is as hereinbefore defined in this section and L^2 represents a displaceable protecting moiety) with a compound of the formula VI as hereinbefore defined, whereby to obtain a compound of formula III in which L^1 is represented by L^2 .

5 A compound of formula XVIII is conveniently used in which L^2 represents a phenoxy group which may if desired carry up to 5 substituents, preferably up to 2 substituents, selected from halogeno, nitro and cyano. The reaction may be conveniently effected under conditions as described for process (b) hereinbefore.

The compounds of formula XVIII and salts thereof may for example be prepared by
10 deprotecting a compound of the formula XIX:



(XIX)

15 (wherein R^2 , s and L^2 are as hereinbefore defined, P^1 is a protecting group and X^1 is as hereinbefore defined in the section describing compounds of the formula XVIII). The choice of protecting group P^1 is within the standard knowledge of an organic chemist, for example those included in standard texts such as "Protective Groups in Organic Synthesis" T.W.

20 Greene and R.G.M. Wuts, 2nd Ed. Wiley 1991, including N-sulphonyl derivatives (for example, p-toluenesulphonyl), carbamates (for example, t-butyl carbonyl), N-alkyl derivatives (for example, 2-chloroethyl, benzyl) and amino acetal derivatives (for example benzyloxymethyl). The removal of such a protecting group may be effected by any of the procedures known for such a transformation, including those reaction conditions indicated in
25 standard texts such as that indicated hereinbefore, or by a related procedure. Deprotection may be effected by techniques well known in the literature, for example where P^1 represents a benzyl group deprotection may be effected by hydrogenolysis or by treatment with trifluoroacetic acid.


One compound of formula III may if desired be converted into another compound of formula III in which the moiety L^1 is different. Thus for example a compound of formula III in which L^1 is other than halogeno, for example optionally substituted phenoxy, may be converted to a compound of formula III in which L^1 is halogeno by hydrolysis of a compound of formula III (in which L^1 is other than halogeno) to yield a compound of formula XI as hereinbefore defined, followed by introduction of halide to the compound of formula XI, thus obtained as hereinbefore defined, to yield a compound of formula III in which L^1 represents halogen.

(ii) Compounds of formula IV and salts thereof in which ring C is an indolyl may be prepared by any of the methods known in the art, such as for example those described in "Indoles Part I", "Indoles Part II", 1972 John Wiley & Sons Ltd and "Indoles Part III" 1979, John Wiley & Sons Ltd, edited by W. J. Houlihan.

Examples of the preparation of indoles are given in the Examples hereinafter, such as Examples 48, 237, 242, 250 and 291.

Compounds of formula IV and salts thereof in which ring C is a quinoliny1 may be prepared by any of the methods known in the art, such as for example those described in "The Chemistry of Heterocyclic Compounds: Quinolines Parts I, II and III", 1982 (Interscience publications) John Wiley & Sons Ltd, edited by G. Jones, and in "Comprehensive Heterocyclic Chemistry Vol II by A. R. Katritzky", 1984 Pergamon Press, edited by A. J. Boulton and A McKillop.

(iii) Compounds of formula V as hereinbefore defined and salts thereof may be made by deprotecting the compound of formula XX:




(XX)

(wherein ring C, Z, R¹, R², P¹, n and s are as hereinbefore defined and X¹ is as hereinbefore defined in the section describing compounds of the formula V) by a process for example as described in (i) above.

Compounds of the formula XX and salts thereof may be made by reacting compounds of the formulae XIX and IV as hereinbefore defined, under the conditions described in (a) hereinbefore, to give a compound of the formula XX or salt thereof.

(iv) Compounds of the formula VII and salts thereof may be made by reacting a compound of the formula XXI:



(XXI)

(wherein R², s and each L¹ are as hereinbefore defined and the L¹ in the 4-position and the other L¹ in a further position on the quinazoline ring may be the same or different) with a compound of the formula IV as hereinbefore defined, the reaction for example being effected by a process as described in (a) above.

(v) Compounds of formula IX as defined hereinbefore and salts thereof may for example be made by the reaction of compounds of formula V as defined hereinbefore with compounds of the formula XXII:



(wherein L¹ is as hereinbefore defined) to give compounds of formula IX or salts thereof. The reaction may be effected for example by a process as described in (b) above.

(vi) Intermediate compounds wherein X¹ is -SO- or -SO₂- may be prepared by oxidation from the corresponding compound in which X¹ is -S- or -SO- (when X¹ is -SO₂- is required in the final product). Conventional oxidation conditions and reagents for such reactions are well known to the skilled chemist.

When a pharmaceutically acceptable salt of a compound of the formula I is required, it may be obtained, for example, by reaction of said compound with, for example, an acid using a conventional procedure, the acid having a pharmaceutically acceptable anion.

Many of the intermediates defined herein, for example, those of the formulae V, VII, IX and XX are novel and these are provided as a further feature of the invention. The preparation of these compounds is as described herein and/or is by methods well known to persons skilled in the art of organic chemistry.

The identification of compounds which potentially inhibit the tyrosine kinase activity associated with VEGF receptors such as Flt and/or KDR and which inhibit angiogenesis and/or increased vascular permeability is desirable and is the subject of the present invention. These properties may be assessed, for example, using one or more of the procedures set out below:

(a) In Vitro Receptor Tyrosine Kinase Inhibition Test

This assay determines the ability of a test compound to inhibit tyrosine kinase activity. DNA encoding VEGF, FGF or EGF receptor cytoplasmic domains may be obtained by total gene synthesis (Edwards M, International Biotechnology Lab 5(3), 19-25, 1987) or by cloning. These may then be expressed in a suitable expression system to obtain polypeptide with tyrosine kinase activity. For example VEGF, FGF and EGF receptor cytoplasmic domains, which were obtained by expression of recombinant protein in insect cells, were found to display intrinsic tyrosine kinase activity. In the case of the VEGF receptor Flt (Genbank accession number X51602), a 1.7kb DNA fragment encoding most of the cytoplasmic domain, commencing with methionine 783 and including the termination codon, described by Shibuya et al (Oncogene, 1990, 5: 519-524), was isolated from cDNA and cloned into a baculovirus transplacement vector (for example pAcYM1 (see The Baculovirus Expression System: A Laboratory Guide, L.A. King and R. D. Possee, Chapman and Hall, 1992) or pAc360 or pBlueBacHis (available from Invitrogen Corporation)). This recombinant construct was co-transfected into insect cells (for example *Spodoptera frugiperda* 21(Sf21)) with viral DNA (eg Pharmingen BaculoGold) to prepare recombinant baculovirus. (Details of the methods for the assembly of recombinant DNA molecules and the preparation and use of recombinant baculovirus can be found in standard texts for example Sambrook et al, 1989, Molecular cloning - A Laboratory Manual, 2nd edition, Cold Spring Harbour Laboratory Press and O'Reilly et al, 1992, Baculovirus Expression Vectors - A Laboratory Manual, W. H.

Freeman and Co, New York). For other tyrosine kinases for use in assays, cytoplasmic fragments starting from methionine 806 (KDR, Genbank accession number L04947), methionine 668 (EGF receptor, Genbank accession number X00588) and methionine 399 (FGF R1 receptor, Genbank accession number X51803) may be cloned and expressed in a similar manner.

For expression of cFIt tyrosine kinase activity, Sf21 cells were infected with plaque-pure cFIt recombinant virus at a multiplicity of infection of 3 and harvested 48 hours later. Harvested cells were washed with ice cold phosphate buffered saline solution (PBS) (10mM sodium phosphate pH7.4, 138mM sodium chloride, 2.7mM potassium chloride) then resuspended in ice cold HNTG/PMSF (20mM Hepes pH7.5, 150mM sodium chloride, 10% v/v glycerol, 1% v/v Triton X100, 1.5mM magnesium chloride, 1mM ethylene glycol-bis(β-aminoethyl ether) N,N,N',N'-tetraacetic acid (EGTA), 1mM PMSF (phenylmethylsulphonyl fluoride); the PMSF is added just before use from a freshly-prepared 100mM solution in methanol) using 1ml HNTG/PMSF per 10 million cells. The suspension was centrifuged for 10 minutes at 13,000 rpm at 4°C, the supernatant (enzyme stock) was removed and stored in aliquots at -70°C. Each new batch of stock enzyme was titrated in the assay by dilution with enzyme diluent (100mM Hepes pH 7.4, 0.2mM sodium orthovanadate, 0.1% v/v Triton X100, 0.2mM dithiothreitol). For a typical batch, stock enzyme is diluted 1 in 2000 with enzyme diluent and 50μl of dilute enzyme is used for each assay well.

A stock of substrate solution was prepared from a random copolymer containing tyrosine, for example Poly (Glu, Ala, Tyr) 6:3:1 (Sigma P3899), stored as 1 mg/ml stock in PBS at -20°C and diluted 1 in 500 with PBS for plate coating.

On the day before the assay 100μl of diluted substrate solution was dispensed into all wells of assay plates (Nunc maxisorp 96-well immunoplates) which were sealed and left overnight at 4°C.

On the day of the assay the substrate solution was discarded and the assay plate wells were washed once with PBST (PBS containing 0.05% v/v Tween 20) and once with 50mM Hepes pH7.4.

Test compounds were diluted with 10% dimethylsulphoxide (DMSO) and 25μl of diluted compound was transferred to wells in the washed assay plates. "Total" control wells contained 10% DMSO instead of compound. Twenty five microlitres of 40mM manganese(II)chloride containing 8μM adenosine-5'-triphosphate (ATP) was added to all test

wells except "blank" control wells which contained manganese(II)chloride without ATP. To start the reactions 50µl of freshly diluted enzyme was added to each well and the plates were incubated at room temperature for 20 minutes. The liquid was then discarded and the wells were washed twice with PBST. One hundred microlitres of mouse IgG anti-phosphotyrosine antibody (Upstate Biotechnology Inc. product 05-321), diluted 1 in 6000 with PBST containing 0.5% w/v bovine serum albumin (BSA), was added to each well and the plates were incubated for 1 hour at room temperature before discarding the liquid and washing the wells twice with PBST. One hundred microlitres of horse radish peroxidase (HRP)-linked sheep anti-mouse Ig antibody (Amersham product NXA 931), diluted 1 in 500 with PBST containing 0.5% w/v BSA, was added and the plates were incubated for 1 hour at room temperature before discarding the liquid and washing the wells twice with PBST. One hundred microlitres of 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) solution, freshly prepared using one 50mg ABTS tablet (Boehringer 1204 521) in 50ml freshly prepared 50mM phosphate-citrate buffer pH5.0 + 0.03% sodium perborate (made with 1 phosphate citrate buffer with sodium perborate (PCSB) capsule (Sigma P4922) per 100ml distilled water), was added to each well. Plates were then incubated for 20-60 minutes at room temperature until the optical density value of the "total" control wells, measured at 405nm using a plate reading spectrophotometer, was approximately 1.0. "Blank" (no ATP) and "total" (no compound) control values were used to determine the dilution range of test compound which gave 50% inhibition of enzyme activity.

(b) In Vitro HUVEC Proliferation Assay

This assay determines the ability of a test compound to inhibit the growth factor-stimulated proliferation of human umbilical vein endothelial cells (HUVEC).

HUVEC cells were isolated in MCDB 131 (Gibco BRL) + 7.5% v/v foetal calf serum (FCS) and were plated out (at passage 2 to 8), in MCDB 131 + 2% v/v FCS + 3µg/ml heparin + 1µg/ml hydrocortisone, at a concentration of 1000 cells/well in 96 well plates. After a minimum of 4 hours they were dosed with the appropriate growth factor (i.e. VEGF 3ng/ml, EGF 3ng/ml or b-FGF 0.3ng/ml) and compound. The cultures were then incubated for 4 days at 37°C with 7.5% CO₂. On day 4 the cultures were pulsed with 1µCi/well of tritiated-thymidine (Amersham product TRA 61) and incubated for 4 hours. The cells were harvested using a 96-well plate harvester (Tomtek) and then assayed for incorporation of

tritium with a Beta plate counter. Incorporation of radioactivity into cells, expressed as cpm, was used to measure inhibition of growth factor-stimulated cell proliferation by compounds.

(c) In Vivo Solid Tumour Disease Model

5 This test measures the capacity of compounds to inhibit solid tumour growth.

CaLu-6 tumour xenografts were established in the flank of female athymic Swiss *nu/nu* mice, by subcutaneous injection of 1×10^6 CaLu-6 cells/mouse in 100 μ l of a 50% (v/v) solution of Matrigel in serum free culture medium. Ten days after cellular implant, mice were allocated to groups of 8-10, so as to achieve comparable group mean volumes. Tumours were
10 measured using vernier calipers and volumes were calculated as: $(l \times w) \times \sqrt{(l \times w) \times (\pi/6)}$, where l is the longest diameter and w the diameter perpendicular to the longest. Test compounds were administered orally once daily for a minimum of 21 days, and control animals received compound diluent. Tumours were measured twice weekly. The level of growth inhibition was calculated by comparison of the mean tumour volume of the control
15 group versus the treatment group using a Student T test and/or a Mann-Whitney Rank Sum Test. The inhibitory effect of compound treatment was considered significant when $p < 0.05$.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the formula I as defined hereinbefore or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable
20 excipient or carrier.

The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) for example as a sterile solution, suspension or emulsion, for topical administration for example as an ointment or cream or for rectal administration for example as
25 a suppository. In general the above compositions may be prepared in a conventional manner using conventional excipients.

The compositions of the present invention are advantageously presented in unit dosage form. The compound will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000mg per square metre body area of the animal, i.e.
30 approximately 0.1-100mg/kg. A unit dose in the range, for example, 1-100mg/kg, preferably 1-50mg/kg is envisaged and this normally provides a therapeutically-effective dose. A unit

dose form such as a tablet or capsule will usually contain, for example 1-250mg of active ingredient.

According to a further aspect of the present invention there is provided a compound of the formula I or a pharmaceutically acceptable salt thereof as defined hereinbefore for use in a method of treatment of the human or animal body by therapy.

We have found that compounds of the present invention inhibit VEGF receptor tyrosine kinase activity and are therefore of interest for their antiangiogenic effects and/or their ability to cause a reduction in vascular permeability.

A further feature of the present invention is a compound of formula I, or a pharmaceutically acceptable salt thereof, for use as a medicament, conveniently a compound of formula I, or a pharmaceutically acceptable salt thereof, for use as a medicament for producing an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human being.

Thus according to a further aspect of the invention there is provided the use of a compound of the formula I, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human being.

According to a further feature of the invention there is provided a method for producing an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof as defined hereinbefore.

As stated above the size of the dose required for the therapeutic or prophylactic treatment of a particular disease state will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. Preferably a daily dose in the range of 1-50mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

The antiangiogenic and/or vascular permeability reducing treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may

be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. In the field of medical oncology it is normal practice to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such conjoint treatment in addition to the
5 antiangiogenic and/or vascular permeability reducing treatment defined hereinbefore may be: surgery, radiotherapy or chemotherapy. Such chemotherapy may cover three main categories of therapeutic agent:

(i) other antiangiogenic agents that work by different mechanisms from those defined hereinbefore (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function, angiostatin, razoxin,
10 thalidomide), and including vascular targeting agents (for example combretastatin phosphate and the vascular damaging agents described in International Patent Application Publication No. WO 99/02166 the entire disclosure of which document is incorporated herein by reference, (for example N-acetylcolchicol-O-phosphate));

(ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene,
15 droloxifene, idoxifene), progestogens (for example megestrol acetate), aromatase inhibitors (for example anastrozole, letrozole, vorazole, exemestane), antiprogestogens, antiandrogens (for example flutamide, nilutamide, bicalutamide, cyproterone acetate), LHRH agonists and antagonists (for example goserelin acetate, luprolide), inhibitors of testosterone 5α -dihydroreductase (for example finasteride), anti-invasion agents (for example

20 metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function) and inhibitors of growth factor function, (such growth factors include for example platelet derived growth factor and hepatocyte growth factor such inhibitors include growth factor antibodies, growth factor receptor antibodies, tyrosine kinase inhibitors and serine/threonine kinase inhibitors); and

25 (iii) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as antimetabolites (for example antifolates like methotrexate, fluoropyrimidines like 5-fluorouracil, purine and adenosine analogues, cytosine arabinoside); antitumour antibiotics (for example anthracyclines like doxorubicin, daunomycin, epirubicin and idarubicin, mitomycin-C, dactinomycin, mithramycin); platinum derivatives (for example
30 cisplatin, carboplatin); alkylating agents (for example nitrogen mustard, melphalan, chlorambucil, busulphan, cyclophosphamide, ifosfamide, nitrosoureas, thiotepa); antimitotic agents (for example vinca alkaloids like vincristine and taxoids like taxol, taxotere);

topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan, and also irinotecan); also enzymes (for example asparaginase); and thymidylate synthase inhibitors (for example raltitrexed); and additional types of chemotherapeutic agent include:

- 5 (iv) biological response modifiers (for example interferon); and
(v) antibodies (for example edrecolomab).

For example such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of a compound of formula I as defined hereinbefore, and a vascular targeting agent described in WO 99/02166 such as N-acetylcolchicol-O-phosphate
10 (Exampe 1 of WO 99/02166).

As stated above the compounds defined in the present invention are of interest for their antiangiogenic and/or vascular permeability reducing effects. Such compounds of the invention are expected to be useful in a wide range of disease states including cancer, diabetes, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic
15 nephropathies, atheroma, arterial restenosis, autoimmune diseases, acute inflammation, excessive scar formation and adhesions, endometriosis, dysfunctional uterine bleeding and ocular diseases with retinal vessel proliferation. In particular such compounds of the invention are expected to slow advantageously the growth of primary and recurrent solid tumours of, for example, the colon, breast, prostate, lungs and skin. More particularly such
20 compounds of the invention are expected to inhibit the growth of those primary and recurrent solid tumours which are associated with VEGF, especially those tumours which are significantly dependent on VEGF for their growth and spread, including for example, certain tumours of the colon, breast, prostate, lung, vulva and skin.

In addition to their use in therapeutic medicine, the compounds of formula I and
25 their pharmaceutically acceptable salts are also useful as pharmacological tools in the development and standardisation of in vitro and in vivo test systems for the evaluation of the effects of inhibitors of VEGF receptor tyrosine kinase activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

It is to be understood that where the term "ether" is used anywhere in this
30 specification it refers to diethyl ether.

The invention will now be illustrated in the following non-limiting Examples in which, unless otherwise stated:-

(i) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids such as drying agents by
5 filtration;

(ii) operations were carried out at ambient temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as argon;

(iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (MPLC) were performed on Merck Kieselgel silica (Art. 9385) or Merck
10 Lichroprep RP-18 (Art. 9303) reversed-phase silica obtained from E. Merck, Darmstadt, Germany;

(iv) yields are given for illustration only and are not necessarily the maximum attainable;

(v) melting points are uncorrected and were determined using a Mettler SP62
15 automatic melting point apparatus, an oil-bath apparatus or a Koffler hot plate apparatus.

(vi) the structures of the end-products of the formula I were confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic resonance chemical shift values were measured on the delta scale and peak multiplicities are shown as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad; q, quartet, quin,
20 quintet;

(vii) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), high-performance liquid chromatography (HPLC), infra-red (IR) or NMR analysis;

(viii) HPLC were run under 2 different conditions:

- 25 1) on a TSK Gel super ODS 2µM 4.6mm x 5cm column, eluting with a gradient of methanol in water (containing 1% acetic acid) 20 to 100% in 5 minutes. Flow rate 1.4 ml/minute. Detection: U.V. at 254 nm and light scattering detections;
- 2) on a TSK Gel super ODS 2µM 4.6mm x 5cm column, eluting with a gradient of methanol in water (containing 1% acetic acid) 0 to 100% in 7 minutes. Flow rate 1.4 ml/minute.
- 30 Detection: U.V. at 254 nm and light scattering detections.

(ix) petroleum ether refers to that fraction boiling between 40-60°C

(x) the following abbreviations have been used:-

DMF *N,N*-dimethylformamide

DMSO dimethylsulphoxide

TFA trifluoroacetic acid

5 NMP 1-methyl-2-pyrrolidinone

THF tetrahydrofuran

HMDS 1,1,1,3,3,3-hexamethyldisilazane.

HPLC RT HPLC retention time

DEAD diethyl azodicarboxylate

10 DMA dimethylacetamide

DMAP 4-dimethylaminopyridine

Example 1

A mixture of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (225mg,
15 0.67mmol), potassium carbonate (106mg, 0.77mmol) and 6-hydroxyquinoline (112mg,
0.77mmol) in DMF (7.5ml) was stirred at 100°C for 5 hours and allowed to cool to ambient
temperature. The reaction mixture was treated with 1M aqueous sodium hydroxide solution
(40ml) and stirred at ambient temperature for a few minutes. The crude solid was collected by
filtration and washed with water. The resultant solid was dissolved in dichloromethane (2ml)
20 and filtered through phase separating paper. The filtrate was evaporated under vacuum and
the residue was triturated with ether, collected by filtration and dried to give **6-methoxy-7-(3-
morpholinopropoxy)-4-(quinolin-6-yloxy)quinazoline** (163mg, 55%).

¹H NMR Spectrum: (DMSO-d₆) 1.98(m, 2H); 2.40(m, 4H); 2.48(t, 2H); 3.59(m, 4H); 4.00(s,
3H); 4.25(t, 2H); 7.40(s, 1H); 7.58(m, 1H); 7.62(s, 1H); 7.74(dd, 1H); 7.92(d, 1H); 8.10(d,
25 1H); 8.38(d, 1H); 8.55(s, 1H); 8.92(m, 1H)

MS (ESI): 447 (MH)⁺

Elemental analysis:	Found	C 65.9	H 5.7	N 12.4
C ₂₅ H ₂₆ N ₄ O ₄ 0.5H ₂ O	Requires	C 65.9	H 6.0	N 12.3%

30 The starting material was prepared as follows:

A mixture of 2-amino-4-benzyloxy-5-methoxybenzamide (10g, 0.04mol), (J. Med.
Chem. 1977, vol 20, 146-149), and Gold's reagent (7.4g, 0.05mol) in dioxane (100ml) was

stirred and heated at reflux for 24 hours. Sodium acetate (3.02g, 0.037mol) and acetic acid (1.65ml, 0.029mol) were added to the reaction mixture and it was heated for a further 3 hours. The volatiles were removed by evaporation, water was added to the residue, the solid was collected by filtration, washed with water and dried. Recrystallisation from acetic acid gave

5 7-benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (8.7g, 84%).

7-Benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (35g, 124mmol) was suspended in thionyl chloride (440ml) and DMF (1.75ml) and heated at reflux for 4 hours. The thionyl chloride was evaporated under vacuum and the residue azeotroped with toluene three times. The residue was dissolved in NMP (250ml) to give a solution of 7-benzyloxy-4-chloro-6-

10 methoxyquinazoline.

Phenol (29.05g, 309mmol) was dissolved in NMP (210ml), sodium hydride (11.025g, 60% dispersion in mineral oil) was added in portions with cooling and the mixture was stirred for 3 hours. The viscous suspension was diluted with NMP (180ml) and stirred overnight. The solution of 7-benzyloxy-4-chloro-6-methoxyquinazoline was added and the suspension

15 stirred at 100°C for 2.5 hours. The suspension was allowed to cool to ambient temperature and poured into water (1.5l) with vigorous stirring. The precipitate was collected by filtration, washed with water and dried under vacuum. The residue was dissolved in dichloromethane, washed with brine and filtered through phase separating paper. The filtrate was evaporated under vacuum then triturated with ether to give 7-benzyloxy-6-methoxy-4-

20 phenoxyquinazoline (87.8g, 83%) as a pale cream solid.

¹H NMR Spectrum: (CDCl₃) 4.09(s, 3H); 5.34(s, 2H); 7.42(m, 12H); 7.63(s, 1H)

MS (ESI): 359 (MH)⁺

7-Benzyloxy-6-methoxy-4-phenoxyquinazoline (36.95g, 105.5mmol) was suspended in TFA (420ml) and heated at reflux for 3 hours. The reaction mixture was allowed to cool and evaporated under vacuum. The residue was stirred mechanically in water then basified

25 with saturated aqueous sodium hydrogen carbonate solution and stirred overnight. The water was decanted and the solid suspended in acetone. After stirring the white solid was collected by filtration, washed with acetone and dried to give 7-hydroxy-6-methoxy-4-

phenoxyquinazoline (26.61g, 96%).

¹H NMR Spectrum: (DMSO-d₆) 3.97(s, 3H); 7.22(s, 1H); 7.30(m, 3H); 7.47(t, 2H); 7.56(s, 1H); 8.47(s, 1H); 10.70(s, 1H)

MS (ESI): 269 (MH)⁺

Morpholine (52.2ml, 600mmol) and 1-bromo-3-chloropropane (30ml, 300mmol) were dissolved in dry toluene (180ml) and heated to 70°C for 3 hours. The solid was removed by filtration and the filtrate evaporated under vacuum. The resulting oil was decanted from the additional solid residue and the oil was vacuum distilled to yield 1-chloro-3-

5 morpholinopropane (37.91g, 77%) as an oil.

¹H NMR Spectrum: (DMSO-d₆) 1.85(m, 2H); 2.30(t, 4H); 2.38(t, 2H); 3.53(t, 4H); 3.65(t, 2H)
MS (ESI): 164 (MH)⁺

7-Hydroxy-6-methoxy-4-phenoxyquinazoline (25.27g, 0.1mol) and 1-chloro-3-morpholinopropane (18.48g, 0.11mol) were taken up in DMF (750ml) and potassium
10 carbonate (39.1g, 0.33mol) was added. The suspension was heated at 90°C for 3 hours then allowed to cool. The suspension was filtered and the volatiles were removed by evaporation. The residue was triturated with ethyl acetate and 6-methoxy-7-(3-morpholinopropoxy)-4-phenoxyquinazoline (31.4g, 84%) was collected by filtration as a yellow crystalline solid.
¹H NMR Spectrum: (DMSO-d₆) 1.97(m, 2H); 2.39(t, 4H); 2.47(t, 2H); 3.58(t, 4H); 3.95(s, 3H);
15 4.23(t, 2H); 7.31(m, 3H); 7.36(s, 1H); 7.49(t, 2H); 7.55(s, 1H); 8.52(s, 1H)
MS (ESI): 396 (MH)⁺

6-Methoxy-7-(3-morpholinopropoxy)-4-phenoxyquinazoline (33.08g, 84mmol) was dissolved in 6M aqueous hydrochloric acid (800ml) and heated at reflux for 1.5 hours. The reaction mixture was decanted and concentrated to 250ml then basified (pH9) with saturated
20 aqueous sodium hydrogen carbonate solution. The aqueous layer was extracted with dichloromethane (4x400ml), the organic layer was separated and filtered through phase separating paper. The solid was triturated with ethyl acetate to give 6-methoxy-7-(3-morpholinopropoxy)-3,4-dihydroquinazolin-4-one (23.9g, 89%) as a white solid.
¹H NMR Spectrum: (DMSO-d₆) 1.91(m, 2H); 2.34(t, 4H); 2.42(t, 2H); 3.56(t, 4H); 3.85(s, 3H);
25 4.12(t, 2H); 7.11(s, 1H); 7.42(s, 1H); 7.96(s, 1H); 12.01(s, 1H)
MS (ESI): 320 (MH)⁺

6-Methoxy-7-(3-morpholinopropoxy)-3,4-dihydroquinazolin-4-one (23.9g, 75mmol) was suspended in thionyl chloride (210ml) and DMF (1.8ml) then heated at reflux for 1.5 hours. The thionyl chloride was removed by evaporation under vacuum and the residue
30 azeotroped with toluene three times. The residue was taken up in water and basified (pH8) with saturated aqueous sodium hydrogen carbonate solution. The aqueous layer was extracted with dichloromethane (4x400ml), the organic layer was washed with water and brine then

dried (MgSO_4). After filtration the organic layer was concentrated under vacuum to give a yellow solid which was triturated with ethyl acetate to give 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (17.39g, 52%) as a pale cream solid.

^1H NMR Spectrum: (CDCl_3) 2.10-2.16(m, 2H); 2.48(br s, 4H); 2.57(t, 2H); 3.73(t, 4H); 4.05(s, 3H); 4.29(t, 2H); 7.36(s, 1H); 7.39(s, 1H); 8.86(s, 1H)

MS-ESI: 337 $[\text{MH}]^+$

Example 2

A mixture of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (225mg, 0.67mmol), (prepared as described for the starting material in Example 1), potassium carbonate (106mg, 0.77mmol) and 7-hydroxyquinoline (112mg, 0.77mmol) in DMF (7.5ml) was stirred at 100°C for 5 hours and allowed to cool to ambient temperature. The reaction mixture was treated with 1M aqueous sodium hydroxide solution (40ml) and stirred at ambient temperature for a few minutes. The crude solid was collected by filtration washing with water. The resultant solid was dissolved in dichloromethane (2ml) and filtered through phase separating paper. The filtrate was evaporated under vacuum to give a solid residue which was triturated with ether, filtered and dried to give **6-methoxy-7-(3-morpholinopropoxy)-4-(quinolin-7-yloxy)quinazoline** (116mg, 39%).

^1H NMR Spectrum: (DMSO-d_6) 1.98(m, 2H); 2.39(m, 4H); 2.48(t, 2H); 3.59(m, 4H); 4.00(s, 3H); 4.25(t, 2H); 7.40(s, 1H); 7.58(m, 2H); 7.62(s, 1H); 7.92(d, 1H); 8.10(d, 1H); 8.44(d, 1H); 8.55(s, 1H); 8.92(m, 1H)

MS (ESI): 447 $(\text{MH})^+$

Elemental analysis:	Found	C 66.6	H 5.7	N 12.4
$\text{C}_{25}\text{H}_{26}\text{N}_4\text{O}_4 \cdot 0.25\text{H}_2\text{O}$	Requires	C 66.6	H 5.9	N 12.4%

Example 3

A mixture of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (225mg, 0.67mmol), (prepared as described for the starting material in Example 1), potassium carbonate (106mg, 0.77mmol) and 1-naphthol (111mg, 0.77mmol) in DMF (7.5ml) was stirred at 100°C for 5 hours then allowed to cool to ambient temperature. The reaction mixture was treated with 1M aqueous sodium hydroxide solution (40ml) and stirred at ambient temperature for a few minutes. The reaction mixture was extracted with ethyl acetate

and the organic extracts were washed with water. The organic extracts were dried (MgSO_4) and the solvent removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (95/5) to give a solid which was triturated with ether, filtered and dried to give **6-methoxy-7-(3-morpholinopropoxy)-4-(1-naphthylloxy)quinazoline** (194mg, 65%).

^1H NMR Spectrum: (DMSO-d_6) 1.98(m, 2H); 2.39(m, 4H); 2.48(t, 2H); 3.59(m, 4H); 4.00(s, 3H); 4.26(t, 2H); 7.40(s, 1H); 7.48(m, 2H); 7.58(m, 2H); 7.74(s, 1H); 7.75(d, 1H); 7.92(d, 1H); 8.03(d, 1H); 8.42(s, 1H)

MS (ESI): 446 (MH)⁺

10	Elemental analysis:	Found	C 69.9	H 6.2	N 9.4
	$\text{C}_{26}\text{H}_{27}\text{N}_3\text{O}_4$	Requires	C 70.1	H 6.1	N 9.4%

Example 4

A mixture of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (225mg, 0.67mmol), (prepared as described for the starting material in Example 1), potassium carbonate (106mg, 0.77mmol) and 7-hydroxy-4-methylquinoline (122mg, 0.77mmol), (Chem. Berich. 1967, 100, 2077), in DMF (7.5ml) was stirred at 100°C for 5 hours then allowed to cool to ambient temperature. The reaction mixture was treated with 1M aqueous sodium hydroxide solution (40ml) and stirred at ambient temperature for a few minutes. The crude solid was collected by filtration washing with water. The resultant solid was dissolved in dichloromethane (2ml) and was filtered through phase separating paper. The filtrate was evaporated under vacuum to give a solid residue which was triturated with ether, filtered and dried to give **6-methoxy-4-(4-methylquinolin-7-yloxy)-7-(3-morpholinopropoxy)quinazoline** (175mg, 57%).

^1H NMR Spectrum: (DMSO-d_6) 1.98(m, 2H); 2.39(m, 4H); 2.48(t, 2H); 2.71(s, 3H); 3.59(m, 4H); 4.00(s, 3H); 4.26(t, 2H); 7.40(s, 1H); 7.41(m, 1H); 7.61(dd, 1H); 7.62(s, 1H); 7.90(d, 1H); 8.20(d, 1H); 8.52(s, 1H); 8.78(d, 1H)

MS (ESI): 461 (MH)⁺

30	Elemental analysis:	Found	C 67.1	H 5.9	N 12.1
	$\text{C}_{26}\text{H}_{28}\text{N}_4\text{O}_4 \cdot 0.2\text{H}_2\text{O}$	Requires	C 67.3	H 6.2	N 12.1%

Example 5

A mixture of 4-chloro-7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxyquinazoline (220mg, 0.57mmol), potassium carbonate (106mg, 0.77mmol) and 7-hydroxyquinoline (111mg, 0.76mmol) in DMF (7.5ml) was stirred at 100°C for 5 hours then allowed to cool to ambient temperature. The reaction mixture was treated with 1M aqueous sodium hydroxide solution (40ml) and stirred at ambient temperature for a few minutes. The crude solid was collected by filtration washing with water. The resultant solid was dissolved in dichloromethane (2ml) and was filtered through phase separating paper. The filtrate was evaporated under vacuum to give a solid residue which was triturated with ether, filtered and dried to give 7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(quinolin-7-yloxy)quinazoline (205mg, 73%).

¹H NMR Spectrum: (DMSO-d₆) 1.98(m, 2H); 2.65(t, 2H); 2.92(m, 4H); 3.10(m, 4H); 4.00(s, 3H); 4.28(t, 2H); 7.42(s, 1H); 7.58(m, 2H); 7.64(s, 1H); 7.92(d, 1H); 8.10(d, 1H); 8.44(d, 1H); 8.55(s, 1H); 8.92(m, 1H)

MS (ESI): 495 (MH)⁺

15	Elemental analysis:	Found	C 60.0	H 5.0	N 11.1
	C ₂₅ H ₂₆ N ₄ O ₅ S 0.25H ₂ O	Requires	C 60.2	H 5.4	N 11.2%

The starting material was prepared as follows:

7-Benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (20.3g, 124mmol), (prepared as described for the starting material in Example 1), was taken up in thionyl chloride (440ml) and DMF (1.75ml) then heated at reflux for 4 hours. The thionyl chloride was evaporated under vacuum and the residue azeotroped with toluene three times to give 7-benzyloxy-4-chloro-6-methoxyquinazoline.

A mixture of the crude 7-benzyloxy-4-chloro-6-methoxyquinazoline, potassium carbonate (50g, 362mmol) and 4-chloro-2-fluorophenol (8.8ml, 83mmol) in DMF (500ml) was stirred at 100°C for 5 hours then allowed to cool to ambient temperature overnight. The reaction mixture was poured into water (2l) and was stirred at ambient temperature for a few minutes. The crude solid was collected by filtration washing with water. The resultant solid was dissolved in dichloromethane and filtered through diatomaceous earth. The filtrate was treated with decolourising charcoal, boiled for a few minutes then filtered through diatomaceous earth. The filtrate was filtered through phase separating paper and then evaporated under vacuum to give a solid residue which was triturated with ether, filtered and

dried to give 7-benzyloxy-4-(4-chloro-2-fluorophenoxy)-6-methoxyquinazoline (23.2g, 76%).

¹H NMR Spectrum: (DMSO-d₆) 3.98(s, 3H); 5.34(s, 2H); 7.42(m, 9H); 7.69(dd, 1H); 8.55(s, 1H)

MS (ESI): 411 (MH)⁺

- 5 7-Benzyloxy-4-(4-chloro-2-fluorophenoxy)-6-methoxyquinazoline (1.4g, 3.4mmol) was suspended in TFA (15ml) and heated at reflux for 3 hours. The reaction mixture was allowed to cool, toluene was added and the volatiles were removed by evaporation under vacuum. The residue was triturated with ether and then acetone. The precipitate was collected by filtration and dried to give 4-(4-chloro-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline (21.8g). This was used without further purification in the next step.

¹H NMR Spectrum: (DMSO-d₆) 3.97(s, 3H); 7.22(s, 1H); 7.39(d, 1H); 7.53(m, 2H); 7.67(dd, 1H); 8.46(s, 1H)

MS (ESI): 321 (MH)⁺

- 15 A mixture of 3-amino-1-propanol (650μl, 8.4mmol) and vinyl sulphone (1g, 8.4mmol) was heated at 110°C for 45 minutes. The mixture was allowed to cool and was purified by column chromatography eluting with methylene chloride/methanol (95/5) to give 3-(1,1-dioxothiomorpholino)-1-propanol (800mg, 90%).

¹H NMR Spectrum: (CDCl₃) 1.7-1.8(m, 2H); 2.73(t, 2H); 3.06(br s, 8H); 3.25(s, 1H); 3.78(t, 2H)

- 20 MS - ESI: 194 [MH]⁺

- 25 4-(4-Chloro-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline (5.0g, 15.6mmol) was suspended in dichloromethane (150ml) and tributylphosphine (11.1ml, 44.6mmol) was added followed by stirring at ambient temperature for 30 minutes. To this mixture was added 3-(1,1-dioxothiomorpholino)-1-propanol (4.2g, 21.8mmol) followed by the addition of 1,1'-(azodicarbonyl)dipiperidine (11.7g, 46.4mmol) in portions. The mixture was stirred at ambient temperature overnight then diluted with ether (300ml) and the precipitate was removed by filtration. The residue was chromatographed on silica eluting with dichloromethane and methanol (95/5). The relevant fractions were combined and evaporated to give a solid which was triturated with ethyl acetate filtered and dried to give 4-(4-chloro-2-fluorophenoxy)-7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxyquinazoline (5.4g, 70%). This was used without further purification in the next step.

¹H NMR Spectrum: (DMSO-d₆) 1.86(m, 2H); 2.65(t, 2H); 2.92(m, 4H); 3.08(m, 4H); 3.97(s,

3H); 4.26(t, 2H); 7.40(m, 1H); 7.42(s, 1H); 7.56(m, 2H); 7.68(dd, 1H); 8.54(s, 1H)

MS (ESI): 496 (MH)⁺

Elemental analysis:	Found	C 52.7	H 4.4	N 8.3
---------------------	-------	--------	-------	-------

C ₂₂ H ₂₃ N ₃ ClFO ₅ S 0.25H ₂ O	Requires	C 52.8	H 4.7	N 8.4%
---	----------	--------	-------	--------

5 4-(4-Chloro-2-fluorophenoxy)-7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxyquinazoline (3.5g, 7mmol) was dissolved in 2M aqueous hydrochloric acid (56ml) and heated at 95°C for 2 hours. The cooled reaction mixture was treated with solid sodium hydrogen carbonate solution to give a thick paste which was diluted with water and filtered. The solid was transferred to a flask and azeotroped with toluene twice to give a dry solid. The
10 solid was flash chromatographed on silica eluting with dichloromethane and methanol (95/5). The relevant fractions were combined and evaporated to give 7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (2.26g, 87%) as a white solid.

MS (ESI): 368 (MH)⁺

15 7-(3-(1,1-Dioxothiomorpholino)propoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (4.2g, 11.4mmol) was suspended in thionyl chloride (45ml) and DMF (0.1ml) then heated at reflux for 2.5 hours. The residue was diluted with toluene, the thionyl chloride was evaporated under vacuum, the residue was then azeotroped with toluene three times. The residue was taken up in water and basified (pH8) with saturated aqueous sodium hydrogen
20 carbonate solution. The aqueous layer was extracted with dichloromethane (x4), the organic layer was washed with water and brine then filtered through phase separating paper. The organic layer was concentrated under vacuum to give an orange solid. The solid was flash chromatographed on silica eluting with dichloromethane and methanol (95/5). The relevant fractions were combined and evaporated to give a solid which was triturated with ether then
25 filtered and dried to give 4-chloro-7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxyquinazoline (2.27g, 52%).

MS (ESI): 386 (MH)⁺

Example 6

30 6,7-Dimethoxy-3,4-dihydroquinazolin-4-one (290mg, 1.4mmol) was suspended in thionyl chloride (5ml) and DMF (2 drops) and heated at reflux for 2 hours. The thionyl chloride was evaporated under vacuum and the residue azeotroped with toluene three times to

give 4-chloro-6,7-dimethoxyquinazoline. A mixture of the crude 4-chloro-6,7-dimethoxyquinazoline, potassium carbonate (970mg, 7mmol) and 7-hydroxyquinoline (235mg, 1.62mmol) in DMF (10ml) was stirred at 100°C for 5 hours and allowed to cool to ambient temperature overnight. The reaction mixture was treated with 1M aqueous sodium hydroxide solution and stirred at ambient temperature for a few minutes. The reaction mixture was extracted with ethyl acetate (x4) and the organic extracts washed with water and brine. The organic extracts were dried (MgSO₄), filtered and the solvent removed under vacuum. The residue was triturated with ethyl acetate and then recrystallised from hot ethyl acetate to give **6,7-dimethoxy-4-(quinolin-7-yloxy)quinazoline** (110mg, 24%) as a white solid.

¹H NMR Spectrum: (DMSO-d₆) 4.00(s, 3H); 4.00(s, 3H); 7.40(s, 1H); 7.59(m, 3H); 7.92(d, 1H); 8.08(d, 1H); 8.42(d, 1H); 8.55(s, 1H); 8.92(dd, 1H)

MS (ESI): 334 (MH)⁺

Elemental analysis:	Found	C 68.2	H 4.3	N 12.5
---------------------	-------	--------	-------	--------

C ₁₉ H ₁₅ N ₃ O ₃	Requires	C 68.5	H 4.5	N 12.6%
---	----------	--------	-------	---------

The starting material was prepared as follows:

A mixture of 4,5-dimethoxyanthranilic acid (19.7g) and formamide (10ml) was stirred and heated at 190°C for 5 hours. The mixture was allowed to cool to approximately 80°C and water (50ml) was added. The mixture was then allowed to stand at ambient temperature for 3 hours. The precipitate was collected by filtration, washed with water and dried to give 6,7-dimethoxy-3,4-dihydroquinazolin-4-one (3.65g).

Example 7

A mixture of (*R,S*)-4-chloro-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline (183mg, 0.57mmol), potassium carbonate (106mg, 0.77mmol) and 7-hydroxyquinoline (111mg, 0.77mmol) in DMF (7ml) was stirred at 100°C for 5 hours and allowed to cool to ambient temperature. The reaction mixture was treated with 1M aqueous sodium hydroxide solution (30ml) and stirred for 10 minutes. The crude solid was collected by filtration washing with water. The resultant solid was dissolved in dichloromethane (2ml) and filtered through phase separating paper. The filtrate was evaporated under vacuum to give a solid residue which was triturated with ether, filtered and dried to give a scalemic

mixture of 6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline (149mg, 61%).

¹H NMR Spectrum: (DMSO-d₆) 1.10(m, 1H); 1.51(m, 1H); 1.64(m, 1H); 1.85(m, 3H); 2.09(m, 1H); 2.15(s, 3H); 2.62(m, 1H); 2.82(m, 1H); 3.99(s, 3H); 4.09(d, 2H); 7.38(s, 1H); 7.55(m, 2H); 7.63(s, 1H); 7.91(d, 1H); 8.10(d, 1H); 8.44(d, 1H); 8.54(s, 1H); 8.93(d, 1H)

MS (ESI): 431 (MH)⁺

Elemental analysis:	Found	C 68.7	H 5.7	N 12.8
C ₂₅ H ₂₆ N ₄ O ₃ 0.3H ₂ O	Requires	C 68.9	H 6.2	N 12.8%

10 The starting material was prepared as follows:

(*R*)-Ethyl nipecotate (5.7g 365mmol), (prepared by resolution of ethyl nipecotate by treatment with L(+)-tartaric acid as described in J. Org. Chem. 1991, (56), 1168), was dissolved in 38.5% aqueous formaldehyde solution (45ml) and formic acid (90ml) and the mixture heated at reflux for 18 hours. The mixture was allowed to cool and added dropwise to cooled saturated aqueous sodium hydrogen carbonate solution. The mixture was adjusted to pH12 by addition of sodium hydroxide and the mixture was extracted with methylene chloride. The organic extract was washed with brine, dried (MgSO₄) and the solvent removed by evaporation to give (*R*)-ethyl 1-methylpiperidine-3-carboxylate (4.51g, 73%) as a colourless oil.

20 MS - ESI: 172 [MH]⁺

A solution of (*R*)-ethyl 1-methylpiperidine-3-carboxylate (5.69g, 33mmol) in ether (20ml) was added dropwise to a stirred solution of lithium aluminium hydride (36.6ml of a 1M solution in THF, 36.6mmol) in ether (85ml) cooled to maintain a reaction temperature of 20°C. The mixture was stirred for 1.5 hours at ambient temperature and then water (1.4ml), 15% aqueous sodium hydroxide solution (1.4ml) and then water (4.3ml) were added. The insolubles were removed by filtration and the volatiles removed from the filtrate by evaporation to give (*R*)-(1-methylpiperidin-3-yl)methanol (4.02g, 94%) as a colourless oil.

¹H NMR Spectrum: (DMSO-d₆) 1.06(q, 1H); 1.51-1.94(m, 5H); 2.04(s, 3H); 2.34(br s, 1H); 2.62(m, 1H); 2.78(d, 1H); 3.49(m, 1H); 3.59(m, 1H)

30 MS - ESI: 130 [MH]⁺

4-(4-Chloro-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline (12.1g, 38mmol), (prepared as described for the starting material in Example 5), was suspended in dichloromethane

(375ml) and treated with triphenylphosphine (29.6g, 113mmol) then stirred at ambient temperature for 30 minutes. (1-Methylpiperidin-3-yl)methanol (8.25g, 63.8mmol) and (*R*)-(1-methylpiperidin-3-yl)methanol (1.46g, 11.3mmol), (CAS 205194-11-2), giving *R*:*S* (57.5:42.5 by chiral HPLC) (9.7g, 75mmol) were dissolved in dichloromethane (75ml) and added to the suspension. Diethyl azodicarboxylate (17.7ml, 75mmol) was added in portions using a syringe pump and the mixture was then allowed to warm to ambient temperature and stirred overnight. The residue was concentrated under vacuum then chromatographed on silica eluting with dichloromethane followed by dichloromethane/methanol /ammonia (93/6/1). The relevant fractions were combined and evaporated to give an oil. The residue was triturated with ether, filtered and dried to give (*R,S*)-4-(4-chloro-2-fluorophenoxy)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline (8.7g, 53%).

¹H NMR Spectrum: (DMSO-d₆) 1.11(m, 1H); 1.50(m, 1H); 1.58-1.98(m, 4H); 2.09(m, 1H); 2.15(s, 3H); 2.62(d, 1H); 2.81(d, 1H); 3.95(s, 3H); 4.09(d, 2H); 7.39(m, 2H); 7.55(m, 2H); 7.67(d, 1H); 8.53(s, 1H)

MS (ESI): 432 (MH)⁺

(*R,S*)-4-(4-Chloro-2-fluorophenoxy)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline (8.7g, 20mmol) was dissolved in 2M aqueous hydrochloric acid (150ml) and heated at reflux for 1.5 hours. The reaction mixture was concentrated then basified (pH9) with saturated aqueous ammonia solution (0.88). The aqueous layer was extracted with dichloromethane (4x400ml) and the organic extracts filtered through phase separating paper then evaporated under vacuum. The solid was triturated with ether to give (*R,S*)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-3,4-dihydroquinazolin-4-one (4.05g, 66%) as a white solid.

¹H NMR Spectrum: (DMSO-d₆) 1.05(m, 1H); 1.40-1.95(m, 5H); 2.02(m, 1H); 2.14(s, 3H); 2.59(d, 1H); 2.78(d, 1H); 3.85(s, 3H); 3.95(d, 2H); 7.09(s, 1H); 7.42(s, 1H); 7.95(s, 1H); 12.00(s, 1H)

MS (ESI): 304 (MH)⁺

(*R,S*)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-3,4-dihydroquinazolin-4-one (2.72g, 8.9mmol) was suspended in thionyl chloride (90ml) and DMF (0.5ml) and heated at reflux for 45 minutes. The thionyl chloride was evaporated under vacuum and the residue azeotroped with toluene three times. The residue was taken up in water and basified (pH8) with saturated aqueous sodium hydrogen carbonate solution. The aqueous layer was extracted

with ethyl acetate (4x400ml). The organic extracts were washed with saturated aqueous sodium hydrogen carbonate solution, water and brine then dried (MgSO₄). After filtration the organic extracts were concentrated under vacuum then dried overnight at 40°C under vacuum to give (*R,S*)-4-chloro-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline (2.62g, 91%) as a solid.

¹H NMR Spectrum: (DMSO-d₆) 1.10(m, 1H); 1.42-1.96(m, 5H); 2.09(m, 1H); 2.15(s, 3H); 2.60(d, 1H); 2.80(d, 1H); 3.98(s, 3H); 4.10(d, 2H); 7.35(s, 1H); 7.42(s, 1H); 8.84(s, 1H)
MS (ESI): 322 (MH)⁺

10 **Example 8**

(*R,S*)-6-Methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline, (prepared as described in Example 7), was chromatographed on Chiral CEL OD (250mm x 4.6mm), (trade mark of Daicel Chemical Industries Ltd), in isohexane/ethanol/triethylamine/TFA (80/20/0.5/0.25). The relevant fractions for *S* (RT 12.55) and *R* (RT 15.88) enantiomers were each combined separately and worked up as follows.

The solution was evaporated under vacuum to give a liquid. This was treated with 5M aqueous sodium hydroxide solution (15ml) and extracted with ethyl acetate. The organic extracts were washed with water then brine and filtered through phase separating paper. The filtrate was evaporated to give (*S*)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline (50mg). The same method was used to give (*R*)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline (71mg).

Example 9

A suspension of 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (0.13g, 0.4mmol), 5-hydroxy-2-methylindole (74mg, 0.5mmol) and potassium carbonate (83mg, 0.6mmol) in DMF (1.5ml) was stirred at 100°C for 2 hours. After cooling to ambient temperature, water (20ml) was added. The precipitate was collected by filtration, washed with water and dried under vacuum at 60°C to give 6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (80mg, 46%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.9-2.0(m, 2H); 2.05-2.2(m, 2H); 2.25-2.4(m, 2H); 2.43(s, 3H); 3.05-3.2(m, 2H); 3.35-3.5(m, 2H); 3.65-3.75(m, 2H); 4.12(s, 3H); 4.35-4.5(t,

2H); 7.0(dd, 1H); 7.35(d, 1H); 7.42(d, 1H); 7.6(s, 1H); 7.85(s, 1H); 9.15(s, 1H)

MS (ESI): 433 (MH)⁺

The starting material was prepared as follows:

5 A mixture of 4-hydroxy-3-methoxybenzoic acid (8.4g, 50mmol), 3-(pyrrolidin-1-yl)propyl chloride (14.75g, 0.1mol), (J. Am. Chem. Soc. 1955, 77, 2272), potassium carbonate (13.8g, 0.1mol) and potassium iodide (1.66g, 10mmol) in DMF (150ml) was stirred and heated at 100°C for 3 hours. The mixture was allowed to cool and the insolubles were removed by filtration and the volatiles were removed from the filtrate by evaporation. The
10 residue was dissolved in ethanol (75ml), 2M aqueous sodium hydroxide (75ml) was added and the mixture was heated at 90°C for 2 hours. The mixture was concentrated by evaporation, acidified with concentrated hydrochloric acid, washed with ether and then subjected to purification on a Diaion (trade mark of Mitsubishi) HP20SS resin column, eluting with water and then with a gradient of methanol (0 to 25%) in dilute hydrochloric acid
15 (pH2.2). The methanol was removed by evaporation and the aqueous residue was freeze dried to give 3-methoxy-4-(3-(pyrrolidin-1-yl)propoxy)benzoic acid hydrochloride (12.2g, 77%).
¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.2(m, 2H); 3.15(t, 2H); 3.3(t, 2H); 3.5(d, 2H); 3.7(t, 2H); 3.82(s, 3H); 4.05(d, 2H); 4.15(t, 2H); 7.07(d, 1H); 7.48(s, 1H); 7.59(d, 1H)
MS - EI: 279 [M]⁺

20 Fuming nitric acid (2.4ml, 57.9mmol) was added slowly at 0°C to a solution of 3-methoxy-4-(3-(pyrrolidin-1-yl)propoxy)benzoic acid hydrochloride (12.15g, 38.17mmol) in TFA (40ml). The cooling bath was removed and the reaction mixture stirred at ambient temperature for 1 hour. The TFA was removed by evaporation and ice/water was added to the residue and the solvent removed by evaporation. The solid residue was dissolved in dilute
25 hydrochloric acid (pH2.2), poured onto a Diaion (trade mark of Mitsubishi) HP20SS resin column and eluted with methanol (gradient 0 to 50%) in water. Concentration of the fractions by evaporation gave a precipitate which was collected by filtration and dried under vacuum over phosphorus pentoxide to give 5-methoxy-2-nitro-4-(3-(pyrrolidin-1-yl)propoxy)benzoic acid hydrochloride (12.1g, 90%).
30 ¹H NMR Spectrum: (DMSO-d₆, TFA) 1.8-1.9 (m, 2H); 2.0-2.1(m, 2H); 2.1-2.2(m, 2H); 3.0-3.1(m, 2H); 3.3(t, 2H); 3.6-3.7(m, 2H); 3.95(s, 3H); 4.25(t, 2H); 7.35(s, 1H); 7.62(s, 1H)

A solution of 5-methoxy-2-nitro-4-(3-(pyrrolidin-1-yl)propoxy)benzoic acid

hydrochloride (9.63g, 24mmol) in thionyl chloride (20ml) and DMF (50 μ l) was heated at 45°C for 1.5 hours. The excess thionyl chloride was removed by evaporation and by azeotroping with toluene (x2). The resulting solid was suspended in THF (250ml) and methylene chloride (100ml) and ammonia was bubbled through the mixture for 30 minutes and the mixture stirred for a further 1.5 hours at ambient temperature. The volatiles were removed by evaporation, the residue was dissolved in water and applied to a Diaion (trade mark of Mitsubishi) HP20SS resin column and eluted with water/methanol (100/0 to 95/5). The solvent was removed by evaporation from the fractions containing product and the residue was dissolved in a minimum of methanol and the solution was diluted with ether. The resulting precipitate was collected by filtration, washed with ether and dried under vacuum to give 5-methoxy-2-nitro-4-(3-(pyrrolidin-1-yl)propoxy)benzamide (7.23g, 73%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.85-1.95(m, 2H); 2-2.1(m, 2H); 2.15-2.25(m, 2H); 3.0-3.1(m, 2H); 3.31(t, 2H); 3.62(t, 2H); 3.93(s, 3H); 4.2(t, 2H); 7.16(s, 1H); 7.60(s, 1H)
MS - EI: 323 [M]⁺

Concentrated hydrochloric acid (5ml) was added to a suspension of 5-methoxy-2-nitro-4-(3-(pyrrolidin-1-yl)propoxy)benzamide (1.5g, 4.64mmol) in methanol (20ml) and the mixture was heated at 50°C to give a solution. Iron powder (1.3g, 23.2mmol) was added in portions and the reaction mixture was then heated at reflux for 1 hour. The mixture was allowed to cool, the insolubles were removed by filtration through diatomaceous earth and the volatiles were removed from the filtrate by evaporation. The residue was purified on a Diaion (trade mark of Mitsubishi) HP20SS resin column, eluting with water and then with dilute hydrochloric acid (pH2). The fractions containing product were concentrated by evaporation and the resulting precipitate was collected by filtration and dried under vacuum over phosphorus pentoxide to give 2-amino-5-methoxy-4-(3-(pyrrolidin-1-yl)propoxy)benzamide hydrochloride (1.44g, 85%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.9(br s, 2H); 2.05(br s, 2H); 2.2(br s, 2H); 3.05(br s, 2H); 3.3(t, 2H); 3.61(br s, 2H); 3.8(s, 3H); 4.11(t, 2H); 7.05(s, 1H); 7.53(s, 1H)
MS - EI: 293 [M]⁺

A mixture of 2-amino-5-methoxy-4-(3-(pyrrolidin-1-yl)propoxy)benzamide hydrochloride (5.92g, 16.2mmol) and Gold's reagent (3.5g, 21.4mmol) in dioxane (50ml) was heated at reflux for 5 hours. Acetic acid (0.7ml) and sodium acetate (1.33g) were added to the reaction mixture which was heated at reflux for a further 5 hours. The mixture was allowed to

cool and the volatiles were removed by evaporation. The residue was dissolved in water, adjusted to pH8 with 2M aqueous sodium hydroxide solution and purified on a Diaion (trademark of Mitsubishi) HP20SS resin column eluting with methanol (gradient 0-50 %) in water. The fractions containing product were concentrated by evaporation and then freeze dried to give 4-hydroxy-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (4.55g, 83%).
¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.9(m, 2H); 2.0-2.1(m, 2H); 2.2-2.3(m, 2H); 3.05(m, 2H); 3.34(t, 2H); 3.6-3.7(br s, 2H); 3.94(s, 3H); 4.27(t, 2H); 7.31(s, 1H); 7.55(s, 1H); 9.02(s, 1H)

A mixture of 4-hydroxy-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (1.7g, 5mmol) and thionyl chloride (25ml) containing DMF (0.2ml) was heated at reflux for 3 hours. Excess thionyl chloride was removed by evaporation and by azeotroping with toluene (x2). The residue was suspended in ether and 10% aqueous solution of sodium hydrogen carbonate was added to the mixture. The organic layer was separated, dried (MgSO₄) and the solvent removed by evaporation to give 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-

yl)propoxy)quinazoline (1.94g, quantitative).
¹H NMR Spectrum: (CDCl₃) 1.8(br s, 4H); 2.17(m, 2H); 2.6(br s, 4H); 2.7(t, 2H); 4.05(s, 3H); 4.3(t, 2H); 7.35(s, 1H); 7.38(s, 1H); 8.86(s, 1H)
MS - ESI: 322 [MH]⁺

Example 10

A suspension of 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (74mg, 0.23mmol), potassium carbonate (48mg, 0.35mmol) and 7-hydroxyquinoline (40.6mg, 0.28mmol) in DMF (1.5ml) was heated at 100°C for 3 hours. After cooling, the mixture was stirred for 10 hours at ambient temperature and then overnight at 5°C. After dilution with methylene chloride (5ml), the mixture was poured onto a column of silica and was eluted with an increasing gradient of methanol/methylene chloride (10/90, 20/80) followed by ammonia/methanol (5%) in methylene chloride (25/75) to give, after removal of the volatiles by evaporation and drying under vacuum, 6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline (82mg, 88%).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5(m, 2H); 1.75-1.9(m, 3H); 1.9-2.05(m, 2H); 2.12(s, 3H); 2.8-2.9(d, 2H); 4.5(s, 3H); 4.1(d, 2H); 7.4(s, 1H); 7.6(dd, 1H); 7.62(dd, 1H)
MS (ESI): 431 [MH]⁺

The starting material was prepared as follows:

To a solution of ethyl 4-piperidinecarboxylate (30g, 0.19mol) in ethyl acetate (150ml) cooled at 5°C was added dropwise a solution of di-tert-butyl dicarbonate (41.7g, 0.19mol) in ethyl acetate (75ml) while maintaining the temperature in the range 0-5°C. After stirring for 48 hours at ambient temperature, the mixture was poured onto water (300ml). The organic layer was separated, washed successively with water (200ml), 0.1M aqueous hydrochloric acid (200ml), saturated sodium hydrogen carbonate (200ml) and brine (200ml), dried (MgSO_4) and evaporated to give ethyl 4-(1-tert-butyloxycarbonylpiperidine)carboxylate (48g, 98%).

^1H NMR Spectrum: (CDCl_3) 1.25(t, 3H); 1.45(s, 9H); 1.55-1.70(m, 2H); 1.8-2.0(d, 2H); 2.35-2.5(m, 1H); 2.7-2.95(t, 2H); 3.9-4.1(br s, 2H); 4.15 (q, 2H)

To a solution of ethyl 4-(1-tert-butyloxycarbonylpiperidine)carboxylate (48g, 0.19mol) in dry THF (180ml) cooled at 0°C was added dropwise a solution of 1M lithium aluminium hydride in THF (133ml, 0.133mol). After stirring at 0°C for 2 hours, water (30ml) was added followed by 2M sodium hydroxide (10ml). The precipitate was filtered through diatomaceous earth and washed with ethyl acetate. The filtrate was washed with water, brine, dried (MgSO_4) and evaporated to give 4-hydroxymethyl-1-tert-butyloxycarbonylpiperidine (36.3g, 89%).

^1H NMR Spectrum: (CDCl_3) 1.05-1.2(m, 2H); 1.35-1.55(m, 10H); 1.6-1.8(m, 2H); 2.6-2.8(t, 2H); 3.4-3.6(t, 2H); 4.0-4.2(br s, 2H)

MS (EI): 215 $[\text{M}]^+$

To a solution of 4-hydroxymethyl-1-tert-butyloxycarbonylpiperidine (52.5g, 0.244mol) in tert-butyl methyl ether (525ml) was added 1,4-diazabicyclo[2.2.2]octane (42.4g, 0.378mol). After stirring for 15 minutes at ambient temperature, the mixture was cooled to 5°C and a solution of toluene sulphonyl chloride (62.8g, 0.33mmol) in tert-butyl methyl ether (525ml) was added dropwise over 2 hours while maintaining the temperature at 0°C. After stirring for 1 hour at ambient temperature, petroleum ether (1l) was added. The precipitate was removed by filtration. The filtrate was evaporated to give a solid. The solid was dissolved in ether and washed successively with 0.5M aqueous hydrochloric acid (2x500ml), water, saturated sodium hydrogen carbonate and brine, dried (MgSO_4) and evaporated to give 4-(4-methylphenylsulphonyloxymethyl)-1-tert-butyloxycarbonylpiperidine (76.7g, 85%).

¹H NMR Spectrum: (CDCl₃) 1.0-1.2(m, 2H); 1.45(s, 9H); 1.65(d, 2H); 1.75-1.9(m, 2H); 2.45(s, 3H); 2.55-2.75(m, 2H); 3.85(d, 1H); 4.0-4.2(br s, 2H); 7.35(d, 2H); 7.8(d, 2H)

MS (ESI): 392 [MNa]⁺

To a suspension of ethyl 3-methoxy-4-hydroxybenzoate (19.6g, 0.1mol) and potassium carbonate (28g, 0.2mol) in dry DMF (200ml) was added 4-(4-methylphenylsulphonyloxymethyl)-1-tert-butyloxycarbonylpiperidine (40g, 0.11mol). After stirring at 95°C for 2.5 hours, the mixture was cooled to ambient temperature and partitioned between water and ethyl acetate/ether. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The resulting oil was crystallised from petroleum ether and the suspension was stored overnight (at 5°C). The solid was collected by filtration, washed with petroleum ether and dried under vacuum to give ethyl 3-methoxy-4-(1-tert-butyloxycarbonylpiperidin-4-ylmethoxy)benzoate (35g, 89%).

m.p. 81-83°C

¹H NMR Spectrum: (CDCl₃) 1.2-1.35(m, 2H); 1.4(t, 3H); 1.48(s, 9H); 1.8-1.9(d, 2H); 2.0-2.15(m, 2H); 2.75(t, 2H); 3.9(d, 2H); 3.95(s, 3H); 4.05-4.25(br s, 2H); 4.35(q, 2H); 6.85(d, 1H); 7.55(s, 1H); 7.65(d, 1H)

MS (ESI): 416 [MNa]⁺

Elemental analysis:	Found	C 63.4	H 8.0	N 3.5
C ₂₁ H ₃₁ NO ₆ 0.3H ₂ O	Requires	C 63.2	H 8.0	N 3.5%

To a solution of ethyl 3-methoxy-4-(1-tert-butyloxycarbonylpiperidin-4-ylmethoxy)benzoate (35g, 89mmol) in formic acid (35ml) was added formaldehyde (12M, 37% in water, 35ml, 420mmol). After stirring at 95°C for 3 hours, the volatiles were removed by evaporation. The residue was dissolved in methylene chloride and 3M hydrogen chloride in ether (40ml, 120mmol) was added. After dilution with ether, the mixture was triturated until a solid was formed. The solid was collected by filtration, washed with ether and dried under vacuum overnight at 50°C to give ethyl 3-methoxy-4-(1-methylpiperidin-4-ylmethoxy)benzoate (30.6g, quant.).

¹H NMR Spectrum: (DMSO-d₆) 1.29(t, 3H); 1.5-1.7(m, 2H); 1.95(d, 2H); 2.0-2.15(br s, 1H); 2.72(s, 3H); 2.9-3.1(m, 2H); 3.35-3.5(br s, 2H); 3.85(s, 3H); 3.9-4.05(br s, 2H); 4.3(q, 2H); 7.1(d, 1H); 7.48(s, 1H); 7.6(d, 1H)

MS (ESI): 308 [MH]⁺

A solution of ethyl 3-methoxy-4-(1-methylpiperidin-4-ylmethoxy)benzoate (30.6g,

89mmol) in methylene chloride (75ml) was cooled to 0-5°C. TFA (37.5ml) was added followed by the dropwise addition over 15 minutes of a solution of fuming 24M nitric acid (7.42ml, 178mmol) in methylene chloride (15ml). After completion of the addition, the solution was allowed to warm up and stirred at ambient temperature for 2 hours. The volatiles were removed under vacuum and the residue was dissolved in methylene chloride (50ml). The solution was cooled to 0-5°C and ether was added. The precipitate was collected by filtration, and dried under vacuum at 50°C. The solid was dissolved in methylene chloride (500ml) and 3M hydrogen chloride in ether (30ml) was added followed by ether (500ml). The solid was collected by filtration and dried under vacuum at 50°C to give ethyl 3-methoxy-4-(1-methylpiperidin-4-ylmethoxy)-6-nitrobenzoate (28.4g, 82%).

¹H NMR Spectrum: (DMSO-d₆) 1.3(t, 3H); 1.45-1.65(m, 2H); 1.75-2.1(m, 3H); 2.75(s, 3H); 2.9-3.05(m, 2H); 3.4-3.5(d, 2H); 3.95(s, 3H); 4.05(d, 2H); 4.3(q, 2H); 7.32(s, 1H); 7.66(s, 1H)
MS (ESI): 353 [MH]⁺

A suspension of ethyl 3-methoxy-4-(1-methylpiperidin-4-ylmethoxy)-6-nitrobenzoate (3.89g, 10mmol) in methanol (80ml) containing 10% platinum on activated carbon (50% wet) (389mg) was hydrogenated at 1.8 atmospheres pressure until uptake of hydrogen ceased. The mixture was filtered and the filtrate was evaporated. The residue was dissolved in water (30ml) and adjusted to pH10 with a saturated solution of sodium hydrogen carbonate. The mixture was diluted with ethyl acetate/ether (1/1) and the organic layer was separated. The aqueous layer was further extracted with ethyl acetate/ether and the organic layers were combined. The organic layers were washed with water, brine, dried (MgSO₄), filtered and evaporated. The resulting solid was triturated in a mixture of ether/petroleum ether, filtered, washed with petroleum ether and dried under vacuum at 60°C to give ethyl 6-amino-3-methoxy-4-(1-methylpiperidin-4-ylmethoxy)benzoate (2.58g, 80%).

m.p. 111-112°C

¹H NMR Spectrum: (CDCl₃) 1.35(t, 3H); 1.4-1.5(m, 2H); 1.85(m, 3H); 1.95(t, 2H); 2.29(s, 3H); 2.9(d, 2H); 3.8(s, 3H); 3.85(d, 2H); 4.3(q, 2H); 5.55(br s, 2H); 6.13(s, 1H); 7.33(s, 1H)
MS (ESI): 323 [MH]⁺

Elemental analysis: Found C 62.8 H 8.5 N 8.3

C₁₇H₂₆N₂O₄ 0.2H₂O Requires C 62.6 H 8.2 N 8.6%

A solution of ethyl 6-amino-3-methoxy-4-(1-methylpiperidin-4-ylmethoxy)benzoate (16.1g, 50mmol) in 2-methoxyethanol (160ml) containing formamidine acetate (5.2g,

50mmol) was heated at 115°C for 2 hours. Formamidinium acetate (10.4g, 100mmol) was added in portions every 30 minutes during 4 hours. Heating was prolonged for 30 minutes after the last addition. After cooling, the volatiles were removed under vacuum. The solid was dissolved in ethanol (100ml) and methylene chloride (50ml). The precipitate was removed by filtration and the filtrate was concentrated to a final volume of 100ml. The suspension was cooled to 5°C and the solid was collected by filtration, washed with cold ethanol followed by ether and dried under vacuum overnight at 60°C to give 6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-3,4-dihydroquinazolin-4-one (12.7g, 70%).

¹H NMR Spectrum: (DMSO-d₆) 1.25-1.4(m, 2H); 1.75(d, 2H); 1.9(t, 1H); 1.9(s, 3H); 2.16(s, 2H); 2.8(d, 2H); 3.9(s, 3H); 4.0(d, 2H); 7.11(s, 1H); 7.44(s, 1H); 7.97(s, 1H)

MS (ESI): 304 [MH]⁺

A solution of 6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-3,4-dihydroquinazolin-4-one (2.8g, 9.24mmol) in thionyl chloride (28ml) containing DMF (280μl) was refluxed at 85°C for 1 hour. After cooling, the volatiles were removed by evaporation. The precipitate was triturated with ether, filtered, washed with ether and dried under vacuum. The solid was dissolved in methylene chloride and saturated aqueous sodium hydrogen carbonate was added. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated to give 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (2.9g, 98%).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5(m, 2H); 1.75-1.9(m, 3H); 2.0(t, 1H); 2.25(s, 3H); 2.85(d, 2H); 4.02(s, 3H); 4.12(d, 2H); 7.41(s, 1H); 7.46(s, 1H); 8.9(s, 1H)

MS (ESI): 322 [MH]⁺

Example 11

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (0.13g, 0.4mmol), (prepared as described for the starting material in Example 10), was reacted with 5-hydroxy-2-methylindole (74mg, 0.5mol) to give 6-methoxy-4-(2-methylindol-5-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (137mg, 79%).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.45(m, 2H); 1.7-1.95(m, 5H); 2.15(s, 3H); 2.4(s, 3H); 2.8(d, 2H); 3.98(s, 3H); 4.05(d, 2H); 6.14(s, 1H); 6.88(d, 1H); 7.29(s, 1H); 7.32(d, 1H); 7.35(s, 1H); 7.6(s, 1H); 8.45(s, 1H)

MS (ESI): 433 [MH]⁺

Example 12

To a solution of 4-chloro-6-methoxy-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)quinazoline (115mg, 0.28mmol) and 7-hydroxyquinoline (50mg, 0.33mmol) in DMF (1.5ml) was added potassium carbonate (60mg, 0.42mmol). The mixture was stirred for 2 hours at 100°C. After cooling, and removal of the volatiles by evaporation, the residue was partitioned between ethyl acetate and water. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with ethylacetate/methylene chloride/methanol (1/1/0 followed by 40/50/10 and 0/9/1). After removal of the volatiles by evaporation, the residue was triturated with pentane, filtered and dried under vacuum to give 6-methoxy-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline (110mg, 76%).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.45(m, 2H); 1.75-1.9(m, 3H); 2.05(t, 2H); 2.72(t, 2H); 2.95(d, 2H); 3.05(s, 3H); 3.35-3.45(m, 2H); 4.00(s, 3H); 4.1(d, 2H); 7.41(s, 1H); 7.57(dd, 1H); 7.62(dd, 1H); 7.65(s, 1H); 7.93(s, 1H); 8.12(d, 1H); 8.45(d, 1H); 8.55(s, 1H); 8.95(d, 1H)

MS (ESI): 523 [MH]⁺

Elemental analysis:	Found	C 61.3	H 6.0	N 10.6
---------------------	-------	--------	-------	--------

C ₂₇ H ₃₀ N ₄ O ₅ S 0.4H ₂ O	Requires	C 61.2	H 5.9	N 10.6%
---	----------	--------	-------	---------

The starting material was prepared as follows:

Sodium hydride (1.44g of a 60% suspension in mineral oil, 36mmol) was added in portions over 20 minutes to a solution of 7-benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (8.46g, 30mmol), (prepared as described for the starting material in Example 1), in DMF (70ml) and the mixture was stirred for 1.5 hours. Chloromethyl pivalate (5.65g, 37.5mmol) was added dropwise and the mixture stirred for 2 hours at ambient temperature. The mixture was diluted with ethyl acetate (100ml) and poured onto ice/water (400ml) and 2M hydrochloric acid (4ml). The organic layer was separated and the aqueous layer extracted with ethyl acetate, the combined extracts were washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was triturated with a mixture of ether and petroleum ether, the solid was collected by filtration and dried under vacuum to give 7-

benzyloxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (10g, 84%).

¹H NMR Spectrum: (DMSO-d₆) 1.11(s, 9H); 3.89(s, 3H); 5.3(s, 2H); 5.9(s, 2H); 7.27(s, 1H); 7.35(m, 1H); 7.47(t, 2H); 7.49(d, 2H); 7.51(s, 1H); 8.34(s, 1H)

A mixture of 7-benzyloxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-

- 5 dihydroquinazolin-4-one (7g, 17.7mmol) and 10% palladium-on-charcoal catalyst (700mg) in ethyl acetate (250ml), DMF (50ml), methanol (50ml) and acetic acid (0.7ml) was stirred under hydrogen at atmospheric pressure for 40 minutes. The catalyst was removed by filtration and the solvent removed from the filtrate by evaporation. The residue was triturated with ether, collected by filtration and dried under vacuum to give 7-hydroxy-6-methoxy-3-
10 ((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (4.36g, 80%).

¹H NMR Spectrum: (DMSO-d₆) 1.1(s, 9H); 3.89(s, 3H); 5.89(s, 2H); 7.0(s, 1H); 7.48(s, 1H); 8.5(s, 1H)

A suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-

- 15 dihydroquinazolin-4-one (6.12g, 20mmol) potassium carbonate (5.52g, 40mmol) in DMF (60ml) was stirred at ambient temperature for 30 minutes. 4-(4-Methylphenylsulphonyloxymethyl)-1-tert-butyloxycarbonylpiperidine (8.86g, 24mmol), (prepared as described for the starting material in Example 10), was added and the mixture was stirred at 100°C for 2 hours. After cooling, the mixture was poured onto water/ice (400ml, 1/1) containing 2M hydrochloric acid (10ml). The precipitate was collected by
20 filtration, washed with water and dried under vacuum over phosphorus pentoxide. The solid was triturated in a mixture of ether/pentane (1/1), collected by filtration and dried to give 6-methoxy-3-((pivaloyloxy)methyl)-7-((1-tert-butyloxycarbonylpiperidin-4-yl)methoxy)-3,4-dihydroquinazolin-4-one (7.9g, 78.5%).

- ¹H NMR Spectrum: (DMSO-d₆) 1.1(s, 9H); 1.1-1.3(m, 2H); 1.42(s, 9H); 1.73(d, 2H); 1.93-
25 2.1(br s, 1H); 2.65-2.9(br s, 2H); 3.9(s, 3H); 3.9-4.1(m, 4H); 5.9(s, 2H); 7.2(s, 1H); 7.5(s, 1H); 8.35(s, 1H)

MS (ESI): 526 [MNa]⁺

A solution of 6-methoxy-3-((pivaloyloxy)methyl)-7-((1-tert-

- 30 butyloxycarbonylpiperidin-4-yl)methoxy)-3,4-dihydroquinazolin-4-one (7.9g, 16mmol) in methylene chloride (80ml) containing 5.5M hydrogen chloride in isopropanol (80ml) was stirred for 1 hour at ambient temperature. Ether was added and the solid was collected by

filtration, washed with ether and dried under vacuum at 60°C to give 6-methoxy-7-((piperidin-4-yl)methoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one hydrochloride (6.9g, 100%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.15(s, 9H); 1.5-1.7(m, 2H); 2.0(d, 2H); 2.2-2.3(br s, 1H); 3.0(t, 2H); 3.4(d, 2H); 3.94(s, 3H); 4.15(d, 2H); 5.97(s, 2H); 7.3(s, 1H); 7.6(s, 1H); 8.65(s, 1H)

MS (ESI): 404 [MH]⁺

To a solution of 6-methoxy-7-((piperidin-4-yl)methoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one hydrochloride (0.88g, 2mmol) and triethylamine (0.3ml, 2.1mmol) in methanol (10ml) and methylene chloride (10ml) was added potassium carbonate (280mg, 2mmol) and methyl vinyl sulfone (0.4ml, 2.1mmol). After stirring for 2 hours at ambient temperature, the volatiles were removed under vacuum. The residue was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried (MgSO₄) and evaporated to give 6-methoxy-7-((1-(2-methylsulphonylethyl)piperidin-4-yl)methoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (0.55g, 54%).

¹H NMR Spectrum: (DMSO-d₆) 1.09(s, 9H); 1.25-1.4(m, 2H); 1.7-1.9(m, 3H); 2.0(t, 2H); 2.7(t, 2H); 2.95(d, 2H); 3.02(s, 3H); 3.25-3.45(m, 2H); 3.9(s, 3H); 4.0(d, 2H); 5.9(s, 2H); 7.15(s, 1H); 7.49(s, 1H); 8.35(s, 1H)

MS (ESI): 510 [MH]⁺

To a suspension of 6-methoxy-7-((1-(2-methylsulphonylethyl)piperidin-4-yl)methoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (90mg, 0.18mmol) in methanol (3ml) was added 2M aqueous sodium hydroxide (180μl, 0.35mmol). After stirring for 2 hours at ambient temperature, the mixture was adjusted to pH10 with 2M hydrochloric acid. The volatiles were removed under vacuum and the residue was suspended in water, filtered, washed with water followed by ether and dried under vacuum at 60°C to give 6-methoxy-7-((1-(2-methylsulphonylethyl)piperidin-4-yl)methoxy)-3,4-dihydroquinazolin-4-one (55mg, 79%).

¹H NMR Spectrum: (DMSO-d₆) 1.2-1.4(m, 2H); 1.7-1.85(m, 3H); 2.0(t, 2H); 2.7(t, 2H); 2.9(d, 2H); 3.02(s, 3H); 3.3-3.5(m, 2H); 3.9(s, 3H); 4.0(d, 2H); 7.11(s, 1H); 7.45(s, 1H); 7.97(s, 1H)

MS (ESI): 396 [MH]⁺

A solution of 6-methoxy-7-((1-(2-methylsulphonylethyl)piperidin-4-yl)methoxy)-3,4-dihydroquinazolin-4-one (335mg, 0.85mmol) in thionyl chloride (5ml) containing DMF

(50µl) was refluxed for 1 hour. After cooling, the volatiles were removed under vacuum and the residue was triturated with ether and filtered. The solid was suspended in methylene chloride and sodium hydrogen carbonate was added. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was triturated with ether, filtered and dried under vacuum to give 4-chloro-6-methoxy-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-ylmethoxy)quinazoline (335mg, 95%).

¹H NMR Spectrum: (DMSO-d₆) 1.25-1.45(m, 2H); 1.75-1.90(m, 3H); 2.0(t, 2H); 2.7(t, 2H); 2.92(d, 2H); 3.03(s, 3H); 3.2-3.35(m, 2H); 4.0(s, 3H); 4.1(d, 2H); 7.40(s, 1H); 7.45(s, 1H); 8.9(s, 1H)

MS (ESI): 414 [MH]⁺

Example 13

Using a procedure analogous to that described for Example 10, 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (130mg, 0.4mmol), (prepared as described for the starting material in Example 10), was reacted with 4-methyl-7-hydroxyquinoline (80mg, 0.5mol), (Chem. Ber. 1967, 100, 2077), to give 6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(4-methylquinolin-7-yloxy)quinazoline (160mg, 90%).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5(m, 2H); 1.7-1.95(m, 3H); 1.9(t, 2H); 2.17(s, 3H); 2.74(s, 3H); 2.8(d, 2H); 4.07(s, 3H); 4.1(d, 2H); 7.4(m, 2H); 7.65(dd, 1H); 7.65(s, 1H); 7.9(s, 1H); 8.21(d, 1H); 8.54(s, 1H); 8.78(d, 1H)

MS (ESI): 445 [MH]⁺

Example 14

A solution of 4-chloro-6-methoxy-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)quinazoline (115mg, 0.28mmol), (prepared as described for the starting material in Example 12), 5-hydroxy-2-methylindole (50mg, 0.33mmol) and potassium carbonate (60mg, 0.42mmol) in DMF (1.5ml) was stirred at 100°C for 2 hours. After cooling, the mixture was partitioned between ethyl acetate and water. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by chromatography eluting with ethyl acetate/methylene chloride (1/1) followed by methanol/ethyl acetate/methylene chloride (1/4/5 and 1/0/9) to give 6-methoxy-4-(2-methylindol-5-yloxy)-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)quinazoline (60mg, 41%).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.45(m, 2H); 1.75-1.92(m, 3H); 2.02(t, 2H); 2.4(s, 3H); 2.7(t, 2H); 2.95(d, 2H); 3.05(s, 3H); 4.0(s, 3H); 4.05(d, 2H); 6.15(s, 1H); 6.85(dd, 1H); 7.25(s, 1H); 7.3(d, 1H); 7.38(s, 1H); 7.6(s, 1H); 8.45(s, 1H)

MS (ESI): 525 [MH]⁺

5	Elemental analysis:	Found	C 60.7	H 6.2	N 10.5
	C ₂₇ H ₃₂ O ₅ S 0.5H ₂ O	Requires	C 60.8	H 6.2	N 10.5%

Example 15

10 Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (0.13g, 0.4mol), (prepared as described for the starting material in Example 9), was reacted with 7-hydroxy-4-methylquinoline (80mg, 0.5mol), (Chem. Berich. 1967, 100, 2077), to give **6-methoxy-4-(4-methylquinolin-7-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (155mg, 87%).

15 ¹H NMR Spectrum: (DMSO-d₆) 1.7(br s, 4H); 2.05(m, 2H); 2.5(br s, 4H); 2.6(t, 2H); 2.75(s, 3H); 4.02(s, 3H); 4.3(t, 2H); 7.41(s, 1H); 7.45(d, 1H); 7.65(s, 1H); 7.65(d, 1H); 7.95(s, 1H); 8.25(d, 1H); 8.55(s, 1H); 8.8(d, 1H)

MS (ESI): 445 [MH]⁺

Example 16

20 Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (0.13g, 0.4mol), (prepared as described for the starting material in Example 9), was reacted with 2,2,4-trimethyl-1,2-dihydroquinolin-6-ol (95mg, 0.5mmol), (IZV. ACAD. NAVK. SSSR. Ser. Khim. 1981, 9, 2008), to give **6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)-4-(2,2,4-trimethyl-1,2-dihydroquinolin-6-yloxy)quinazoline** (90mg, 47%).

25 ¹H NMR Spectrum: (DMSO-d₆) 1.23(s, 6H); 1.7(br s, 4H); 1.85(s, 3H); 2.0(m, 2H); 2.45(br s, 4H); 2.57(t, 2H); 3.95(s, 3H); 4.25(t, 2H); 5.35(s, 1H); 5.9(s, 1H); 6.5(d, 1H); 6.8(dd, 1H); 6.85(s, 1H); 7.32(s, 1H); 7.52(s, 1H); 8.5(s, 1H)

MS (ESI): 475 [MH]⁺

30

Example 17

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-

((1-methylpiperidin-4-yl)methoxy)quinazoline (0.13g, 0.4mmol), (prepared as described for the starting material in Example 10), was reacted with 2,2,4-trimethyl-1,2-dihydroquinolin-6-ol (95mg, 0.5mmol), (IZV. ACAD. NAVK. SSSR. Ser. Khim. 1981, 9, 2008), to give **6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(2,2,4-trimethyl-1,2-dihydroquinolin-6-yloxy)quinazoline** (140mg, 74%).

¹H NMR Spectrum: (DMSO-d₆) 1.15(s, 6H); 1.3-1.45(m, 2H); 1.7-2.0(m, 8H); 2.16(s, 3H); 2.65-2.85(d, 2H); 4.0(s, 3H); 4.05(d, 2H); 5.35(s, 1H); 5.9(s, 1H); 6.5(d, 1H); 6.80(d, 1H); 6.82(s, 1H); 7.33(s, 1H); 7.5(s, 1H); 8.52(s, 1H)

MS (ESI): 475 [MH]⁺

Example 18

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (0.13g, 0.4mol), (prepared as described for the starting material in Example 10), was reacted with 2,4-dimethyl-7-hydroxyquinoline (87mg, 0.5mmol), (Chem. Berichte, 1903, 36, 4016), to give **4-(2,4-dimethylquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline** (61mg, 33%).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5(m, 2H); 1.7-1.95(m, 5H); 2.2(s, 3H); 2.65(s, 3H); 2.7(s, 3H); 2.75-2.9(br d, 2H); 4.05(s, 3H); 4.1(d, 2H); 7.3(s, 1H); 7.4(s, 1H); 7.52(d, 1H); 7.65(s, 1H); 7.8(s, 1H); 8.15(d, 1H); 8.55(s, 1H)

MS (ESI): 459 [MH]⁺

Example 19

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (0.13g, 0.4mol), (prepared as described for the starting material in Example 10), was reacted with 6-hydroxy-2*H*-4*H*-1,4-benzoxazin-3-one (83mg, 0.5mol), (J. Chem. Soc. C, 1971, 2696), to give **6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(3-oxo-2*H*-4*H*-1,4-benzoxazin-6-yloxy)quinazoline** (158mg, 88%).

¹H NMR Spectrum: (DMSO-d₆) 1.25-1.45(m, 2H); 1.8(d, 2H); 1.7-1.9(m, 1H); 1.9(t, 2H); 2.2(s, 3H); 2.8(d, 2H); 3.97(s, 3H); 4.05(d, 2H); 4.65(s, 2H); 6.8(s, 1H); 6.85(d, 1H); 7.05(d, 1H); 7.35(s, 1H); 7.52(s, 1H); 8.55(s, 1H)

MS (ESI): 451 [MH]⁺

Example 20

Using a procedure analogous to that described for Example 9, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (0.13g, 0.4mol), (prepared as described for the starting material in Example 9), was reacted with 6-hydroxy-2*H*-4*H*-1,4-benzoxazin-3-one (83mg, 0.5mol), (J. Chem. Soc. C, 1971, 2696), to give **6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)-4-(3-oxo-2*H*-4*H*-1,4-benzoxazin-6-yloxy)quinazoline** (170mg, 94%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 1.8-2.0(m, 2H); 2.0-2.15(m, 2H); 2.2-2.35(m, 2H); 3.0-3.2(m, 2H); 3.4(t, 2H); 3.6-3.75(m, 2H); 4.05(s, 3H); 4.35(t, 2H); 4.65(s, 2H); 6.85(s, 1H); 6.9(d, 1H); 7.1(d, 1H); 7.5(s, 1H); 7.7(s, 1H); 8.9(s, 1H)

MS (ESI): 451 [MH]⁺

Example 21

Using a procedure analogous to that described for Example 10, 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (74mg, 0.23mmol), (prepared as described for the starting material in Example 10), was reacted with 6-hydroxyquinoline (41mg, 0.28mol) to give **6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(quinolin-6-yloxy)quinazoline** (89mg, 94%).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5(m, 2H); 1.8(d, 2H); 1.9(t, 2H); 1.8-1.9(m, 1H); 2.2(s, 3H); 2.82(d, 2H); 4.02(s, 3H); 4.1(d, 2H); 7.4(s, 1H); 7.6(dd, 1H); 7.65(s, 1H); 7.75(d, 1H); 7.95(s, 1H); 8.15(d, 1H); 8.4(d, 1H); 8.55(s, 1H); 8.95(d, 1H)

MS (ESI): 431 [MH]⁺

Example 22

To 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (250mg, 0.74mmol), (prepared as described for the starting material in Example 1), in suspension in DMF (4ml) were successively added 4-chloro-7-hydroxyquinoline (133mg, 0.74mmol) and potassium carbonate (153mg, 1mmol) and the reaction mixture heated to 100°C. More 4-chloro-7-hydroxyquinoline (27mg, 0.15mmol) was added after one hour and heating was continued for a further 30 minutes. The product precipitated upon cooling to ambient temperature. The reaction mixture was diluted with water, the product was collected by filtration and washed with more water. The dried solid was triturated with ether and filtered to give **4-(4-chloroquinolin-7-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline** (166mg, 47%).

¹H NMR Spectrum: (DMSO-d₆, CF₃CO₂D) 2.3(m, 2H); 3.2(m, 2H); 3.4(m, 2H); 3.5(m, 2H); 3.7(m, 2H); 4.0(m, 2H); 4.1(s, 3H); 4.4(m, 2H); 7.55(s, 1H); 7.75(s, 1H); 7.90(dd, 1H); 7.95(d, 1H); 8.15(d, 1H); 8.45 (d, 1H); 8.80(s, 1H); 9.05(d, 1H)

MS - ESI: 481 [MH]⁺

5	Elemental analysis:	Found	C 61.8	H 5.1	N 11.5
	C ₂₅ H ₂₅ ClN ₄ O ₄	Requires	C 62.4	H 5.2	N 11.7%

The starting material was prepared as follows:

A solution of 7-benzyloxy-4-chloroquinoline (17g, 56mmol), (Konishi et al. WO 96/11187), in TFA (170ml) was heated at reflux for 2 hours. The solvent was removed under vacuum and the residue was triturated with ether, filtered and washed with ether. The solid was suspended in an aqueous solution of sodium hydrogen carbonate (5.5g, 65mmol in 200ml of water) and stirred at ambient temperature for 30 minutes. The solid was collected by filtration, washed with water and dried overnight under vacuum and over phosphorus pentoxide to give 4-chloro-7-hydroxyquinoline (9.85g, 98%).

¹H NMR Spectrum: (DMSO-d₆) 7.37(s, 1H); 7.39(d, 1H); 7.62(d, 1H); 8.15(d, 1H); 8.8(d, 1H)
MS - EI: m/z 179 [M.]⁺

Example 23

A solution of 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (74mg, 0.23mmol), (prepared as described for the starting material in Example 10), and 2-hydroxynaphthalene (40mg, 0.28mmol) in DMF (1.5ml) containing potassium carbonate (48mg, 0.35mmol) was stirred at 100°C for 3.5 hours. After cooling, methylene chloride (4.5ml) was added and the mixture was poured onto a column of silica (SiO₂ Isolute®) and eluted with, successively, methylene chloride, methylene chloride/methanol (9/1), methylene chloride/methanol/3M ammonia in methanol (75/20/5). The fractions containing the product were evaporated under vacuum. The residues was triturated with ether, filtered and dried under vacuum to give 6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(2-naphthyloxy)quinazoline (80mg, 83%).

MS - ESI: 430 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.35-1.45 (m, 2H), 1.8 (d, 2H), 2.0 (t, 1H), 2.2 (s, 3H), 2.85

(d, 2H), 3.3-3.4 (m, 2H), 4.02 (s, 3H), 4.1 (d, 2H), 7.4 (s, 1H), 7.5 (dd, 1H), 7.55 (m, 2H), 7.65 (s, 1H), 7.88 (s, 1H), 7.98 (d, 1H), 8.0 (d, 1H), 8.1 (d, 1H), 8.55 (s, 1H)

Example 24

5 A solution of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (74mg, 0.23mmol), (prepared as described for the starting material in Example 1), and 3,4-(methylenedioxy)aniline (53mg, 0.24mmol) in a solution of isopropanol (3.5ml) containing 5.5M hydrogen chloride in isopropanol (42μl) was heated for 3 hours. After cooling to ambient temperature, the reaction mixture was cooled to 0°C and maintained at this
10 temperature overnight. The precipitate was collected by filtration, washed with ethyl acetate and dried under vacuum to give 4-(1,3-benzodioxol-5-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (82mg, 76%).


MS - ESI: 439 [MH]⁺

15 ¹H NMR Spectrum: (DMSO-d₆) 2.3-2.4 (m, 2H), 3.05-3.2 (m, 2H), 3.25-3.35 (m, 2H), 3.5 (d, 2H), 3.82 (t, 2H), 4.0 (d, 2H), 4.05 (s, 3H), 4.32 (t, 2H), 6.1 (s, 2H), 7.02 (d, 1H), 7.1 (dd, 1H), 7.3 (s, 1H), 7.4 (s, 1H), 8.32 (s, 1H), 8.8 (s, 1H)

Examples 25-29

20 Using an analogous procedure to that described in Example 24, 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline, (prepared as described for the starting material in Example 1), was used in the synthesis of the compounds described in Table I hereinafter as detailed in the notes a)-e) to Table I.

Table I



25

Example No.	Weight (mg)	yield %	MS-ESI [MH] ⁺	note	R
25	104	90	435.1	a	1- <i>H</i> -indazol-6-yl

26	102	89	435.1	b	1- <i>H</i> -indazol-5-yl
27	99	84	452	c	1,3-benzothiazol-6-yl
28	108	91	466	d	2-methyl-1,3-benzothiazol-5-yl
29	102	95	435.1	e	2,3-dihydro-1 <i>H</i> -inden-5-yl

Notes

- a) 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (74mg) was reacted with 6-aminoindazole (32mg) to give **4-(1-*H*-indazol-6-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline**.
- ¹H NMR Spectrum: (DMSO-d₆) 2.3-2.4 (m, 2H), 3.05-3.2 (m, 2H), 3.2-3.3 (m, 2H), 3.52 (d, 2H), 3.85 (t, 2H), 4.0 (d, 2H), 4.05 (s, 3H), 4.32 (t, 2H), 7.42 (s, 1H), 7.45 (d, 1H), 7.85 (d, 1H), 7.98 (s, 1H), 8.1 (s, 1H), 8.42 (s, 1H), 8.85 (s, 1H)
- 5 b) 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (74mg) was reacted with 5-aminoindazole (32mg) to give **4-(1-*H*-indazol-5-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline**.
- ¹H NMR Spectrum: (DMSO-d₆) 2.3-2.4 (m, 2H), 3.05-3.2 (m, 2H), 3.25-3.3 (m, 2H), 3.45-3.55 (m, 2H), 3.8-3.9 (m, 2H), 3.9-4.02 (m, 2H), 4.05 (s, 3H), 4.32 (t, 2H), 7.42 (s, 1H), 7.65 (m, 2H), 8.05 (s, 1H), 8.15 (s, 1H), 8.4 (s, 1H), 8.75 (s, 1H)
- 15 c) 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (74mg) was reacted with 6-aminothiazole (36mg) to give **4-(1,3-benzothiazol-6-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline**.
- 20 ¹H NMR Spectrum: (DMSO-d₆) 2.3-2.4 (m, 2H), 3.05-3.2 (m, 2H), 3.2-3.3 (m, 2H), 3.55 (d, 2H), 3.8 (t, 2H), 4.0 (d, 2H), 4.08 (s, 3H), 4.32 (t, 2H), 7.4 (s, 1H), 7.88 (dd, 1H), 8.2 (d, 1H), 8.4 (s, 1H), 8.55 (s, 1H), 8.85 (s, 1H), 9.42 (s, 1H)
- 25 d) 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (74mg) was reacted with 6-amino-2-methylthiazole (57mg) to give **6-methoxy-4-(2-methyl-1,3-benzothiazol-5-ylamino)-7-(3-morpholinopropoxy)quinazoline**.
- ¹H NMR Spectrum: (DMSO-d₆) 2.3-2.4 (m, 2H), 2.85 (s, 3H), 3.05-3.2 (m, 2H), 3.3 (t, 2H), 3.4-3.5 (m, 2H), 3.85 (t, 2H), 4.0 (d, 2H), 4.05 (s, 3H), 4.35 (t, 2H), 7.42 (s, 1H), 7.75 (dd,

1H), 8.15 (d, 1H), 8.3 (s, 1H), 8.42 (s, 1H), 8.85 (s, 1H)

e) 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (74mg) was reacted with 5-aminoindan (32mg) to give **4-(2,3-dihydro-1H-inden-5-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.08 (m, 2H), 2.3-2.4 (m, 2H), 2.9 (m, 4H), 3.05-3.2 (m, 2H), 3.2-3.3 (m, 2H), 3.5 (d, 2H), 3.82 (t, 2H), 4.0 (d, 2H), 4.05 (s, 3H), 4.3 (t, 2H), 7.32 (d, 1H), 7.4 (m, 2H), 7.55 (s, 1H), 8.32 (s, 1H), 8.8 (s, 1H)

10 Example 30

A suspension of 4-chloro-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (130mg, 0.4mmol), (prepared as described for the starting material in Example 10), 7-hydroxy-2-methylchromone (88mg, 0.5 mmol), (Bull Soc. Chim. Fr. 1995, 132, 233), and potassium carbonate (83mg, 0.6 mmol) was heated at 100°C for 1.5 hours. After cooling, the mixture was partitioned between water and ethyl acetate. The organic layer was washed with water, brine, dried (MgSO₄), and the volatiles were removed by evaporation. The residue was triturated with ether, collected by filtration, washed with ether and dried under vacuum to give **6-methoxy-4-(2-methyl-4-oxo-4H-chromen-7-yloxy)-7-(1-methylpiperidin-4-ylmethoxy)quinazoline** (170mg, 92%).

20 MS - ESI: 462 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5 (m, 2H); 1.75-1.95 (m, 5H); 2.2 (s, 3H), 2.42 (s, 3H); 4.0 (s, 3H); 4.1 (d, 2H); 6.3 (s, 2H); 7.4 (s, 1H); 7.45 (dd, 1H); 7.6 (s, 1H); 7.7 (s, 1H); 8.15 (d, 1H); 8.61 (s, 1H)

25 Examples 31-33

Using an analogous procedure to that described in Example 30, the compounds described in Table II hereinafter and detailed in the notes a)-c) to Table II, were made.

Table II




Table II

Example No.	Weight (mg)	yield %	MS-ESI [MH] ⁺	note	Q	R
31	180	85	451	a	1-methylpiperidin-4-ylmethoxy	4-methyl-3,4-dihydro-2 <i>H</i> -1,4-benzoxazin-6-yloxy
32	160	87	462	b	3-pyrrolidin-1-ylpropoxy	2-methyl-4-oxo-4 <i>H</i> -chromen-7-yloxy
33	100	56	451	c	3-pyrrolidin-1-ylpropoxy	4-methyl-3,4-dihydro-2 <i>H</i> -1,4-benzoxazin-6-yloxy

a) 4-Chloro-6-methoxy-7-(1-methylpiperidin-4-yloxy)quinazoline (130mg), (prepared as described for the starting material in Example 10), was reacted with 3,4-dihydro-4-methyl-2*H*-1,4-benzoxazin-6-ol (83mg), (J. Org. Chem. 1971, 36 (1)), to give **6-methoxy-4-(4-methyl-3,4-dihydro-2*H*-1,4-benzoxazin-6-yloxy)-7-(1-methylpiperidin-4-ylmethoxy)quinazoline**

¹H NMR Spectrum: (DMSO-d₆) 1.6-1.75 (m, 2H); 1.9-2.3 (m, 5H); 2.8 (s, 3H); 2.9 (s, 3H); 3.0-3.15 (m, 2H); 3.3 (br s, 2H); 3.5-3.6 (d, 2H); 4.1 (s, 3H); 4.2 (d, 2H); 4.3 (t, 2H); 6.55 (m, 1H); 6.75 (s, 1H); 6.8 (d, 1H); 7.6 (s, 1H); 7.75 (s, 1H); 9.15 (s, 1H)

b) 4-Chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (130mg), (prepared as described for the starting material in Example 9), was reacted with 7-hydroxy-2-methylchromone (88mg), (Bull Soc. Chim Fr. 1995, 132, 233). After cooling, water was added (20ml) and the precipitate was collected by filtration and dried under vacuum over phosphorus pentoxide at 60°C to give **6-methoxy-4-(2-methyl-4-oxo-4*H*-chromen-7-yloxy)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.8-2.0 (m, 2H); 2.0-2.15 (m, 2H); 2.2-2.3 (m, 2H); 2.4 (s, 3H); 3.05-3.15 (m, 2H); 3.3-3.4 (m, 2H); 3.6-3.7 (m, 2H); 4.05 (s, 3H); 4.35 (t, 2H); 6.3 (s, 1H); 7.45 (d, 1H); 7.5 (s, 1H); 7.65 (s, 1H); 7.72 (s, 1H); 8.15 (d, 1H); 8.75 (s, 1H)

c) 4-Chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (130mg), (prepared as described for the starting material in Example 9), was reacted with 3,4-dihydro-4-methyl-2H-1,4-benzoxazin-6-ol (83mg), (J. Org. Chem. 1971, 36 (1)), to give **6-methoxy-4-(4-methyl-3,4-dihydro-2H-1,4-benzoxazin-6-yloxy)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline**.

5 ¹H NMR Spectrum: (DMSO-d₆) 1.85-2.0 (m, 2H); 2.0-2.15 (m, 2H); 2.25-2.35 (m, 2H); 2.83 (s, 3H); 3.05-3.15 (m, 2H); 3.3 (t, 2H); 3.4 (t, 2H); 3.7 (br m, 2H); 4.1 (s, 3H); 4.3 (t, 2H); 4.4 (t, 2H); 6.52 (d, 1H); 6.7 (s, 1H); 6.8 (d, 1H); 7.55 (s, 1H); 7.75 (s, 1H); 9.1 (s, 1H)

Example 34

10 A solution of 4-chloro-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (110mg, 0.34mmol), (prepared as described for the starting material in Example 10), and 5-hydroxyindole (55mg, 0.41mmol) in DMF (1.5ml) containing potassium carbonate (70mg, 0.51mmol) was heated at 100°C for 2 hours. After cooling, water was added and the precipitate was collected by filtration, washed with water followed by ether, and dried under
15 vacuum over phosphorus pentoxide to give **4-(indol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline** (90mg, 64%).

MS - ESI: 419 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.35-1.5 (m, 2H); 1.8 (d, 2H); 1.95 (t, 2H); 1.7-2.0 (m, 1H); 2.2 (s, 3H); 2.85 (d, 2H); 4.02 (s, 3H); 4.1 (d, 2H); 6.45 (s, 1H); 7.0 (d, 1H); 7.35 (s, 1H); 7.4-
20 7.5 (m, 3H); 7.6 (s, 1H); 8.5 (s, 1H)

Elemental analysis:

Found C 67.4 H 6.5 N 13.1

C₂₄H₂₆N₄O₃ · 0.5H₂O

Requires C 67.4 H 6.4 N 13.1%

Example 35

25 Using an analogous procedure to that described in Example 34, 4-chloro-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (110mg, 0.34mmol), (prepared as described for the starting material in Example 10), was reacted with 2,3-dimethyl-5-hydroxyindole (66mg, 0.41mmol), (Arch. Pharm. 1972, 305, 159). The crude product was purified by column chromatography eluting with methanol/methylene chloride (1/9) followed by 3M ammonia in
30 methanol/methanol/methylene chloride (5/15/80) to give **4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline** (60mg, 40%).

MS - ESI: 447 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.2-1.4 (m, 2H); 1.7 (d, 2H); 1.8 (t, 2H); 1.7-1.9 (m, 1H); 2.05 (s, 3H); 2.12 (s, 3H); 2.25 (s, 3H); 2.75 (d, 2H); 3.9 (s, 3H); 4.0 (d, 2H); 6.8 (d, 1H); 7.15 (s, 1H); 7.2 (d, 1H); 7.3 (s, 1H); 7.52 (s, 1H); 8.45 (s, 1H)

Elemental analysis:

Found C 68.6 H 6.9 N 12.5

5 C₂₆H₃₀N₄O₃ · 0.4H₂O

Requires C 68.8 H 6.8 N 12.4%

Example 36

Using an analogous procedure to that described in Example 34, 4-chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (110mg, 0.34mmol), (prepared as described for the starting material in Example 9), was reacted with 5-hydroxyindole (55mg, 0.41mmol). The crude product was purified by chromatography on alumina, eluting with methanol/ethyl acetate/methylene chloride (5/45/50) to give 4-(indol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (70mg, 50%).

MS - ESI 419 [MH]⁺

15 ¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.9-2.0 (m, 2H); 2.1 (m, 2H); 2.3 (t, 2H); 3.0-3.15 (m, 2H); 3.4 (t, 2H); 3.6-3.75 (m, 2H); 4.1 (s, 3H); 4.4 (t, 2H); 6.5 (s, 1H); 7.05 (d, 1H); 7.5 (s, 1H); 7.5-7.6 (m, 2H); 7.85 (s, 1H); 9.11 (s, 1H)

Elemental analysis:

Found C 63.7 H 6.4 N 12.1

20 C₂₄H₂₆N₄O₃ · 1.9H₂O

Requires C 63.7 H 6.6 N 12.4%

Example 37

A suspension of 4-chloro-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (100mg, 0.31mmol), (prepared as described for the starting material in Example 10), and 5-amino-2,3-dimethylindole (55mg, 0.34mmol) in isopropanol (6ml) containing 5.5M hydrogen chloride in isopropanol (60μL) was heated for 30 minutes at 70°C. After cooling, the solid was collected by filtration, washed with isopropanol, followed by ether and dried under vacuum to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline hydrochloride (118mg, 74%).

MS - ESI: 446 [MH]⁺

30 ¹H NMR Spectrum: (DMSO-d₆): 1.8-1.9 (m, 2H); 2.0 (d, 2H); 2.1-2.2 (m, 1H); 2.16 (s, 3H); 2.33 (s, 3H); 2.75 (br s, 3H); 2.95-3.05 (m, 2H); 3.5 (d, 2H); 4.0 (s, 3H); 4.07 (d, 2H); 7.25 (d, 1H); 7.4 (d, 1H); 7.42 (s, 1H); 7.52 (s, 1H); 8.25 (s, 1H); 8.75 (s, 1H); 10.0 (br s, 1H); 10.9 (s,

1H); 11.25 (br s, 1H)

Elemental analysis:

Found C 58.5 H 6.8 N 12.9

C₂₆H₃₁N₅O₂ 1H₂O 1.9HCl

Requires C 58.6 H 6.6 N 13.1%

5 **Example 38**

Using an analogous procedure to that described in Example 37, 4-chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (100mg, 0.31mmol), (prepared as described for the starting material in Example 9), was reacted with 5-amino-2,3-dimethylindole (55mg, 0.34mmol) to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline hydrochloride** (114mg, 72%).

MS - ESI: 446 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.85-2.0 (m, 2H); 2.05-2.15 (m, 2H); 2.1 (s, 3H); 2.2 (s, 3H); 2.25-2.35 (m, 2H); 2.35 (s, 3H); 3.0-3.15 (m, 2H); 3.32-3.42 (m, 2H); 3.6-3.7 (m, 2H); 4.05 (s, 3H); 4.3 (t, 2H); 7.2 (d, 1H); 7.3 (s, 1H); 7.35 (d, 1H); 7.57 (s, 1H); 8.2 (s, 1H); 8.8 (s, 1H)

Elemental analysis:

Found C 58.8 H 7.0 N 12.5

C₂₆H₃₁N₅O 1.9H₂O 1.9HCl 0.1isopropanol

Requires C 58.6 H 7.1 N 12.9%

Example 39

Using an analogous procedure to that described in Example 38, 4-chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (100mg, 0.31mmol), (prepared as described for the starting material in Example 9), was reacted with 5-amino-2-methylindole (50mg, 0.34mmol) to give **6-methoxy-4-(2-methylindol-5-ylamino)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline hydrochloride** (138mg, 89%).

MS - ESI: 432 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.8-1.9 (m, 2H); 2.0-2.1 (m, 2H); 2.15-2.35 (m, 2H); 2.4 (s, 3H); 3.0-3.1 (m, 2H); 3.2-3.3 (m, 2H); 3.5-3.6 (m, 2H); 4.0 (s, 3H); 4.32 (t, 2H); 6.2 (s, 1H); 7.2 (d, 1H); 7.3 (m, 2H); 7.65 (s, 1H); 8.25 (s, 1H); 8.75 (s, 1H); 10.75 (br s, 1H); 11.15 (s, 1H); 11.25 (br s, 1H)

Elemental analysis:

Found C 58.9 H 6.6 N 13.5

C₂₅H₂₉N₅O₂ 2.2HCl 0.1isopropanol

Requires C 58.7 H 6.2 N 13.5%

Example 40

A mixture of 4-chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (100mg, 0.31mmol), (prepared as described for the starting material in Example 9), and 7-hydroxy-2,4-dimethylquinoline (64mg, 0.36mmol), (Chem. Berichte, 1903, 36, 4016), in DMF (3ml) containing potassium carbonate (86mg, 0.62mmol) was heated at 90°C for 3 hours. After cooling, the mixture was poured onto a column of silica and eluted with 2.5M ammonia in methanol/methylene chloride (5/95) to give **4-(2,4-dimethylquinolin-7-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline** (50mg, 35%).

MS - ESI: 459 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 1.8 (br s, 4H); 2.2 (m, 4H); 2.55 (br s, 4H); 2.7 (2s, 6H); 2.68 (m, 2H); 4.05 (s, 3H); 4.3 (t, 2H); 7.15 (s, 1H); 7.35 (s, 1H); 7.45 (d, 1H); 7.6 (s, 1H); 7.9 (s, 1H); 8.05 (d, 1H); 8.6 (s, 1H)

Elemental analysis:	Found	C 70.4	H 7.1	N 12.1
C ₂₇ H ₃₀ N ₄ O ₃ 0.2ether	Requires	C 70.5	H 6.8	N 11.8%

Example 41

Using an analogous procedure to that described in Example 37, 4-chloro-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (50mg, 0.155mmol), (prepared as described for the starting material in Example 10), was reacted with 5-amino-2-methylindole (0.171mmol) to give **6-methoxy-4-(2-methylindol-5-ylamino)-7-(1-methylpiperidin-4-ylmethoxy)quinazoline hydrochloride** (72mg, quant.).

MS - ESI: 432 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.5-1.7 (m, 2H); 2.05 (d, 2H); 2.1-2.2 (m, 1H); 2.45 (s, 3H); 2.8 (s, 3H); 3.05 (t, 2H); 3.5 (d, 2H); 4.0 (s, 3H); 4.1 (d, 2H); 6.2 (s, 1H); 7.2 (d, 1H); 7.32 (d, 1H); 7.4 (d, 1H); 7.6 (s, 1H); 8.2 (s, 1H); 8.85 (s, 1H)

Elemental analysis:	Found	C 53.9	H 6.8	N 12.4
C ₂₅ H ₂₉ N ₅ O ₂ 2.6H ₂ O 2.07HCl	Requires	C 54.2	H 6.6	N 12.6%

Example 42

A suspension of 4-chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (100mg, 0.31 mmol), (prepared as described for the starting material in Example 9), and 7-hydroxy-2-methylquinoline (54mg, 0.34mmol), (J. Med. Chem. 1998, 41, 4062), in DMF

(3ml) containing potassium carbonate (86mg, 0.62mmol) was heated at 90°C for 2 hours. After cooling, the mixture was partitioned between ethyl acetate and water. The organic layer was separated, washed with water, brine, dried and the volatiles were removed by evaporation. The residue was triturated with minimal ether, collected by filtration and dried under vacuum to give **6-methoxy-4-(2-methylquinolin-7-yloxy)-7-(3-pyrrolidin-1-ylpropoxy)quinazoline** (95mg, 69%).

MS - ESI: 445 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 1.8 (br s, 4H); 2.2 (m, 2H); 2.5 (br s, 4H); 2.7 (t, 2H); 2.8 (s, 3H); 4.1 (s, 3H); 4.3 (t, 2H); 7.3 (d, 1H); 7.35 (s, 1H); 7.45 (dd, 1H); 7.6 (s, 1H); 7.85 (d, 1H); 7.9 (s, 1H); 8.1 (d, 1H); 8.6 (s, 1H)

Example 43

Using an analogous procedure to that described in Example 42, 4-chloro-6-methoxy-7-(1-(2-methylsulphonyl)ethyl)piperidin-4-ylmethoxy)quinazoline (156mg, 0.38mmol), (prepared as described for the starting material in Example 12), was reacted with 7-hydroxy-2-methylquinoline (66mg, 0.4 mmol), (J. Med. Chem. 1998, 41, 4062), to give **6-methoxy-7-(1-(2-methylsulphonyl)ethyl)piperidin-4-ylmethoxy)-4-(2-methylquinolin-7-yloxy)quinazoline** (166mg, 82%).

MS - ESI: 537 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5 (m, 2H); 1.75-1.95 (m, 3H); 1.95-2.15 (m, 2H); 2.7 (s, 3H); 2.7-2.8 (m, 2H); 2.9-3.0 (m, 2H); 3.05 (s, 3H); 3.2-3.35 (m, 2H); 4.02 (s, 3H); 4.1 (d, 2H); 7.4 (s, 1H); 7.45 (d, 1H); 7.55 (d, 1H); 7.65 (s, 1H); 7.8 (s, 1H); 8.05 (d, 1H); 8.35 (d, 1H); 8.55 (s, 1H)

Elemental analysis:

Found C 62.2 H 6.3 N 10.4

C₂₈H₃₂N₄O₅S 0.35ether 0.2DMF

Requires C 62.4 H 6.4 N 10.2%

Example 44

A suspension of 4-chloro-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (50mg, 0.155mmol), (prepared as described for the starting material in Example 10), and 5-hydroxy-2-trifluoromethylindole (34mg, 0.17mmol) in DMF (1.5ml) containing potassium carbonate (43mg, 0.31mmol) was heated at 90°C for 2 hours. After cooling, the mixture was partitioned between ethyl acetate and water. The organic layer was separated, washed with

brine, dried (MgSO_4) and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with methanol/ethyl acetate/methylene chloride (10/50/40) followed by 2.5M ammonia in methanol/ethyl acetate/methylene chloride (10/50/40) to give 6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)-4-(2-

5 trifluoromethylindol-5-yloxy)quinazoline (35mg, 48%).

MS - ESI: 487 $[\text{MH}]^+$

^1H NMR Spectrum: (DMSO-d_6) 1.25-1.4 (m, 2H); 1.75 (d, 2H); 1.8 (t, 2H); 1.7-2.0 (m, 1H); 2.2 (s, 3H); 2.75 (d, 2H); 4.0 (s, 3H); 4.1 (d, 2H); 7.0 (s, 1H); 7.25 (d, 1H); 7.4 (s, 1H); 7.6 (d, 1H); 7.8 (s, 1H); 8.5 (s, 1H); 12.5 (s, 1H)

10	Elemental analysis:	Found	C 60.2	H 5.8	N 10.9
	$\text{C}_{25}\text{H}_{25}\text{F}_3\text{N}_4\text{O}_3 \cdot 0.7\text{H}_2\text{O} \cdot 0.2\text{ether}$	Requires	C 60.3	H 5.6	N 10.9%

The starting material was prepared as follows:

15 A solution of (4-methoxy-2-methylphenyl)-carbamic acid-1,1-dimethylethyl ester (2g, 8.43mmol), (J. Med. Chem. 1996, 39, 5119), in dry THF (25ml) was cooled to -40°C and sec-butyllithium (15ml, 19.5mmol) was added. After stirring for 15 minutes at this temperature, N-methyl-N-methoxytrifluoroacetamide (1.32g, 8.43mmol) in THF (20ml) was added in portions. Stirring was continued for 1 hour at -40°C and then the mixture was allowed to warm to ambient temperature. The mixture was poured onto ether/1M hydrochloric acid. The
20 organic layer was separated, washed with water, brine, dried (MgSO_4) and the volatiles were removed by evaporation.

The crude residue (1.4g) was dissolved in methylene chloride (8ml) and TFA was added (1.5ml). After stirring for 3 hours at ambient temperature, the volatiles were removed under vacuum. The crude product was partitioned between methylene chloride and water. The
25 organic layer was separated, washed with water, brine, dried (MgSO_4) and the volatiles were removed by evaporation. The residue was purified by column chromatography, eluting with ether/petroleum ether (1/9) to give 5-methoxy-2-trifluoromethylindole (845mg, 47% over 2 steps).

^1H NMR Spectrum: (CDCl_3) 3.83 (s, 3H), 6.82 (s, 1H), 7.0 (dd, 1H), 7.1 (s, 1H), 7.3 (d, 1H),
30 8.15 (br s, 1H)

A solution of 5-methoxy-2-trifluoromethylindole (800mg, 3.7mmol) in methylene chloride (6ml) was cooled to -15°C and a solution of 1M boron tribromide in methylene

chloride (7.44 ml, 7.4mmol) was added in portions. The mixture was allowed to warm to ambient temperature and was stirred for 45 minutes. After cooling to 0°C, saturated aqueous sodium hydrogen carbonate (25ml) was added. The mixture was extracted with ethyl acetate. The organic layer was dried (MgSO₄) and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with ethyl acetate/petroleum ether. After removal of the volatiles by evaporation, the solid was triturated with pentane, collected by filtration and dried under vacuum to give 5-hydroxy-2-trifluoromethylindole (290mg, 39%).

MS - EI: 201 [M.]⁺

¹H NMR Spectrum: (CDCl₃) 4.64 (s, 1H), 6.8 (s, 1H), 6.92 (dd, 1H), 7.1 (s, 1H), 7.3 (d, 1H), 8.3 (br s, 1H)

Elemental analysis:

Found C 53.3 H 2.9 N 6.8

C₉H₆F₃NO 0.1 H₂O

Requires C 53.3 H 3.1 N 6.9%

15 **Example 45**

Using an analogous procedure to that described in Example 44, 4-chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (100mg, 0.3mmol), (prepared as described for the starting material in Example 9), was reacted with 5-hydroxy-2-trifluoromethylindole (75mg, 0.37 mmol), (prepared as described for the starting material in Example 44), to give 6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline (105mg, 70%).

MS - ESI: 487 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 1.8 (m, 4H); 2.1-2.3 (m, 2H); 2.55 (br s, 4H); 2.7 (t, 2H); 4.1 (s, 3H); 4.3 (t, 2H); 6.95 (s, 1H); 7.2 (dd, 1H); 7.35 (s, 1H); 7.5 (d, 1H); 7.55 (s, 1H); 7.6 (s, 1H); 8.6 (s, 1H); 8.8 (s, 1H)

Elemental analysis:

Found C 61.7 H 5.5 N 11.5

C₂₅H₂₅F₃N₄O₃

Requires C 61.7 H 5.2 N 11.5%

Example 46

Using an analogous procedure to that described in Example 42, 4-chloro-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (100mg, 0.31mmol), (prepared as described for the starting material in Example 10), was reacted with 7-hydroxy-2-methylquinoline (54mg,

0.34mmol), (J. Med. Chem. 1998, **41**, 4062), to give **6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)-4-(2-methylquinolin-7-yloxy)quinazoline** (86mg, 63%).

MS - ESI: 445 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 1.4-1.6 (m, 2H); 1.95 (d, 2H); 2.05 (t, 2H); 1.9-2.1 (m, 1H); 2.35 (s, 3H); 2.8 (s, 3H); 2.95 (d, 2H); 4.1 (s, 3H); 4.15 (d, 2H); 7.3 (m, 2H); 7.45 (dd, 1H); 7.6 (s, 1H); 7.9 (d, 1H); 7.95 (s, 1H); 8.1 (d, 1H); 8.6 (s, 1H)

Elemental analysis:

Found C 69.7 H 6.5 N 12.8

C₂₆H₂₈N₄O₃ 0.2H₂O

Requires C 69.7 H 6.4 N 12.5%

10 **Example 47**

A suspension of 4-chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (110mg, 0.34 mmol), (prepared as described for the starting material in Example 9), and 2,3-dimethyl-5-hydroxyindole (66mg, 0.41mmol), (Arch. Pharm. 1972, **305**, 159), in DMF (1.5ml) containing potassium carbonate (70mg, 0.51mmol) was heated at 100°C for 2 hours. After cooling, the residue was purified by chromatography, eluting with methanol/methylene chloride (1/9) followed by 2.5M ammonia in methanol/methanol/methylene chloride (5/10/85) to give **4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline** (50mg, 33%).

MS - ESI: 447 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.9-2.0 (m, 2H); 2.05-2.15 (m, 2H); 2.15 (s, 3H); 2.3-2.4 (m, 2H); 2.4 (s, 3H); 3.05-3.15 (m, 2H); 3.35-3.45 (t, 2H); 3.7 (br s, 2H); 4.1 (s, 3H); 4.4 (t, 2H); 6.95 (d, 1H); 7.3 (s, 1H); 7.35 (d, 1H); 7.55 (s, 1H); 7.85 (s, 1H); 9.15 (s, 1H)

Elemental analysis:

Found C 67.7 H 6.8 N 12.2

C₂₆H₃₀N₄O₃ 0.8H₂O

Requires C 67.8 H 6.9 N 12.2%

Example 48

Using an analogous procedure to that described in Example 32, 7-benzyloxy-4-chloro-6-methoxyquinazoline (1g, 3.33mmol), (prepared as described for the starting material in Example 1), was reacted with 5-hydroxy-2-methylindole (0.59g, 4mmol) to give **7-benzyloxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (1.25g, 91%).

MS - ESI: 412 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H); 4.0 (s, 3H); 5.35 (s, 2H); 6.15 (s, 1H); 6.85 (s,

1H); 7.2-7.6 (m, 9H); 8.5 (s, 1H)

Elemental analysis:

Found C 72.2 H 5.1 N 10.2

C₂₅H₂₁N₃O₃ · 0.2H₂O

Requires C 72.3 H 5.2 N 10.1%

5 The starting material may be prepared as follows:

A solution of boron tribromide (32.5ml, 341mmol) in methylene chloride (60ml) was added in portions to a solution of 5-methoxy-2-methylindole (25g, 155mmol) in methylene chloride (250ml) cooled at -45°C. After stirring for 15 minutes at -30°C, the mixture was warmed up to ambient temperature and stirred for 1 hour. Methylene chloride (300ml) was added in portions and the mixture was cooled to 0°C. Water was added in portions and the mixture was adjusted to pH6 with 4N sodium hydroxide. The organic layer was separated. The aqueous layer was extracted with methylene chloride and the organic layers were combined, washed with water, brine, dried (MgSO₄) and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with ethyl acetate/methylene chloride (1/9 followed by 15/85) to give 5-hydroxy-2-methylindole (21.2g, 93%).

¹H NMR Spectrum: (DMSO-d₆) 2.35 (s, 3H) ; 5.95 (s, 1H) ; 6.5 (dd, 1H) ; 6.7 (s, 1H) ; 7.05 (d, 1H) ; 8.5 (s, 1H)

20 Example 49

A solution of 7-benzyloxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (0.2g, 0.5mmol), (prepared as described in Example 48), in a mixture of methylene chloride (5ml) and DMF (2ml) containing 10% palladium-on-charcoal (50mg) was treated with hydrogen at 1.8 atmospheres pressure for 2 hours. The suspension was filtered and the catalyst was washed with methanol followed by methylene chloride. The volatiles were removed from the filtrate by evaporation. The residue was triturated with water. The resulting solid was washed with water and dried under vacuum over phosphorus pentoxide at 60°C to give **7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (140mg, 89%).

MS - ESI: 322 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H); 4.0 (s, 3H); 6.15 (s, 1H); 6.9 (d, 1H); 7.2 (s, 1H); 7.25 (s, 1H); 7.3 (d, 1H); 7.6 (s, 1H); 8.4 (s, 1H)

Example 50

A suspension of 4-chloro-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline (150mg, 0.45mmol) and 5-hydroxy-2-trifluoromethylindole (109mg, 0.54mmol), (prepared as described for the starting material in Example 44), in DMF (1.5ml) containing potassium carbonate (94mg, 0.67mmol) was heated at 100°C for 1 hour. After cooling, the precipitate was collected by filtration, washed with ether, and dried under vacuum to give 6-methoxy-7-(3-methylsulphonylpropoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline (195mg, 87%).

MS - ESI: 496 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.25-2.4 (m, 2H), 3.1 (s, 3H), 3.35 (t, 2H), 4.1 (s, 3H), 4.4 (t, 2H), 7.1 (s, 1H), 7.3 (d, 1H), 7.5 (s, 1H), 7.6 (d, 1H), 7.7 (s, 1H), 7.78 (s, 1H), 8.9 (s, 1H)

The starting material was prepared as follows:

A solution of 3-(methylthio)-1-propanol (5.3g, 50mmol) in methanol (500ml) was added to a solution of OXONE, (trade mark of E.I. du Pont de Nemours & Co., Inc), (30g) in water (150ml) and the mixture stirred at ambient temperature for 24 hours. The precipitated solid was removed by filtration and the methanol removed from the filtrate by evaporation. The aqueous residue was saturated with sodium chloride and extracted with methylene chloride (4x25ml). The aqueous residue was then saturated with ammonium chloride and extracted with ethyl acetate (4x25ml). The extracts were combined, dried (MgSO₄) and the solvent removed by evaporation to give 3-(methylsulphonyl)-1-propanol (610mg, 9%) as an oil.

¹H NMR Spectrum: (CDCl₃) 2.10(m, 2H); 2.96(s, 3H); 3.20(t, 2H); 3.80(t, 2H)

MS - ESI: 139 [MH]⁺

Alternatively the 3-(methylsulphonyl)-1-propanol may be prepared as follows:

m-Chloroperoxybenzoic acid (67%, 25 g, 97.2 mmol) was added in portions to 3-(methylthio)-1-propanol (5 ml, 48.6 mmol) in solution in dichloromethane. Some m-chlorobenzoic acid precipitated out and was removed by filtration. The filtrate was evaporated and the residue was purified over alumina using first dichloromethane (100%) then dichloromethane/methanol (95/5) to give 3-(methylsulphonyl)-1-propanol (4.18 g, 62%) as an oil.

Triphenylphosphine (8.9g, 35.2mmol) was added to a suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (6g, 19.6mmol), (prepared as described for the starting material in Example 12), in methylene chloride (150ml). This was followed by the addition of 3-(methylsulphonyl)-1-propanol (3.5g, 25.4mmol) and diethyl azodicarboxylate (5.55ml, 35.2mmol) in portions. The reaction was complete once the reaction became homogeneous. Silica was added and the volatiles were removed by evaporation. The free flowing powder was placed on the top of a flash chromatography column pre-equilibrated with ethyl acetate (100%). Elution was done using ethyl acetate (100%) followed by methylene chloride/ethyl acetate/methanol (60/35/5). The volatiles were removed by evaporation to give 6-methoxy-7-(3-methylsulphonylpropoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (7.58 g, 91%) as a white solid.

¹H NMR Spectrum: (CDCl₃) 1.2(s, 9H); 2.4-2.5(m, 2H); 3.0(s, 3H); 3.25-3.35(t, 2H); 5.95(s, 1H); 7.1(s, 1H); 7.65(s, 1H); 8.2(s, 1H)

6-Methoxy-7-(3-methylsulphonylpropoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (7g, 17mmol) was suspended in methanol and 2M sodium hydroxide (3.3ml, 6.6mmol) was added with continuous stirring. The reaction mixture became homogeneous after 15 minutes. After a further 45 minutes water was added (7ml) and the reaction mixture was adjusted to pH10 with 2M hydrochloric acid. The precipitate (a white solid) was collected by filtration, washed with water and dried over phosphorus pentoxide under vacuum to give 6-methoxy-7-(3-methylsulphonylpropoxy)-3,4-dihydroquinazolin-4-one (5 g, 90%).


¹H NMR Spectrum: (DMSO-d₆) 2.2-2.3(m, 2H); 3.05(s, 3H); 3.35(t, 2H); 3.9(s, 3H); 4.25(t, 2H); 7.15(s, 1H); 7.5(s, 1H); 8.0(s, 1H)

6-Methoxy-7-(3-methylsulphonylpropoxy)-3,4-dihydroquinazolin-4-one (3.6g, 11.5mmol) was suspended in thionyl chloride (40ml). DMF (1.8ml) was added under argon and the mixture was heated at reflux for 1.5 hours. The thionyl chloride was eliminated by several azeotropic distillations using toluene. The solid residue was suspended in ice/water and a saturated solution of sodium hydrogen carbonate was added to adjust the mixture to pH7. The solid was collected by filtration, washed with water and dried in a vacuum dessicator over phosphorus pentoxide to give 4-chloro-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline (3.35g, 88%).

Examples 51-52

Using an analogous procedure to that described in Example 50, 4-chloro-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline, (prepared as described for the starting material in Example 50), was reacted with the appropriate phenol to give the compounds described in

5 Table III:

Table III

Example No.	Weight (mg)	Yield %	MS-ESI [MH] ⁺	Ar	Note
51	189	92	454	2-methylquinolin-7-yl	a
52	175	90	428	indol-5-yl	b

10

a) 4-Chloro-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline (150mg, 0.45mmol) was reacted with 7-hydroxy-2-methylquinoline (86.6mg, 0.54mmol), (J. Med. Chem. 1998, 41, 4062). After cooling, water was added and the precipitate was collected by filtration, washed with water, followed by ether and dried under vacuum to give 6-methoxy-7-(3-

15 **methylsulphonylpropoxy)-4-(2-methylquinolin-7-yloxy)quinazoline.**

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.2-2.35 (m, 2H), 2.95 (s, 3H), 3.1 (s, 3H), 3.35 (m, 2H), 4.05 (s, 3H), 4.4 (t, 2H), 7.5 (s, 1H), 7.7 (s, 1H), 7.95 (dd, 1H), 8.02 (d, 1H), 8.2 (s, 1H), 8.48 (d, 1H), 8.7 (s, 1H), 9.12 (d, 1H)

20 b) Using an analogous procedure to that described in note a), 4-chloro-6-methoxy-7-(3-(methylsulphonyl)propoxy)quinazoline (150mg, 0.45mmol) was reacted with 5-hydroxyindole (72.4mg, 0.54mmol) to give 4-(indol-5-yloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline.

25 ¹H NMR Spectrum: (DMSO-d₆) 2.2-2.35 (m, 2H), 3.1 (s, 3H), 3.3-3.4 (t, 2H), 4.0 (s, 3H), 4.4 (t, 2H), 6.5 (s, 1H), 7.0 (dd, 1H), 7.4 (s, 1H), 7.4-7.5 (m, 3H), 7.6 (s, 1H), 8.5 (s, 1H), 11.25

(s, 1H)

Example 53


0.5M Triphenylphosphine in methylene chloride and diisopropyl azodicarboxylate (150 μ l, 0.75mmol) were added in portions to a suspension of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (112mg, 0.35mmol), (prepared as described in Example 49), and *N,N*-dimethylethanolamine (62mg, 0.7mmol) in methylene chloride (2ml). After stirring for 2 hours at ambient temperature, the reaction mixture was poured onto an isolute® column (10g of silica) and eluted with ethyl acetate/methylene chloride (1/1) followed by methanol/ethyl acetate/methylene chloride (10/40/50), methanol/methylene chloride (10/90), and 3M ammonia in methanol/methanol/methylene chloride (5/15/80). After removal of the volatiles by evaporation, the residue was dissolved in the minimum amount of methylene chloride (about 3ml) and ether and petroleum ether (about 10ml) was added. The resulting precipitate was collected by filtration and dried under vacuum to give 7-(2-(*N,N*-dimethylamino)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (52mg, 38%).

MS - ESI: 393 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.25 (s, 6H), 2.4 (s, 3H), 2.75 (t, 2H), 4.0 (s, 3H), 4.3 (t, 2H), 6.15 (s, 1H), 6.87 (d, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 7.5 (s, 1H)

Examples 54-56

Using an analogous procedure to that described in Example 53, the appropriate alcohols were reacted with 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline, (prepared as described in Example 49), in analogous proportions to give the compounds described in Table IV:

Table IV

Example	Weight	Yield	MS-ESI	R	Note
---------	--------	-------	--------	---	------

No.	(mg)	%	[MH] ⁺		
54	25	17	419	2-pyrrolidin-1-ylethoxy	a
55	112	74	433	1-methylpiperidin-3-ylmethoxy	b
56	115	72	456	2-(<u>N</u> -methyl- <u>N</u> -(4-pyridyl)amino)ethoxy	c

a) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 1-(2-hydroxyethyl)pyrrolidine (81mg) to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-pyrrolidin-1-ylethoxy)quinazoline**.

5 ¹H NMR Spectrum: (DMSO-d₆) 1.65-1.8 (m, 4H), 2.4 (s, 3H), 2.6 (br s, 4H), 2.9 (t, 2H), 4.0 (s, 3H), 4.3 (t, 2H), 6.15 (s, 1H), 6.9 (d, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

10 b) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 1-methyl-3-piperidinemethanol (90mg) to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-methylpiperidin-3-ylmethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.45-2.2 (m, 7H), 2.18 (s, 3H), 2.4 (s, 3H), 2.6 (br d, 1H), 2.85 (br d, 1H), 4.0 (s, 3H), 4.1 (d, 2H), 6.15 (s, 1H), 6.9 (d, 1H), 7.25 (d, 1H), 7.3 (d, 1H), 7.35 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)


15

c) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 2-(N-methyl-N-(4-pyridyl)amino)ethanol (106mg), (EP 0359389), to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-(4-pyridyl)amino)ethoxy)quinazoline**.

20 ¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H), 3.1 (s, 3H), 3.9 (t, 2H), 3.97 (s, 3H), 4.4 (t, 2H), 6.15 (s, 1H), 6.75 (d, 2H), 6.87 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.35 (s, 1H), 7.6 (s, 1H), 8.15 (d, 2H), 8.5 (s, 1H)

Examples 57-66

25 Using an analogous procedure to that described in Example 53, except that ammonia in methanol was not necessary during the column chromatography, the appropriate alcohols were reacted with 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline, (prepared as described in Example 49), in analogous proportions to give the compounds described in Table V:

Table V

Example No.	Weight (mg)	Yield %	MS-ESI [MH] ⁺	R	Note
57	115	76	435	2-morpholinoethoxy	a
58	64	42	433	2-piperidinoethoxy	b
59	66	43	437	2-(N-(2-methoxyethyl)-N-methylamino)ethoxy	c
60	118	75	449	3-morpholinopropoxy	d
61	101	68	424	2-(2-methoxyethoxy)ethoxy	e
62	81	57	407	3-(N,N-dimethylamino)propoxy	f
63	160	92	497	3-(1,1-dioxothiomorpholino)propoxy	g
64	121	83	417	2-(1H-1,2,4-triazol-1-yl)ethoxy	h
65	38	22	492	2-(2-(4-methylpiperazin-1-yl)ethoxy)ethoxy	i
66	80	48	479	2-(2-morpholinoethoxy)ethoxy	j

5

a) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 4-(2-hydroxyethyl)morpholine (92mg) to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-morpholinoethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H), 2.5-2.7 (m, 4H), 2.8 (t, 2H), 3.6 (t, 4H), 4.0 (s, 3H), 4.35 (t, 2H), 6.15 (s, 1H), 6.87 (dd, 1H), 7.25 (s, 1H), 7.32 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

10

b) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 1-(2-hydroxyethyl)piperidine (90mg) to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-**

piperidinoethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.45 (m, 2H), 1.4-1.6 (m, 4H), 2.4 (s, 3H), 2.4-2.5 (m, 4H), 2.75 (t, 2H), 3.97 (s, 3H), 4.3 (t, 2H), 6.15 (s, 1H), 6.9 (d, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

5

c) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 2-(N-(2-methoxyethyl)-N-methylamino)ethanol (93mg) to give **6-methoxy-7-(2-(N-(2-methoxyethyl)-N-methylamino)ethoxy)-4-(2-methylindol-5-yloxy)quinazoline.**

¹H NMR Spectrum: (DMSO-d₆) 2.35 (s, 3H), 2.4 (s, 3H), 2.65 (t, 2H), 2.85 (t, 2H), 3.25 (s, 3H), 3.45 (t, 2H), 3.97 (s, 3H), 4.25 (t, 2H), 6.15 (s, 1H), 6.9 (dd, 1H), 7.25 (s, 1H), 7.32 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

10

The starting material was prepared as follows:

A mixture of 2-(methylamino)ethanol (5.4g, 72 mmol), 2-bromoethyl methyl ether
15 (10g, 72 mmol) and triethylamine (10ml, 72 mmol) in acetonitrile (70ml) was refluxed overnight. After cooling, the solid was filtered and the filtrate was evaporated. The residue was triturated with ether. The ether layer was separated and evaporated to give 2-(N-(2-methoxyethyl)-N-methylamino)ethanol (3g, 31%).

MS-EI : 134 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 2.35 (s, 3H) ; 2.6 (t, 2H) ; 2.65 (t, 2H) ; 3.35 (s, 3H) ; 3.5 (t, 2H)
20 ; 3.6 (t, 2H)

d) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 4-(3-hydroxypropyl)morpholine (102mg) to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline.**

25

¹H NMR Spectrum: (DMSO-d₆) 1.9-2.1 (m, 2H), 2.4 (s, 3H), 2.45 (t, 2H), 2.45-2.6 (s, 4H), 3.6 (t, 4H), 4.0 (s, 3H), 4.25 (t, 2H), 6.15 (s, 1H), 6.9 (d, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.38 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

30

The starting material was prepared as follows:

Morpholine (94g, 1.08mol) was added dropwise to a solution of 3-bromo-1-propanol (75g, 0.54mol) in toluene (750ml) and the reaction then heated at 80°C for 4 hours. The

mixture was allowed to cool to ambient temperature and the precipitated solid was removed by filtration. The volatiles were removed from the filtrate and the resulting yellow oil was purified by distillation at 0.4-0.7 mmHg to give 4-(3-hydroxypropyl)morpholine (40g, 50%) as a colourless oil.

5 b.p. 68-70°C (~0.5mmHg)

¹H NMR Spectrum: (DMSO-d₆) 1.65-1.78(m, 2H); 2.50(t, 4H); 2.60(t, 2H); 3.68(t, 4H); 3.78(t, 2H); 4.90(br d, 1H)

10 e) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 2-(2-methoxyethoxy)ethanol (84mg) to give **6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(2-methylindol-5-yloxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.42 (s, 3H), 3.27 (s, 3H), 3.5 (t, 2H), 3.65 (t, 2H), 3.85 (t, 2H), 4.0 (s, 3H), 4.32 (t, 2H), 6.15 (s, 1H), 6.9 (d, 1H), 7.3 (s, 1H), 7.35 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

15

f) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 3-(N,N-dimethylamino)propanol (72mg) to give **7-(3-N,N-dimethylaminopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline**.

20 ¹H NMR Spectrum: (DMSO-d₆) 1.9-2.0 (m, 2H), 2.17 (s, 6H), 2.4 (s, 3H), 3.98 (s, 3H), 4.22 (t, 2H), 6.14 (s, 1H), 6.88 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.35 (s, 1H), 7.6 (s, 1H), 8.47 (s, 1H)

25 g) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 3-(1,1-dioxothiomorpholino)-1-propanol (135mg), (prepared as described for the starting material in Example 5), to give **7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.9-2.0 (m, 2H), 2.38 (s, 3H), 2.65 (t, 2H), 2.9 (br s, 4H), 3.1 (br s, 4H), 3.96 (s, 3H), 4.25 (t, 2H), 6.12 (s, 1H), 6.85 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.37 (s, 1H), 7.56 (s, 1H), 8.46 (s, 1H)

30

h) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 2-(1*H*-1,2,4-triazol-1-yl)ethanol (79mg), (Ann. Phar. Fr. 1977, 35, 503-508), to give **6-methoxy-4-**

(2-methylindol-5-yloxy)-7-(2-(1*H*-1,2,4-triazol-1-yl)ethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.42 (s, 3H), 3.96 (s, 3H), 4.62 (m, 2H), 4.75 (m, 2H), 6.15 (s, 1H), 6.9 (dd, 1H), 7.27 (s, 1H), 7.32 (d, 1H), 7.47 (s, 1H), 7.63 (s, 1H), 8.03 (s, 1H), 8.51 (s, 1H), 8.60 (s, 1H)

5

i) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 2-(2-(4-methylpiperazin-1-yl)ethoxy)ethanol (132mg), (Arzneim. Forsch. 1966, 16, 1557-1560), to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(2-(4-methylpiperazin-1-yl)ethoxy)ethoxy)quinazoline.**

10 ¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.2-2.6 (m, 10H), 2.4 (s, 3H), 3.65 (t, 2H), 3.85 (t, 2H), 4.03 (s, 3H), 4.35 (m, 2H), 6.16 (s, 1H), 6.9 (dd, 1H), 7.3 (s, 1H), 7.35 (d, 1H), 7.4 (s, 1H), 7.61 (s, 1H), 8.5 (s, 1H)

15 j) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline was reacted with 2-(2-morpholinoethoxy)ethanol (123mg) to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(2-morpholinoethoxy)ethoxy)quinazoline.**

¹H NMR Spectrum: (DMSO-d₆) 2.40 (s, 3H), 2.4-2.5 (m, 4H), 2.4-2.6 (m, 2H), 3.55 (t, 4H), 3.6 (t, 2H), 3.85 (t, 2H), 3.97 (br s, 3H), 4.15 (br s, 2H), 6.15 (s, 1H), 6.9 (d, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.48 (s, 1H)

20

The starting material was prepared as follows:

2-(2-Chloroethoxy)ethanol (1.25g, 10mmol) was added to a mixture of morpholine (2.58g, 30mmol) and potassium carbonate (5.5g, 40mmol) in acetonitrile (50ml). The mixture was heated at reflux for 6 hours and then stirred for 18 hours at ambient temperature. The insolubles were removed by filtration and the volatiles were removed from the filtrate by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (95/5 followed by 90/10 and then 80/20) to give 2-(2-morpholinoethoxy)ethanol (600mg, 34%).

25

MS - (EI): 175 [M.]⁺

30 ¹H NMR Spectrum: (CDCl₃) 2.5(br s, 4H); 2.59(t, 2H); 3.6-3.85(m, 10H)

Example 67

A solution of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (100mg, 0.29mmol), 5-hydroxy-2-methylindole (53mg, 0.36mmol), (prepared as described for the starting material in Example 48), and potassium carbonate (62mg, 0.44mmol) in DMF (2ml) was heated at 85°C for 3 hours, followed by heating at 95°C for 2 hours. After cooling, ice/water (15ml) was added and the precipitate was collected by filtration and dried under vacuum. The solid was purified by column chromatography eluting with methylene chloride/methanol (95/5) followed by methylene chloride/methanol/3M ammonia in methanol (95/3/2) to give 6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline (71mg, 54%).

MS - ESI: 447 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.35-1.4 (m, 2H), 1.45-1.55 (m, 4H), 1.92-2.0 (m, 2H), 2.3-2.4 (m, 4H), 2.40 (s, 3H), 2.4-2.5 (m, 2H), 3.97 (s, 3H), 4.22 (t, 2H), 6.15 (s, 1H), 6.9 (d, 1H), 7.27 (s, 1H), 7.8 (d, 1H), 7.35 (s, 1H), 7.58 (s, 1H), 8.48 (s, 1H)

The starting material was prepared as follows:

Diethyl azodicarboxylate (3.9ml, 24.5mmol) was added in portions to a suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (5g, 16.3mmol), (prepared as described for the starting material in Example 12), 3-bromo-1-propanol (2.21ml, 24.5mmol) and triphenylphosphine (6.42g, 24.5mmol) in methylene chloride (50ml). After stirring for 2 hours at ambient temperature, the volatiles were removed under vacuum and the residue was purified by column chromatography eluting with methylene chloride followed by methylene chloride/methanol (95/5) to give 7-(3-bromopropoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (6g, 86%).

MS - ESI: 427-429 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.12 (s, 9H), 2.32 (t, 2H), 3.7 (t, 2H), 3.9 (s, 3H), 4.25 (t, 2H), 5.9 (s, 2H), 7.20 (s, 1H), 7.51 (s, 1H), 8.36 (s, 1H)

Elemental analysis:	Found	C 50.1	H 5.4	N 6.4
C ₁₈ H ₂₃ BrN ₂ O ₅ · 0.2H ₂ O	Requires	C 50.2	H 5.5	N 6.5%

A solution of 7-(3-bromopropoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (2.89g, 6.78mmol) in piperidine (10ml) was heated at 100°C for 1 hour. After cooling, the volatiles were removed under vacuum. The residue was dissolved in methylene chloride, and washed with saturated ammonium chloride and brine. The organic

layer was dried (MgSO_4) and the volatiles were removed by evaporation. The residue was dried under vacuum to give 6-methoxy-7-(3-piperidinopropoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (2.4g, 83%).

MS - ESI: 432 $[\text{MH}]^+$

- 5 ^1H NMR Spectrum: (DMSO-d_6) 1.15 (s, 9H), 1.35-1.5 (m, 1H), 1.6-1.8 (m, 3H), 1.8-1.9 (d, 2H), 2.2-2.3 (m, 2H), 2.95 (t, 2H), 3.25 (t, 2H), 3.55 (d, 2H), 3.95 (s, 3H), 4.25 (t, 2H), 5.94 (s, 2H), 7.24 (s, 1H), 7.56 (s, 1H), 8.46 (s, 1H)

- 10 A solution of 6-methoxy-7-(3-piperidinopropoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (2.35g, 5.45mmol) in 7M ammonia in methanol (50ml) was stirred overnight at ambient temperature. The volatiles were removed under vacuum and the residue was triturated with ether, filtered and washed with ether followed by ether/methylene chloride (1/1) and dried under vacuum to give 6-methoxy-7-(3-piperidinopropoxy)-3,4-dihydroquinazolin-4-one (1.65g, 95%).

MS - ESI: 318 $[\text{MH}]^+$

- 15 ^1H NMR Spectrum: (DMSO-d_6) 1.3-1.4 (m, 2H), 1.4-1.55 (m, 4H), 1.85-1.95 (m, 2H), 2.35 (br s, 4H), 2.4 (t, 2H), 3.9 (s, 3H), 4.15 (t, 2H), 7.11 (s, 1H), 7.44 (s, 1H), 7.9 (s, 1H)

Elemental analysis:

Found C 63.5 H 7.4 N 13.1

$\text{C}_{17}\text{H}_{23}\text{N}_3\text{O}_3 \cdot 0.2\text{H}_2\text{O}$

Requires C 63.6 H 7.4 N 13.0%

- 20 A solution of 6-methoxy-7-(3-piperidinopropoxy)-3,4-dihydroquinazolin-4-one (1.5g, 4.7mmol) in thionyl chloride (15ml) containing DMF (1.5ml) was heated at reflux for 3 hours. After cooling, the volatiles were removed under vacuum. The residue was azeotroped with toluene. The solid was partitioned between methylene chloride and sodium hydrogen carbonate. The aqueous layer was adjusted to pH10 with 6M aqueous sodium hydroxide. The organic layer was separated, washed with brine, dried (MgSO_4) and the volatiles were removed by evaporation. The residue was purified by column chromatography to give 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (1.21g, 76%).

MS - ESI: 336 $[\text{MH}]^+$

^1H NMR Spectrum: (DMSO-d_6) 1.35-1.45 (m, 2H), 1.5-1.6 (m, 4H), 1.9-2.05 (m, 2H), 2.4 (br s, 4H), 2.45 (t, 2H), 4.0 (s, 3H), 4.29 (t, 2H), 7.41 (s, 1H), 7.46 (s, 1H), 8.9 (s, 1H)

Example 68

Using an analogous procedure to that described in Example 67, 4-chloro-6-methoxy-7-

(3-piperidinopropoxy)quinazoline (100mg), (prepared as described for the starting material in Example 67), was reacted with 5-hydroxyindole (48mg, 0.36mmol) to give **4-(indol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline** (57mg, 45%).

MS - ESI: 433 [MH]⁺

- 5 ¹H NMR Spectrum: (DMSO-d₆) 1.4 (br s, 2H), 1.45-1.6 (br s, 4H), 1.9-2.1 (m, 2H), 2.4 (br s, 4H), 2.45 (t, 2H), 4.0 (s, 3H), 4.25 (t, 2H), 6.47 (s, 1H), 7.0 (d, 1H), 7.35 (s, 1H), 7.45 (s, 2H), 7.47 (d, 1H), 7.61 (s, 1H), 8.49 (s, 1H)

Example 69

- 10 A solution of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (161mg, 0.5mmol), (prepared as described in Example 49), 4-(4-methylphenylsulphonyloxymethyl)-1-tert-butoxycarbonylpiperidine (222mg, 0.6mmol), (prepared as described for the starting material in Example 10), and potassium carbonate (188mg, 1mol) in DMF (1.6ml) was heated at 100°C for 2 hours. After cooling, water was added. The precipitate was collected by
15 filtration, washed with water, and dried under vacuum over phosphorus pentoxide at 60°C. The solid was triturated with petroleum ether, collected by filtration, washed with a mixture of ether/petroleum ether (1/1) and dried under vacuum to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-tert-butoxycarbonylpiperidin-4-ylmethoxy)quinazoline** (200mg, 77%).

MS - ESI: 541 [MNa]⁺

- 20 ¹H NMR Spectrum: (DMSO-d₆) 1.1-1.3 (m, 2H), 1.4 (s, 9H), 1.8 (d, 2H), 1.95-2.1 (m, 1H), 2.4 (s, 1H), 2.7-2.85 (br s, 2H), 3.95 (s, 3H), 4.05 (d, 2H), 6.12 (s, 1H), 6.85 (d, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.35 (s, 1H), 7.55 (s, 1H), 8.45 (s, 1H)

Example 70

- 25 A solution of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-tert-butoxycarbonylpiperidin-4-ylmethoxy)quinazoline (155mg, 0.3mmol), (prepared as described in Example 69), in methylene chloride (5ml) containing TFA (1ml) was stirred at ambient temperature for 30 minutes. The volatiles were removed under vacuum and the residue was treated with water and adjusted to pH12 with 2M sodium hydroxide. The mixture was
30 extracted with methylene chloride. The organic layer was dried (MgSO₄), and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/ethyl acetate/methanol (5/4/1) followed by methylene

chloride/methanol (9/1) and by 3M ammonia in methanol/methanol/methylene chloride (5/15/80). After removal of the solvent by evaporation, the residue was dissolved in the minimum of methylene chloride, ether was added followed by petroleum ether. The precipitate was collected by filtration, washed with ether and dried under vacuum to give 6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline (120mg, 96%).
MS - ESI: 419 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.5-1.7 (m, 2H), 2.05 (br d, 2H), 2.3-2.4 (m, 1H), 2.4 (s, 3H), 3.05 (t, 2H), 3.4 (d, 2H), 4.09 (s, 3H), 4.25 (d, 2H), 6.95 (dd, 1H), 7.35 (s, 1H), 7.4 (d, 1H), 7.6 (s, 1H), 7.85 (s, 1H), 9.15 (s, 1H)

Example 71

Methoxyacetaldehyde (368mg, 3.47mol) (freshly distilled) followed by sodium triacetoxymethylborohydride (552mg, 2.6mol) were added to a solution of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline (726mg, 1.74mmol), (prepared as described in Example 70), in a mixture of methylene chloride (15ml) and methanol (15ml). After stirring for 1.5 hours at ambient temperature, saturated sodium hydrogen carbonate was added. The volatiles were removed under vacuum and the residue was partitioned between methylene chloride and water. The organic layer, was separated, washed with water, brine, dried (MgSO₄) and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (80/20). After removal of the solvent, the residue was triturated with ether, collected by filtration, washed with ether and dried under vacuum at 60°C to give 6-methoxy-7-(1-(2-methoxyethyl)piperidin-4-ylmethoxy)-4-(2-methylindol-5-yloxy)quinazoline (392mg, 47%).

MS - ESI 477 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.6-1.8 (m, 2H), 2.05 (br d, 2H), 2.15-2.3 (m, 1H), 2.4 (s, 3H), 3.05 (t, 2H), 3.3 (br s, 2H), 3.32 (s, 3H), 3.58 (d, 2H), 3.65 (br s, 2H), 4.05 (s, 3H), 4.18 (d, 2H), 6.2 (s, 0.5 H (partly exchanged)), 6.92 (dd, 1H), 7.32 (s, 1H), 7.35 (d, 1H), 7.55 (s, 1H), 7.8 (s, 1H), 9.15 (s, 1H)

Elemental analysis:

Found C 68.0 H 6.8 N 11.8

C₂₇H₃₂N₄O₄

Requires C 68.1 H 6.8 N 11.8%

The starting material was prepared as follows :

A solution of 1,1,2-trimethoxyethane (90g, 750mmol) in water (570ml) containing 12 N hydrochloric acid (3.75ml) was stirred at 40°C for 1.5 hours. After cooling, solid sodium chloride was added and the mixture was extracted with ether. The organic layer was dried (MgSO₄). The organic layer was distilled and the fraction from 70-90°C was collected to give
5 methoxyacetaldehyde (20.3g) which was used directly in the next step.

Example 72

Diphenylphosphoryl azide (83mg, 0.3mmol) was added in portions to a solution of 7-(2-carboxyvinyl)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (75mg, 0.2mmol),
10 triethylamine (40mg, 0.4mmol) and 1-(2-aminoethyl)pyrrolidine (46mg, 0.4mmol) in DMF (1.5ml). After stirring for 5 hours at ambient temperature, the mixture was partitioned between ethyl acetate and water. The organic layer was separated, washed with water, brine, dried (MgSO₄) and the volatiles were removed by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (9/1) followed by
15 methylene chloride/3M ammonia in methanol (9/1). After removal of the solvent, the solid was triturated with ether, collected by filtration, washed with ether and dried under vacuum to give 6-methoxy-4-(2-methylindol-5-yloxy)-7-((2-(2-pyrrolidin-1-ylethyl)carbamoyl)vinyl)quinazoline (25mg, 26%).

MS - ESI: 472 [MH]⁺

20 ¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.8-1.95 (m, 2H), 1.95-2.1 (m, 2H), 2.48 (s, 3H), 3.0-3.2 (m, 2H), 3.35 (t, 2H), 3.6 (t, 2H), 3.65 (br s, 2H), 4.11 (s, 3H), 6.18 (s, 0.5H, partially exchanged), 6.95 (dd, 1H), 7.05 (d, 1H), 7.35 (s, 1H), 7.37 (d, 1H), 7.8 (s, 1H), 7.86 (d, 1H), 8.2 (s, 1H), 8.76 (s, 1H)

25 The starting material was prepared as follows:

Trifluoromethanesulphonic anhydride (338mg, 1.2mmol) was added to a suspension of 4-(4-chloro-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline (320mg, 1mmol), (prepared as described for the starting material in Example 5), in methylene chloride (2ml) containing pyridine (2ml) cooled at 5°C. When the addition was complete, the mixture was
30 left to warm to ambient temperature and stirred for 1 hour. After removal of the volatiles by evaporation, the residue was partitioned between ethyl acetate/ether and water. The organic layer was separated, washed with 0.5M hydrochloric acid, followed by water, brine, dried

(MgSO₄) and evaporated to give 4-(4-chloro-2-fluorophenoxy)-6-methoxy-7-(trifluoromethylsulphonyloxy)quinazoline (400mg, 88%).

MS - ESI: 453 - 455 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 4.15 (s, 3H), 7.5 (d, 1H), 7.62 (t, 1H), 7.78 (d, 1H), 8.02 (s, 1H), 8.27 (s, 1H), 8.77 (s, 1H)

Triethylamine (33mg, 0.33mmol) and *tert*-butyl acrylate (77mg, 0.6mmol) followed by diphenylpropylphosphine (3.4mg, 0.008mmol) and palladium(II) acetate (1.7mg, 0.0075mmol) were added to a solution of 4-(4-chloro-2-fluorophenoxy)-6-methoxy-7-(trifluoromethylsulphonyloxy)quinazoline (136mg, 0.3mmol) in DMF (1.5ml) under argon.

When the addition was complete the reaction flask was purged with argon. The mixture was stirred at 80-85°C for 6 hours. After cooling, the mixture was partitioned between ethyl acetate and water. The aqueous layer was adjusted to pH6 with 2M hydrochloric acid. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with methylene chloride followed by methylene chloride/ether (95/5). After removal of the solvent under vacuum, the solid was triturated with pentane/ether, collected by filtration and dried under vacuum to give 4-(4-chloro-2-fluorophenoxy)-6-methoxy-7-(2-(*tert*-butoxycarbonyl)vinyl)quinazoline (63mg, 49%).

MS - ESI: 431 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.51 (s, 9H), 4.07 (s, 3H), 6.87 (d, 1H), 7.45 (d, 1H), 7.6 (t, 1H), 7.7 (s, 1H), 7.75 (d, 1H), 7.91 (d, 1H), 8.39 (s, 1H), 8.65 (s, 1H)

Elemental analysis:

Found C 61.1 H 4.8 N 6.6

C₂₂H₂₀ClFN₂O₃

Requires C 61.3 H 4.7 N 6.5%

A solution of 4-(4-chloro-2-fluorophenoxy)-6-methoxy-7-(2-(*tert*-butoxycarbonyl)vinyl)quinazoline (581mg, 1.31mmol) in a mixture of methylene chloride/TFA (2.5ml/2.5ml) was stirred at ambient temperature for 1.5 hours. After removal of the volatiles under vacuum, the residue was partitioned between ethyl acetate and water. The aqueous layer was adjusted to pH3 with 0.5M sodium hydroxide. The organic layer was separated and the aqueous layer was further extracted with ethyl acetate. The combined organic layers were washed with brine, dried (MgSO₄) and evaporated to give 7-(2-carboxyvinyl)-4-(4-chloro-2-fluorophenoxy)-6-methoxyquinazoline (430mg, 85%).

¹H NMR Spectrum: (DMSO-d₆) 4.08 (s, 3H), 6.9 (d, 1H), 7.45 (s, 1H), 7.6 (t, 1H), 7.70 (s,

1H), 7.73 (d, 1H), 7.95 (d, 1H), 8.39 (s, 1H), 8.66 (s, 1H)

1M Sodium HMDS in THF (0.84ml, 8.4mmol) was added to a suspension of 7-(2-carboxyvinyl)-4-(4-chloro-2-fluorophenoxy)-6-methoxyquinazoline (105mg, 0.28mmol) and 5-hydroxy-2methylindole (82mg, 0.56mmol), (prepared as described for the starting material in Example 48), in DMSO (1.5ml). After stirring for 2 hours at ambient temperature, the mixture was partitioned between ethyl acetate and water. The aqueous layer was adjusted to pH3 with 2M hydrochloric acid. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with methylene chloride/methanol (95/5 followed by 90/10 and 70/30) to give 7-(2-carboxyvinyl)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (75mg, 71%).

¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H), 4.06 (s, 3H), 6.15 (s, 1H), 6.82 (d, 1H), 6.9 (dd, 1H), 7.3 (s, 1H), 7.35 (d, 1H), 7.68 (s, 1H), 7.84 (d, 1H), 8.25 (s, 1H), 8.55 (s, 1H)

Example 73

A suspension of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (321mg, 1mmol), (prepared as described in Example 49), 1-bromo-3-chloropropane (120μl, 1.2mmol) and potassium carbonate (359mg, 2.6mmol) in DMF (5ml) was stirred at ambient temperature overnight. After addition of water, the precipitate was collected by filtration, washed with water and dried over phosphorus pentoxide at 60°C to give 7-(3-chloropropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (280mg, 70%).

MS - ESI: 398 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.2-2.35 (m, 2H), 2.4 (s, 3H), 3.85 (t, 2H), 4.0 (s, 3H), 4.32 (t, 2H), 6.15 (s, 1H), 6.88 (d, 1H), 7.27 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

Example 74

A solution of 7-(3-chloropropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (150mg, 0.38mmol), (prepared as described in Example 73), in 1-methylpiperazine (2ml) was heated at 100°C for 2 hours. After cooling, the mixture was partitioned between ethyl acetate and aqueous 5% sodium hydrogen carbonate. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography on an isolate column eluting with methanol/ethyl acetate/methylene chloride (1/4/5 followed by 1/9/0) and 3M ammonia in methanol/methanol/methylene chloride

(5/10/80). After removal of the solvent under vacuum, the solid was dissolved in the minimum of methylene chloride and ether/petroleum ether was added. The precipitate was collected by filtration, and dried under vacuum to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline** (55mg, 32%).

5 MS - ESI: 462 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD, 60°C) 2.2-2.3 (m, 2H), 2.4 (s, 3H), 2.9 (s, 3H), 3.4-3.5 (m, 4H), 3.5-3.8 (m, 6H), 4.07 (s, 3H), 4.4 (t, 2H), 6.95 (d, 1H), 7.35 (s, 1H), 7.4 (d, 1H), 7.55 (s, 1H), 7.8 (s, 1H), 8.95 (s, 1H)

10 **Example 75**

Triphenylphosphine (262mg, 1mmol) and *N,N*-diethylethanolamine (88mg, 0.75mmol) were added to a suspension of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (160mg, 0.5mmol), (prepared as described in Example 49), in methylene chloride (5ml), followed by the addition, in portions, of diethyl azodicarboxylate (165μl, 15 1mmol). After stirring for 1 hour at ambient temperature, the volatiles were removed under vacuum. The residue was purified by column chromatography eluting with methylene chloride/methanol (95/5) followed by methylene chloride/3M ammonia in methanol (90/10) to give **7-(2-(*N,N*-diethylamino)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (147mg, 70%).

20 MS - ESI 421 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.0 (t, 6H), 2.41 (s, 3H), 2.6 (q, 4H), 2.88 (t, 2H), 3.97 (s, 3H), 4.24 (t, 2H), 6.14 (s, 1H), 6.89 (dd, 1H), 7.25 (s, 1H), 7.32 (d, 1H), 7.38 (s, 1H), 7.58 (s, 1H), 8.48 (s, 1H)

Elemental analysis:	Found	C 66.2	H 6.9	N 13.1
---------------------	-------	--------	-------	--------

25 C ₂₄ H ₂₈ N ₄ O ₃ 0.8H ₂ O	Requires	C 66.3	H 6.9	N 12.9%
--	----------	--------	-------	---------

Example 76

Using an analogous procedure to that described in Example 75, 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (321mg, 1mmol), (prepared as described in Example 30 49), was reacted with 2-((1-*tert*butoxycarbonyl)piperidin-4-yloxy)ethanol (294mg, 1.2mmol) to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-((1-*tert*butoxycarbonyl)piperidin-4-yloxy)ethoxy)quinazoline** (420mg, 76%).

MS - ESI: 549 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.4 (s, 9H), 1.3-1.5 (m, 2H), 1.7-1.9 (m, 2H), 2.38 (s, 3H), 3.0 (br t, 2H), 3.5-3.7 (m, 3H), 3.85 (m, 2H), 3.98 (s, 3H), 4.3 (t, 2H), 6.12 (s, 1H), 6.85 (d, 1H), 7.22 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.55 (s, 1H), 8.48 (s, 1H)

5

The starting material was prepared as follows:

tert-Butoxycarbonyl anhydride (1.52g, 7mmol) in acetone (3.5ml) was added to a solution of 4,4-(ethylenedioxy)piperidine (1g, 7mmol) in acetone/trichloromethane (3.5ml/3.5ml) cooled at 0°C. After stirring for 4 hours at ambient temperature, the volatiles were removed under vacuum. The residue was dissolved in ether and the ether solution was washed with water, brine, dried (MgSO₄) and evaporated to give 4,4-(ethylenedioxy)-1-*tert*butoxycarbonylpiperidine (1.7g, quant.).

10

¹H NMR Spectrum: (CDCl₃): 1.46 (s, 9H), 1.65 (t, 4H), 3.5 (t, 4H), 3.97 (s, 4H)

Freshly distilled boron trifluoride etherate (52μl, 0.41mmol), followed by sodium cyanoborohydride (38mg, 0.6mmol) were added to a solution of 4,4-(ethylenedioxy)-1-*tert*butoxycarbonylpiperidine (100mg, 0.41mmol) in THF (1.4ml) cooled at 0°C. After stirring for 6 hours at ambient temperature, boron trifluoride etherate (52μl) and sodium cyanoborohydride (26mg, 0.41mmol) were added. After stirring overnight at ambient temperature, the mixture was partitioned between ethyl acetate and 2M sodium hydroxide. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with methylene chloride/methanol (95/5) followed by methylene chloride/methanol/3M ammonia in methanol (80/15/5) to give 2-((1-*tert*butoxycarbonyl)piperidin-4-yloxy)ethanol (42mg, 42%).

15

20

MS - ESI: 268 [MNa]⁺

¹H NMR Spectrum: (CDCl₃) 1.48 (s, 9H), 1.5-1.6 (m, 2H), 1.8-1.9 (m, 2H), 2.0 (t, 1H), 3.05-3.15 (m, 2H), 3.5 (m, 1H), 3.57 (t, 2H), 3.7-3.9 (m, 4H)

25

Example 77

A solution of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-((1-*tert*butoxycarbonyl)piperidin-4-yloxy)ethoxy)quinazoline (379 mg, 0.69 mmol), (prepared as described in Example 76), in methylene chloride (7ml) containing TFA (2.5ml) was stirred for 1.5 hours at ambient temperature. After removal of the volatiles under vacuum, the residue

30

was partitioned between ethyl acetate and water. Solid sodium hydrogen carbonate and 2N sodium hydroxide were added to adjust the aqueous layer to about pH10. The organic layer was washed with water, followed by brine, dried (MgSO₄) and evaporated. The residue was triturated with ether, filtered, washed with ether and dried under vacuum to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(piperidin-4-yloxy)ethoxy)quinazoline** (164 mg, 53 %).

¹HNMR Spectrum: (DMSO-d₆) 1.2-1.4 (m, 2H), 1.8-1.9 (m, 2H), 2.47 (s, 3H), 2.4-2.5 (m, 2H), 2.9-3.0 (d, 2H), 3.3-3.5 (m, 1H), 3.95 (s, 2H), 4.0 (s, 3H), 4.35 (s, 2H), 6.15 (s, 1H), 6.9 (dd, 1H), 7.28 (s, 1H), 7.32 (d, 1H), 7.41 (s, 1H), 7.60 (s, 1H), 8.49 (s, 1H)

MS-ESI : 448 [M.]⁺

Example 78

A solution of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (193 mg, 0.6 mmol), (prepared as described in Example 49), 4-(2-hydroxyethoxy)pyridine (166 mg, 1.2 mmol), (J. Chem. Soc. Perkin II, 1987, 1867), in methylene chloride (5 ml) containing triphenylphosphine (330 mg, 1.26 mmol) and diisopropyl azodicarboxylate (255 mg, 1.26 mmol) was stirred at ambient temperature for 2 hours. The precipitate was filtered, triturated with ether followed by ethyl acetate, and dried under vacuum to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(4-pyridyloxy)ethoxy)quinazoline** (142 mg, 54 %).

¹HNMR Spectrum: (DMSO-d₆) 2.40 (s, 3H), 3.97 (s, 3H), 4.52 (t, 2H), 4.58 (t, 2H), 6.14 (s, 1H), 6.89 (dd, 1H), 7.07 (d, 2H), 7.26 (s, 1H), 7.31 (d, 1H), 7.46 (s, 1H), 7.61 (s, 1H), 8.41 (d, 2H), 8.5 (s, 1H)

MS-ESI : 443 [MH]⁺

Elemental analysis	Found	C 66.6	H 5.0	N 12.5
C ₂₅ H ₂₂ N ₄ O ₄ 0.12 CH ₂ Cl ₂	Requires	C 66.9	H 5.0	N 12.4%

Example 79

A suspension of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-tert-butoxycarbonylamino)ethoxy)quinazoline (148 mg, 0.31 mmol), (prepared as described in Example 149), in methylene chloride (4 ml) containing TFA (1 ml) was stirred for 1 hour. After removing the volatiles under vacuum, the residue was azeotroped with toluene. The residue was dissolved in methylene chloride (3 ml) and triethylamine (215 µl, 1.5 mmol) was added followed by methanesulphonyl chloride (48 µl, 0.62 mmol). After stirring for 1 hour at

ambient temperature, the mixture was partitioned between methylene chloride and water. The organic layer was separated, washed with water, brine, dried (MgSO_4) and evaporated. The residue was purified by column chromatography eluting with ethylacetate/methanol (99/1 followed by 97/3). After evaporation of the solvent, the solid was triturated with ether,

5 filtered, washed with ether and dried under vacuum to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-methylsulphonylamino)ethoxy)quinazoline** (54 mg, 38 %).

^1H NMR Spectrum: (DMSO-d_6) 2.4 (s, 3H), 2.93 (s, 3H), 3.0 (s, 3H), 3.62 (t, 2H), 4.0 (s, 3H), 4.38 (t, 2H), 6.14 (s, 1H), 6.88 (dd, 1H), 7.26 (s, 1H), 7.3 (d, 1H), 7.43 (s, 1H), 7.61 (s, 1H), 8.49 (s, 1H)

10 MS-ESI : 457 $[\text{MH}]^+$

Example 80

A solution of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(piperidin-4-yloxy)ethoxy)quinazoline (76 mg, 0.17 mmol), (prepared as described in Example 77), in acrylonitrile (0.5 ml), methylene chloride (1 ml) and methanol (1 ml) was stirred overnight at ambient temperature. After removal of the volatiles under vacuum the residue was purified by column chromatography eluting with methylene chloride/methanol (98/2 followed by 95/5 and 90/10). The residue was triturated with ethyl acetate and ether. The resulting solid was filtered and dried under vacuum to give **7-(2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (73 mg, 86 %).

^1H NMR Spectrum: (DMSO-d_6) 1.4-1.55 (m, 2H), 1.8-1.9 (m, 2H), 2.15 (t, 2H), 2.4 (s, 3H), 2.55 (t, 2H), 2.65 (t, 2H), 2.7-2.8 (m, 2H), 3.4-3.5 (m, 1H), 3.85 (m, 2H), 4.0 (s, 3H), 4.3 (t, 2H), 6.15 (s, 1H), 6.9 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

MS-ESI : 502 $[\text{MH}]^+$

25	Elemental analysis	Found	C 67.0	H 6.2	N 14.0
	$\text{C}_{28}\text{H}_{31}\text{N}_5\text{O}_4$	Requires	C 67.1	H 6.2	N 14.0%

Example 81

A solution of 4-chloro-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (100 mg, 0.31 mmol), (prepared as described for the starting material in Example 9), 6-hydroxyindole (50 mg, 0.37 mmol) and potassium carbonate (64 mg, 0.466 mmol) in DMF (1 ml) was heated at 95°C for 4 hours. After cooling, the mixture was diluted with methylene chloride and

poured onto a silica column. The product was eluted with methylene chloride, followed by methylene chloride/methanol (80/20 followed by 70/30 and 50/50). After removal of the solvent by evaporation, the precipitate was triturated with ether, filtered and dried under vacuum to give 6-methoxy-4-(indol-6-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (90 mg, 69 %).

¹H NMR Spectrum: (DMSO-d₆) 1.85 (br s, 4H), 2.15-2.25 (m, 2H), 2.85-3.15 (m, 6H), 4.01 (s, 3H), 4.32 (t, 2H), 6.5 (s, 1H), 6.95 (dd, 1H), 7.32 (s, 1H), 7.4 (s, 2H), 7.6 (d, 1H), 7.65 (s, 1H), 8.52 (s, 1H)

MS-ESI : 419 [MH]⁺

Example 82

Diisopropyl azodicarboxylate (146 mg, 0.72 mmol) was added to a solution of 7-hydroxy-4-(2-methylindol-5-yloxy)quinazoline (100 mg, 0.34 mmol), triphenyl phosphine (189 mg, 0.72 mol), and 3-pyrrolidinopropan-1-ol (89 mg, 0.686 mmol), (J. Org. Chem. 1988, 53, 3164), in methylene chloride (2.5 ml). After stirring overnight at ambient temperature, the solid was filtered. The filtrate was purified by column chromatography eluting with ethyl acetate/methylene chloride (1/1) followed by ethyl acetate/methylene chloride/methanol (4/5/1), methylene chloride/methanol (9/1) and 3N ammonia in methanol/methylene chloride (1/9). After removal of the solvent, the residue was triturated with ether, filtered, and dried under vacuum to give 4-(2-methylindol-5-yloxy)-7-(3-(pyrrolidin-yl)propoxy)quinazoline (49 mg, 35 %).

¹H NMR Spectrum: (DMSO-d₆) 1.8-2.0 (m, 2H), 2.0-2.15 (m, 2H), 2.2-2.32 (m, 2H), 2.41 (s, 3H), 3.0-3.2 (m, 2H), 3.4 (t, 2H), 3.6-3.7 (m, 2H), 4.35 (t, 2H), 6.2 (s, 1H), 6.95 (dd, 1H), 7.3 (s, 1H), 7.35 (d, 1H), 7.5 (s, 1H), 7.57 (dd, 1H), 8.5 (d, 1H), 9.15 (s, 1H)

MS-ESI : 403 [MH]⁺

The starting material was prepared as follows:

Sodium (368mg, 16mmol) was added to benzyl alcohol (10ml, 96mmol) and the mixture was heated at 148°C for 30 minutes. 7-Fluoro-3,4-dihydroquinazolin-4-one (656mg, 4mmol), (J. Chem. Soc. section B 1967, 449), was added and the mixture maintained at 148°C for 24 hours. The reaction mixture was allowed to cool, the solution was poured on to water (170ml) and the aqueous mixture adjusted to pH3 with concentrated hydrochloric acid. The

precipitate was collected by filtration, washed with water, ether and dried under vacuum to give 7-benzyloxy-3,4-dihydroquinazolin-4-one (890mg, 89%) as a white solid.

m.p. 267-269°C

¹H NMR Spectrum: (DMSO-d₆; CF₃COOD) 5.32(s, 2H); 7.25(d, 1H); 7.32-7.52(m, 6H);
5 8.12(d, 1H); 8.99(s, 1H)

MS - ESI: 252 [MH]⁺

Elemental analysis:	Found	C 71.4	H 4.9	N 10.7
C ₁₅ H ₁₂ N ₂ O ₂ 0.04H ₂ O	Requires	C 71.2	H 4.8	N 11.1%

A mixture of 7-benzyloxy-3,4-dihydroquinazolin-4-one (11g, 43.6mmol) and DMF
10 (1ml) in thionyl chloride (100ml) was heated at reflux for 1.5 hours. Excess thionyl chloride was removed by evaporation and the residue azeotroped with toluene. The residue was partitioned between methylene chloride and water and saturated aqueous sodium hydrogen carbonate was added until the aqueous layer was at about pH9. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated to give 7-benzyloxy-4-
15 chloroquinazoline (10.5g, 89%).

¹H NMR Spectrum: (DMSO-d₆) 5.4 (s, 2H); 7.35-7.65 (m, 6H); 8.2 (d, 1H); 9.0 (s, 1H)

MS - ESI: 270 [MH]⁺

A solution of 7-benzyloxy-4-chloroquinazoline (2g, 7.4mmol), 5-hydroxy-2-methylindole (1.3 g, 8.9 mmol), (prepared as described for the starting material in Example
20 48), in DMF (20 ml) containing potassium carbonate (1.53 g, 11.1 mmol) was stirred at 80°C for 3 hours. After cooling, the mixture was poured in portions into ice/water. The precipitate was filtered and washed with water and dried under vacuum. The solid was dissolved in methylene chloride and was purified by column chromatography eluting with ethyl acetate and methylene chloride (1/1) to give 7-benzyloxy-4-(2-methylindol-5-yloxy)quinazoline (2.28
25 g, 81 %).

MS-ESI : 382 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.41 (s, 3H), 5.4 (s, 2H), 6.15 (s, 1H), 6.9 (dd, 1H), 7.3 (s, 1H), 7.35 (d, 1H), 7.4 (d, 1H), 7.4-7.5 (m, 4H), 7.55 (d, 2H), 8.32 (d, 1H), 8.6 (s, 1H).

10% Palladium on charcoal (200 mg) followed by ammonium formate (4.34 g, 69
30 mmol) were added to a solution of 7-benzyloxy-4-(2-methylindol-5-yloxy)quinazoline (1.75 g, 4.58 mmol) in DMF (60 ml). After stirring for 1 hour at ambient temperature, the mixture


was filtered. The filtrate was evaporated. The residue was triturated with water, filtered, washed with ethyl acetate, and dried under vacuum to give 7-hydroxy-4-(2-methylindol-5-yloxy)quinazoline (1.24 g, 93 %).

¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H), 6.14 (s, 1H), 6.88 (dd, 1H), 7.17 (s, 1H), 7.25-7.3 (m, 2H), 7.30 (d, 1H), 8.24 (d, 1H), 8.5 (s, 1H)



Examples 83-89

Using an analogous procedure to that described in Example 82, the appropriate alcohols were reacted with 7-hydroxy-4-(2-methylindol-5-yloxy)quinazoline, (prepared as described for the starting material in Example 82), to give the compounds described in Table VI below.

Table VI



Example number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	R	Note
83	34	24	412		a
84	45	32	405		b
85	5	3	417		c
86	56	35	467		d
87	63	44	419		e

88	24	17	403		f
89	84	63	387		g

a) 7-Hydroxy-4-(2-methylindol-5-yloxy)quinazoline (100 mg) was reacted with 3-(methylsulphonyl)-1-propanol (95 mg), (prepared as described for the starting material in Example 50), to give 7-(3-(methylsulphonyl)propoxy)-4-(2-methylindol-5-

5 **yloxy)quinazoline.**

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.2-2.3 (m, 2H), 2.4 (s, 3H), 3.05 (s, 3H), 3.3-3.45 (m, 2H), 4.4 (t, 2H), 6.2 (s, 1H), 6.95 (dd, 1H), 7.38 (s, 1H), 7.4 (d, 1H), 7.5 (s, 1H), 7.6 (dd, 1H), 8.5 (d, 1H), 9.2 (s, 1H)

Elemental analysis

Found C 60.2 H 5.3 N 10.6

10 C₂₁H₂₁N₃O₄S 0.4 DMF

Requires C 60.5 H 5.4 N 10.8%

b) 7-Hydroxy-4-(2-methylindol-5-yloxy)quinazoline (100 mg) was reacted with 4-(2-hydroxyethyl)morpholine (90 mg) to give 4-(2-methylindol-5-yloxy)-7-(2-morpholinoethoxy)quinazoline.

15 ¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.4 (s, 3H), 3.1-3.3 (m, 2H), 3.62 (d, 2H), 3.7-3.9 (m, 4H), 4.05 (d, 2H), 4.7 (t, 2H), 6.2 (s, 0.5 H, partially exchanged), 6.95 (dd, 1H), 7.35 (s, 1H), 7.39 (d, 1H), 7.6 (s, 1H), 7.65 (dd, 1H), 8.55 (d, 1H), 9.15 (s, 1H)

Elemental analysis

Found C 67.2 H 6.0 N 13.5

C₂₃H₂₄N₄O₃ 0.3 H₂O

Requires C 67.4 H 6.1 N 13.7%

20

c) 7-Hydroxy-4-(2-methylindol-5-yloxy)quinazoline (100 mg) was reacted with 1-(3-hydroxypropyl)piperidine (98 mg) to give 4-(2-methylindol-5-yloxy)-7-(3-(piperidin-1-yl)propoxy)quinazoline.

25 ¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.2-1.5 (m, 2H), 1.6-1.8 (m, 2H), 1.8-1.9 (m, 2H), 2.25-2.35 (m, 2H), 2.45 (s, 3H), 2.95 (t, 2H), 3.25-3.3 (m, 2H), 3.55 (d, 2H), 4.4 (t, 2H), 6.95 (dd, 1H), 7.4 (s, 1H), 7.45 (d, 1H), 7.5 (s, 1H), 7.6 (d, 1H), 8.55 (d, 1H), 9.15 (s, 1H)

d) 7-Hydroxy-4-(2-methylindol-5-yloxy)quinazoline (100 mg) was reacted with 3-(1,1-dioxothiomorpholino)-1-propanol (133 mg), (prepared as described for the starting material in Example 5), to give **4-(2-methylindol-5-yloxy)-7-(3-(1,1-dioxothiomorpholino)propoxy)quinazoline**.

5 ¹H NMR Spectrum: (DMSO-d₆) 1.9-2.0 (m, 2H), 2.4 (s, 3H), 1.6-1.7 (m, 2H), 2.9 (br s, 4H), 3.1 (br s, 4H), 4.25 (t, 2H), 6.12 (s, 1H), 6.85 (d, 1H), 7.22 (s, 1H), 7.3 (d, 1H), 7.3-7.4 (m, 2H), 8.25 (d, 1H), 8.55 (s, 1H)

10 e) 7-Hydroxy-4-(2-methylindol-5-yloxy)quinazoline (100 mg) was reacted with 4-(3-hydroxypropyl)morpholine (100 mg), (prepared as described for the starting material in Example 60), to give **4-(2-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.95-2.05 (m, 2H), 2.42 (s, 3H), 2.5 (t, 2H), 2.55 (t, 4H), 3.6 (t, 4H), 4.3 (t, 2H), 6.18 (s, 1H), 6.9 (dd, 1H), 7.3 (s, 1H), 7.35 (d, 1H), 7.3-7.4 (m, 2H), 8.3 (d, 1H), 8.6 (s, 1H)

15	Elemental analysis	Found	C 66.5	H 6.2	N 12.7
	C ₂₄ H ₂₆ N ₄ O ₃ 0.14 CH ₂ Cl ₂ 0.7 H ₂ O	Requires	C 66.7	H 6.4	N 13.0%

f) 7-Hydroxy-4-(2-methylindol-5-yloxy)quinazoline (100 mg) was reacted with 1-(2-hydroxyethyl)piperidine (89 mg) to give **4-(2-methylindol-5-yloxy)-7-(2-(piperidin-1-yl)ethoxy)quinazoline**.

20 ¹H NMR Spectrum: (DMSO-d₆) 1.4-1.5 (br s, 2H), 1.5-1.7 (br s, 4H), 2.42 (s, 3H), 2.5-2.7 (br s, 4H), 2.8-3.0 (br s, 2H), 4.35 (br s, 2H), 6.18 (s, 1H), 6.9 (dd, 1H), 7.3 (s, 1H), 7.35 (d, 1H), 7.4 (d, 1H), 7.42 (s, 1H), 8.3 (d, 1H), 8.6 (s, 1H)

	Elemental analysis	Found	C 69.0	H 6.6	N 13.4
25	C ₂₄ H ₂₆ N ₄ O ₂ 0.8 H ₂ O	Requires	C 69.1	H 6.7	N 13.4%

g) 7-Hydroxy-4-(2-methylindol-5-yloxy)quinazoline (100 mg) was reacted with 2-(1*H*-1,2,4-triazol-1-yl)ethanol (78 mg), (Ann. Phar. Fr. 1977, 35, 503-508), to give **4-(2-methylindol-5-yloxy)-7-(2-(1*H*-1,2,4-triazol-1-yl)ethoxy)quinazoline**.

30 ¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H), 4.6 (m, 2H), 4.7 (m, 2H), 6.15 (s, 1H), 6.9 (dd, 1H), 7.28 (s, 1H), 7.3 (d, 2H), 7.4 (s, 1H), 8.02 (s, 1H), 8.3 (d, 1H), 8.6 (s, 1H), 8.65 (s, 1H)

	Elemental analysis	Found	C 63.7	H 4.8	N 21.5
--	--------------------	-------	--------	-------	--------

$C_{21}H_{18}N_6O_2 \cdot 0.5 H_2O$

Requires C 63.8 H 4.8 N 21.3%

Example 90

A solution of 7-hydroxy-4-(2-methylindol-5-yloxy)quinazoline (423 mg, 1.45 mol),
5 (prepared as described for the starting material in Example 82), triphenylphosphine (685 mg, 2.61 mmol), 4-hydroxymethyl-1-*tert*-butoxycarbonylpiperidine (500 mg, 2.32 mmol),
(prepared as described for the starting material in Example 10), and diisopropyl
azodicarboxylate (528 mg, 2.61 mmol) in methylene chloride (18 ml) was stirred overnight at
ambient temperature. The mixture was then poured onto a column of silica and eluted with
10 ethyl acetate. After evaporation of the solvent, the residue was triturated with ether, filtered,
and dried under vacuum to give **7-(1-*tert*-butoxycarbonylpiperidin-4-ylmethoxy)-4-(2-
methylindol-5-yloxy)quinazoline** (478 mg, 68 %).

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.4 (m, 2H), 1.42 (s, 9H), 1.85 (d, 2H), 2.0-2.1 (m, 1H),
2.42 (s, 3H), 2.7-2.9 (br s, 2H), 3.95-4.05 (m, 2H), 4.1 (d, 2H), 6.15 (s, 1H), 6.9 (dd, 1H), 7.3
15 (s, 1H), 7.33 (d, 1H), 7.38 (s, 1H), 7.35-7.4 (m, 1H), 8.3 (d, 1H), 8.6 (s, 1H)

MS-ESI : 489 [MH]⁺

Elemental analysis

Found C 68.7 H 6.7 N 11.3

 $C_{28}H_{32}N_4O_4$

Requires C 68.8 H 6.6 N 11.5%

Example 91

To a suspension of 4-(2,3-dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline
(124 mg, 0.32 mmol) in methylene chloride (2.5 ml) was added triphenylphosphine (179 mg,
0.628 mmol), 1-(2-hydroxyethyl)pyrrolidine (75 mg, 0.65 mmol) followed by diisopropyl
azodicarboxylate (134 µl, 0.68 mmol) in portions. After stirring overnight at ambient
25 temperature the mixture was poured onto a column of silica and eluted with ethyl
acetate/methylene chloride (1/1) followed by ethyl acetate/methylene chloride/methanol
(4/5/1) followed by methylene chloride/methanol (9/1). After removal of the solvent, the
solid was triturated with ether, filtered, washed with ether and dried under vacuum to give **4-
(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline** (51 mg,
30 37 %).

¹H NMR Spectrum: (DMSO-d₆) 1.6-1.75 (m, 4H), 2.12 (s, 3H), 2.28 (s, 3H), 2.52 (br s, 4H),
3.85 (t, 2H), 3.93 (s, 3H), 4.25 (t, 2H), 6.8 (d, 1H), 7.17 (s, 1H), 7.22 (d, 1H), 7.33 (s, 1H),

7.54 (s, 1H), 8.43 (s, 1H)

The starting material was prepared as follows:

To a solution of 2,3-dimethyl-5-methoxyindole (175mg, 1 mmol), (J. Chem. Soc. 1957, 3175-3180) in methylene (5 ml) cooled at -60°C was added boron tribromide (210 µl, 2.2 mmol) dropwise. After completion of addition, the mixture was left to warm up to ambient temperature and was stirred for 1 hour. Water was added and the pH was adjusted to 6 with 2N sodium hydroxide. The mixture was extracted with ethyl acetate and the organic layer was separated, washed with brine, dried (MgSO₄) and evaporated to give 2,3-dimethyl-5-hydroxyindole (124mg, 77%).

¹H NMR Spectrum: (DMSO-d₆) 2.1 (s, 3H) ; 2.3 (s, 3H) ; 6.5 (dd, 1H) ; 6.65 (d, 1H) ; 7.0 (d, 1H) ; 8.45 (s, 1H)

Under nitrogen, to a solution of 2,3-dimethyl-5-hydroxyindole (643mg, 4 mmol), in DMF (10 ml) was added potassium carbonate (690mg, 5 mmol). After stirring for 15 minutes at ambient temperature, 7-benzyloxy-4-chloro-6-methoxyquinazoline (1g, 3.33 mmol), (prepared as described for the starting material in Example 1), was added. The mixture was heated at 90°C for 2 hours followed by 30 minutes at 95°C. After cooling, the mixture was poured onto water (100ml) cooled at 5 °C. The precipitate was filtered, washed with water, followed by ether and dried under vacuum to give 7-benzyloxy-4-(2,3-dimethylindol-5-yloxy)-6-methoxyquinazoline (1.4 g, 95%).

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H) ; 2.35 (s, 3H) ; 4.02 (s, 3H) ; 5.4 (s, 2H) ; 6.9 (dd, 1H) ; 7.22 (d, 1H) ; 7.3 (d, 1H) ; 7.35-7.6 (m, 6H) ; 7.65 (s, 1H) ; 8.5 (s, 1H)

A solution of 7-benzyloxy-4-(2,3-dimethylindol-5-yloxy)-6-methoxyquinazoline (2g, 4.7 mmol) in DMF (120 ml) containing ammonium formate (11gr, 174 mmol) and 10% palladium on charcoal (200mg) was stirred for 2.5 hours at ambient temperature. The mixture was filtered, and the filtrate was evaporated under vacuum. The residue was triturated with ether and the solid was filtered, washed with water followed by ether and dried under vacuum at 50°C to give 4-(2,3-dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (1.1 g, 69%).


¹H NMR Spectrum: (DMSO-d₆) 2.1 (s, 3H) ; 2.32 (s, 3H) ; 3.97 (s, 3H) ; 7.85 (dd, 1H) ; 7.2 (bs, 2H) ; 7.25 (d, 1H) ; 7.58 (s, 1H) ; 8.4 (s, 1H)

Examples 92-106










Using an analogous procedure to that described in Example 91, the appropriate alcohol was reacted with 4-(2,3-dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline, (prepared as described for the starting material in Example 91), to give the compounds described in the Table VII below.

5 **Table VII**



Example number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	R	HPLC* RT (min)	Note
92	91	65	431		-	a
93	78	55	438		-	b
94	34	27	435		-	c
95	39	33	407		-	d
96	58	44	449		-	e
97	58	47	421		-	f
98	85	66	447		-	g
99	24	18	447		-	h

100	110	82	461		-	i
101	9	7	447		-	j
102	81	62	463		3.4	k
103	75	57	451		-	l
104	96	65	511		-	m
105	103	78	457		-	n
106	64	49	456		-	o

* HPLC conditions 2) as described hereinbefore.

a) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (124 mg) was reacted with 2-(1*H*-1,2,4-triazol-1-yl)ethanol (74 mg), (Ann. Phar. Fr. 1977, 35, 503-508), to give 4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(2-(1*H*-1,2,4-triazol-1-yl)ethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.10 (s, 3H), 2.30 (s, 3H), 3.93 (s, 3H), 4.52 (m, 2H), 4.55-4.65 (m, 2H), 6.85 (d, 1H), 7.2 (s, 1H), 7.25 (d, 1H), 7.4 (d, 1H), 7.58 (s, 1H), 8.0 (s, 1H), 8.48 (s, 1H), 8.58 (s, 1H)

b) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (124 mg) was reacted with 2-(2-methoxyethoxy)ethanol (78 mg) to give 4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.14 (s, 3H), 2.35 (s, 3H), 3.3 (s, 3H), 3.5 (t, 2H), 3.65 (t, 2H), 3.85 (t, 2H), 4.0 (s, 3H), 4.32 (t, 2H), 6.9 (d, 1H), 7.25 (d, 1H), 7.28 (d, 1H), 7.4 (s, 1H), 7.6

(s, 1H), 8.5 (s, 1H)

c) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with N,N-diethylethanolamine (68 mg) to give 7-(2-(N,N-diethylamino)ethoxy)-4-(2,3-

5 **dimethylindol-5-yloxy)-6-methoxyquinazoline.**

¹H NMR Spectrum: (DMSO-d₆) 1.05 (t, 6H), 2.15 (s, 3H), 2.35 (s, 3H), 2.6-2.7 (m, 4H), 2.92 (br s, 2H), 4.0 (s, 3H), 4.25 (t, 2H), 6.9 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

10 d) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with N,N-dimethylethanolamine (52 mg) to give 7-(2-(N,N-dimethylamino)ethoxy)-4-(2,3-

dimethylindol-5-yloxy)-6-methoxyquinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.35 (s, 9H), 2.85 (br s, 2H), 4.0 (s, 3H), 4.35 (t, 2H), 6.87 (dd, 1H), 7.22 (s, 1H), 7.3 (d, 1H), 7.42 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

15

e) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 4-(2-hydroxyethyl)morpholine (59 mg) to give 4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(2-morpholinoethoxy)quinazoline.

20 ¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.35 (s, 3H), 3.25-3.4 (m, 2H), 3.65 (d, 2H), 3.7-3.8 (m, 4H), 4.0-4.1 (m, 2H), 4.1 (s, 3H), 4.7 (t, 2H), 6.95 (dd, 1H), 7.3 (s, 1H), 7.35 (d, 1H), 7.6 (s, 1H), 7.8 (s, 1H), 9.0 (s, 1H)

25 f) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 3-(N,N-dimethylamino)propan-1-ol (60 mg) to give 7-(3-(N,N-dimethylamino)propoxy)-4-(2,3-dimethylindol-5-yloxy)-6-methoxyquinazoline.

¹H NMR Spectrum: (DMSO-d₆) 1.95-2.05 (m, 2H), 2.15 (s, 3H), 2.2 (s, 6H), 2.35 (s, 3H), 2.45 (t, 2H), 4.0 (s, 3H), 4.25 (t, 2H), 6.9 (dd, 1H), 7.22 (d, 1H), 7.3 (d, 1H), 7.37 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

30 g) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 1-(2-hydroxyethyl)-2-pyrrolidinone (75 mg) to give 4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(2-(2-oxopyrrolidin-1-yl)ethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 1.9-2.05 (m, 4H), 2.15 (s, 3H), 2.25 (t, 2H), 2.35 (s, 3H), 3.65 (t, 2H), 4.0 (s, 3H), 4.35 (t, 2H), 6.9 (d, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.45 (s, 1H), 7.62 (s, 1H), 8.5 (s, 1H)

- 5 h) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 2-(2-hydroxyethyl)piperidine (75 mg) to give **4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(2-(piperidin-2-yl)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.0-1.15 (m, 1H), 1.25-1.4 (m, 2H), 1.5 (br s, 1H), 1.65 (d, 1H), 1.7-1.8 (m, 1H), 1.8-1.9 (m, 2H), 2.15 (s, 3H), 2.35 (s, 3H), 2.5 (d, 1H), 2.6-2.7 (m, 1H),
10 2.9-3.0 (m, 1H), 4.0 (s, 3H), 4.2-4.35 (m, 2H), 6.88 (dd, 1H), 7.2 (s, 1H), 7.27 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

- i) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 1-(2-hydroxyethyl)pyrrolidin-2,5-dione (83 mg) to give **4-(2,3-dimethylindol-5-yloxy)-7-(2-(2,5-dioxopyrrolidin-1-yl)ethoxy)-6-methoxyquinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.12 (s, 3H), 2.35 (s, 3H), 2.68 (s, 4H), 3.85 (t, 2H), 3.95 (s, 3H), 4.35 (t, 2H), 6.88 (dd, 1H), 7.22 (s, 1H), 7.25 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

- 20 j) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 1-methyl-3-piperidinemethanol (75 mg) to give **4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-3-ylmethoxy)quinazoline**.

- 25 k) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 4-(3-hydroxypropyl)morpholine (75 mg), (prepared as described for the starting material in Example 60), to give **4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.95-2.05 (m, 2H), 2.15 (s, 3H), 2.35 (s, 3H), 2.42 (br s, 4H), 2.5 (t, 2H), 3.6 (m, 4H), 4.0 (s, 3H), 4.25 (t, 2H), 6.85 (dd, 1H), 7.25 (d, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H).

30

- l) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 2-(N-(2-methoxyethyl)-N-methylamino)ethanol (77 mg), (prepared as described for the

starting material in Example 59), to give **4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(2-(N-(2-methoxyethyl)-N-methylamino)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.35 (s, 6H), 2.65 (t, 2H), 2.9 (t, 2H), 3.25 (s, 3H), 3.45 (t, 2H), 4.0 (s, 3H), 4.3 (t, 2H), 6.9 (dd, 1H), 7.22 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

m) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 3-(1,1-dioxothiomorpholino)-1-propanol (112 mg), (prepared as described for the starting material in Example 5), to give **4-(2,3-dimethylindol-5-yloxy)-7-(3-(1,1-**

dioxothiomorpholino)propoxy)-6-methoxyquinazoline.

¹H NMR Spectrum: (DMSO-d₆) 1.95-2.05 (m, 2H), 2.15 (s, 3H), 2.35 (s, 3H), 2.7 (t, 2H), 2.95 (br s, 4H), 3.15 (br s, 4H), 4.0 (s, 3H), 4.29 (t, 2H), 6.9 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.61 (s, 1H), 8.5 (s, 1H)

n) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 2-(4-pyridyloxy)ethanol (81 mg), (J. Chem. Soc. Perkin Trans 2, 1987, 12, 1867), to give **4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(2-(4-pyridyloxy)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.35 (s, 3H), 4.0 (s, 3H), 4.55 (m, 2H), 4.6 (m, 2H), 6.88 (dd, 1H), 7.08 (d, 2H), 7.22 (s, 1H), 7.28 (d, 1H), 7.48 (s, 1H), 7.6 (s, 1H), 8.42 (d, 2H), 8.5 (s, 1H), 10.78 (s, 1H)

o) 4-(2,3-Dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (97 mg) was reacted with 3-(methylsulphonyl)-1-propanol (80 mg), (prepared as described for the starting material in Example 50), to give **4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(3-**

methylsulphonylpropoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 1.8-1.9 (m, 2H), 2.15 (s, 3H), 2.25-2.35 (m, 2H), 2.35 (s, 3H), 3.0 (s, 3H), 4.02 (s, 3H), 4.35 (t, 2H), 6.9 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.7 (s, 1H), 8.52 (s, 1H)

Example 107

Using an analogous procedure to that described in Example 91, 7-hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89mg) was reacted with 2-(2-methoxyethoxy)ethanol (70mg)

to give 4-(indol-5-yloxy)-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline (50mg, 42%).

¹H NMR Spectrum: (DMSO-d₆) 3.3 (s, 3H), 3.5 (m, 2H), 3.65 (m, 2H), 3.85 (m, 2H), 4.02 (s, 3H), 4.35 (t, 2H), 6.58 (s, 1H), 7.0 (dd, 1H), 7.4 (s, 1H), 7.45 (br s, 2H), 7.47 (d, 1H), 7.61 (s, 1H), 8.5 (s, 1H)

MS-ESI : 410 [MH]⁺

The starting material was prepared as follows:

A mixture of 7-benzyloxy-4-chloro-6-methoxyquinazoline (3g, 10 mmol), (prepared as described for the starting material in Example 1), 5-hydroxyindole (1.46g, 11 mmol) in DMF (30ml) containing potassium carbonate (2.75g, 20 mmol) was heated at 95°C for 2 hours. After cooling the mixture was poured onto water (100ml). The precipitate was filtered, washed with water and dried under vacuum at 50°C over phosphorus pentoxide. The solid was triturated with ether, filtered, washed with ether and dried under vacuum to give 7-benzyloxy-4-(indol-5-yloxy)-6-methoxyquinazoline (3.5g, 88%).

¹H NMR Spectrum: (DMSO-d₆) 4.02 (s, 3H), 5.4 (s, 2H), 6.5 (s, 1H), 7.0 (dd, 1H), 7.4-7.6 (m, 9H), 7.65 (s, 1H), 8.5 (s, 1H), 11.23 (s, 1H)

MS-ESI : 398 [MH]⁺

A solution of 7-benzyloxy-4-(indol-5-yloxy)-6-methoxyquinazoline (8 g, 20 mmol) in DMF (50 ml) and methylene chloride (100 ml) containing 10% palladium on charcoal (2 g) was hydrogenated at 1.8 atmospheres pressure until uptake of hydrogen had ceased. The solution was filtered, the catalyst was washed with DMF and the filtrate was evaporated. The residue was purified by column chromatography eluting with methylene chloride, followed by methylene chloride/methanol (95/5 and 90/10). After evaporation of the solvent, the residue was triturated with ether, filtered and dried under vacuum to give 7-hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (2.7 g; 44 %).


¹H NMR Spectrum: (DMSO-d₆) 4.0 (s, 3H), 6.46 (s, 1H), 7.01 (dd, 1H), 7.2 (s, 1H), 7.4-7.5 (m, 3H), 7.6 (s, 1H), 8.41 (s, 1H)

Examples 108-118

Using an analogous procedure to that described in Example 107, the appropriate alcohol was reacted with 7-hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline, (prepared as



described for the starting material in Example 107), to give the compounds described in the Table VIII below.

Table VIII



5

Example number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	R	Note
108	58	49	407		r
109	14	13	379		s
110	55	48	393		t
111	27	23	405		u
112	58	47	421		v
113	63	52	419		w
114	64	53	419		x
115	106	84	435		y
116	76	62	423		z

117	113	81	483		aa
118	24	19	429		bb

r) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with N,N-diethylethanolamine (68 mg) to give **7-(2-(N,N-diethylamino)ethoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline**.

5

s) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with N,N-dimethylethanolamine (52 mg) to give **7-(2-(N,N-dimethylamino)ethoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.3 (s, 6H), 2.8 (t, 2H), 4.0 (s, 3H), 4.3 (t, 2H), 6.45 (s, 1H), 7.0 (dd, 1H), 7.4-7.5 (m, 4H), 7.6 (s, 1H), 8.5 (s, 1H)

10

t) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with 3-(N,N-dimethylamino)propan-1-ol (60 mg) to give **7-(3-(N,N-dimethylamino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline**.

15

¹H NMR Spectrum: (DMSO-d₆) 1.9-2.05 (m, 2H), 2.21 (s, 6H), 2.45 (t, 2H), 4.02 (s, 3H), 4.25 (t, 2H), 6.47 (s, 1H), 7.0 (dd, 1H), 7.38 (s, 1H), 7.35-7.4 (m, 2H), 7.45 (d, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

20

u) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with (2*S*)-2-(hydroxymethyl)-1-methylpyrrolidine (67 mg) to give **(2*S*)-4-(indol-5-yloxy)-6-methoxy-7-(1-methylpyrrolidin-2-yl)quinazoline**.

25

v) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with 3-(N,N-diethylamino)-1-propanol (76 mg) to give **7-(3-(N,N-diethylamino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 0.95 (t, 6H), 1.9-2.0 (m, 2H), 2.5 (m, 4H), 2.6 (t, 2H), 4.0 (s, 3H), 4.25 (t, 2H), 6.48 (s, 1H), 7.0 (dd, 1H), 7.38 (s, 1H), 7.42-7.5 (m, 3H), 7.6 (s, 1H), 8.5 (s, 1H)

w) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with 2-(2-hydroxyethyl)piperidine (75 mg) to give **4-(indol-5-yloxy)-6-methoxy-7-(2-(piperidin-2-yl)ethoxy)quinazoline**.

5 ¹H NMR Spectrum: (DMSO-d₆) 1.45-1.75 (m, 3H), 1.75-1.85 (m, 2H), 2.0-2.1 (m, 1H), 2.1-2.2 (m, 1H), 2.25-2.35 (m, 1H), 2.95 (t, 1H), 3.3-3.4 (m, 2H), 4.1 (s, 3H), 4.4-4.5 (m, 2H), 6.5 (s, 1H), 7.05 (dd, 1H), 7.45-7.6 (m, 4H), 7.75 (s, 1H), 9.0 (s, 1H)

10 x) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with 1-(2-hydroxyethyl)piperidine (75 mg) to give **4-(indol-5-yloxy)-6-methoxy-7-(2-(piperidin-1-yl)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.1-1.3 (m, 1H), 1.35-1.5 (m, 1H), 1.65-1.8 (m, 2H), 1.8-1.9 (m, 2H), 3.1 (t, 2H), 3.6 (d, 2H), 3.65 (t, 2H), 4.1 (s, 3H), 4.7 (t, 2H), 6.5 (d, 1H), 7.05 (dd, 1H), 7.45 (s, 1H), 7.5-7.55 (m, 2H), 7.61 (s, 1H), 7.8 (s, 1H), 9.0 (m, 1H)

15

y) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with 4-(3-hydroxypropyl)morpholine (84 mg), (prepared as described for the starting material in Example 60), to give **4-(indol-5-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline**.

20 ¹H NMR Spectrum: (DMSO-d₆) 1.9-2.1 (m, 2H), 2.4 (br s, 4H), 2.5 (t, 2H), 3.6 (t, 4H), 4.0 (s, 3H), 4.25 (t, 2H), 6.45 (s, 1H), 7.0 (dd, 1H), 7.4 (s, 1H), 7.4-7.45 (m, 2H), 7.47 (d, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

25 z) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with 2-(N-(2-methoxyethyl)-N-methylamino)ethanol (77 mg), (prepared as described for the starting material in Example 59), to give **4-(indol-5-yloxy)-6-methoxy-7-(2-(N-(2-methoxyethyl)-N-methylamino)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.35 (s, 3H), 2.65 (t, 2H), 2.9 (t, 2H), 3.25 (s, 3H), 3.45 (t, 2H), 4.0 (s, 3H), 4.3 (t, 2H), 6.45 (s, 1H), 7.05 (dd, 1H), 7.4-7.5 (m, 4H), 7.6 (s, 1H), 8.5 (s, 1H)

30

aa) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with 3-(1,1-dioxothiomorpholino)-1-propanol (112 mg), (prepared as described for the starting material in

Example 5), to give 7-(3-(1,1-dioxothiomorpholino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.0 (m, 2H), 2.65 (m, 2H), 2.9 (br s, 4H), 3.15 (br s, 4H), 4.0 (s, 3H), 4.25 (t, 2H), 6.5 (s, 1H), 7.0 (dd, 1H), 7.35-7.5 (m, 4H), 7.65 (s, 1H), 8.5 (s, 1H)

5

bb) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (89 mg) was reacted with 2-(4-pyridyloxy)ethanol (81 mg), (J. Chem. Soc. Perkin Trans 2, 1987, 12, 1867), to give 4-(indol-5-yloxy)-6-methoxy-7-(2-(4-pyridyloxy)ethoxy)quinazoline.

10 Example 119

A solution of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (200 mg, 0.59 mmol), (prepared as described for the starting material in Example 67), 6-hydroxyindole (96 mg, 0.715 mmol) in DMF (3 ml) containing cesium carbonate (291 mg, 0.894 mmol) was heated at 90°C for 4 hours. After cooling, the mixture was diluted with water, the precipitate was filtered, washed with water and dried under vacuum. The solid was purified by column chromatography eluting with methylene chloride/methanol (90/10 increasing to 50/50) to give 4-(indol-6-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline (240 mg, 93 %).

15

¹H NMR Spectrum: (DMSO-d₆) 1.35-1.45 (m, 2H), 1.45-1.55 (m, 4H), 1.9-2.05 (m, 2H), 2.3-2.4 (m, 4H), 2.45 (t, 2H), 4.0 (s, 3H), 4.22 (t, 2H), 6.5 (s, 1H), 6.9 (dd, 1H), 7.3 (s, 1H), 7.35-7.40 (m, 2H), 7.55-7.65 (m, 2H), 8.5 (s, 1H)

20

MS-ESI : 433 [MH]⁺

Elemental analysis	Found	C 68.4	H 6.4	N 12.8
C ₂₅ H ₂₈ N ₄ O ₃ 0.4 H ₂ O	Requires	C 68.3	H 6.6	N 12.7%

25 Example 120

A solution of 4-chloro-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline (200 mg, 0.6 mmol), (prepared as described for the starting material in Example 50), and 6-hydroxyindole (97 mg, 0.73 mmol) in DMF (3 ml) containing potassium carbonate (125 mg, 0.91 mmol) was heated at 90°C for 2.5 hours. After cooling, water was added. The precipitate was filtered, washed with water and dried under vacuum. The residue was triturated with ether, filtered, washed with ether and dried under vacuum to give 4-(indol-6-yloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline (130 mg, 50 %).

30

¹H NMR Spectrum: (DMSO-d₆) 2.2-2.35 (m, 2H), 3.05 (s, 3H), 3.3 (m, 2H), 4.0 (s, 3H), 4.35 (t, 2H), 6.48 (s, 1H), 6.9 (dd, 1H), 7.3 (s, 1H), 7.4 (2s, 2H), 7.6 (d, 1H), 7.65 (s, 1H), 7.9 (s, 1H)

MS-ESI : 428 [MH]⁺

5	Elemental analysis	Found	C 56.2	H 4.9	N 9.3
	C ₂₁ H ₂₁ N ₃ O ₅ S 1.1 H ₂ O	Requires	C 56.4	H 5.2	N 9.4%

Example 121

Using an analogous procedure to that described for Example 120, 4-chloro-6-methoxy-

- 10 7-(3-morpholinopropoxy)quinazoline (200 mg, 0.59 mmol), (prepared as described for the starting material in Example 1), was reacted with 6-hydroxyindole (95 mg, 0.71 mmol) to give **4-(indol-6-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline** (155 mg, 60 %).

¹H NMR Spectrum: (DMSO-d₆) 1.95-2.05 (m, 2H), 2.4 (br s, 4H), 2.48 (t, 2H), 3.6 (t, 4H), 4.0 (s, 3H), 4.27 (t, 2H), 6.5 (s, 1H), 6.93 (dd, 1H), 7.3 (s, 1H), 7.4 (br s, 2H), 7.6 (d, 1H), 7.61 (s, 1H), 8.5 (s, 1H)

MS-ESI : 435 [MH]⁺

Elemental analysis	Found	C 62.0	H 6.2	N 12.1
C ₂₄ H ₂₆ N ₄ O ₄ 1.6 H ₂ O	Requires	C 62.2	H 6.4	N 12.1%

20 Example 122

- A suspension of 7-(1-*tert*-butoxycarbonylpiperidin-4-ylmethoxy)-4-(2-methylindol-5-yloxy)quinazoline (150 mg, 0.31 mmol), (prepared as described in Example 90), in methylene chloride (2 ml) and TFA (1.5 ml) was stirred for 1 hour at ambient temperature. After removal of the volatiles under vacuum the residue was azeotroped with toluene. The residue was partitioned between methylene chloride and water and the aqueous layer was adjusted to pH 11. The organic layer was separated, washed with brine, dried (MgSO₄), and evaporated. The residue was triturated with ether, filtered, washed with ether and dried under vacuum to give **4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline** (80 mg, 67 %).

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.5-1.65 (m, 2H), 2.0 (d, 2H), 2.15-2.3 (m, 1H), 2.4 (s, 3H), 2.95 (t, 2H), 3.38 (d, 2H), 4.2 (d, 2H), 6.2 (s, 0.5H, partially exchanged), 6.9 (dd, 1H), 7.35 (s, 1H), 7.4 (d, 1H), 7.5 (s, 1H), 7.58 (dd, 1H), 8.5 (d, 1H), 9.1 (s, 1H)

MS-ESI : 389 [MH]⁺

- 166 -

Elemental analysis	Found	C 68.9	H 6.2	N 13.7
$C_{23}H_{24}N_4O_2 \cdot 0.2 H_2O \cdot 0.12 CH_2Cl_2$	Requires	C 69.0	H 6.2	N 13.9%

Example 123

- 5 Using an analogous procedure to that described for Example 71, 4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline (150 mg, 0.386 mmol), (prepared as described in Example 122), was reacted with methoxyacetaldehyde (83 mg, 0.772 mmol), (prepared as described for the starting material in Example 71), to give **7-(1-(2-methoxyethyl)piperidin-4-ylmethoxy)-4-(2-methylindol-5-yloxy)quinazoline** (80 mg, 46 %).
- 10 ¹H NMR Spectrum: (DMSO-d₆) 1.3-1.42 (m, 2H), 1.7-1.9 (m, 3H), 2.0 (t, 2H), 2.4 (s, 3H), 2.48 (t, 2H), 2.92 (d, 2H), 3.22 (s, 3H), 3.42 (t, 2H), 4.05 (d, 2H), 6.15 (s, 1H), 6.88 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.35 (s, 1H), 7.37 (d, 1H), 8.28 (d, 1H), 8.6 (s, 1H)
- MS-ESI : 447 [MH]⁺
- | | | | | |
|--|----------|--------|-------|---------|
| Elemental analysis | Found | C 68.4 | H 6.7 | N 12.2 |
| 15 $C_{26}H_{30}N_4O_3 \cdot 0.5 H_2O$ | Requires | C 68.6 | H 6.9 | N 12.3% |

Example 124

- Diethyl azodicarboxylate (117 mg, 0.67 mmol) was added in portions to a solution of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (120 mg, 0.37 mmol), (prepared as described in Example 49), and 3-(ethylsulphonyl)-1-propanol (74 mg, 0.48 mmol) in methylene chloride (3.5 ml) and triphenylphosphine (176 mg, 0.67 mmol). After stirring for 2 hours at ambient temperature, the residue was poured onto a column of silica and eluted with ethyl acetate/methylene chloride (1/1) followed by methylene chloride/methanol (97/3 followed by 95/5). After removal of the solvent under vacuum, the residue was triturated with ether, filtered and dried under vacuum to give **7-(3-(ethylsulphonyl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (93 mg, 55 %).
- 25 ¹H NMR Spectrum: (DMSO-d₆) 1.25 (t, 3H), 2.2-2.3 (m, 2H), 2.4 (s, 3H), 3.2 (q, 2H), 3.3 (t, 2H), 4.0 (s, 3H), 4.35 (t, 2H), 6.15 (s, 1H), 6.9 (dd, 1H), 7.28 (s, 1H), 7.32 (d, 1H), 7.4 (s, 1H), 7.62 (s, 1H), 8.5 (s, 1H)
- 30 MS-ESI : 456 [MH]⁺
- | | | | | |
|-----------------------|----------|--------|-------|--------|
| Elemental analysis | Found | C 60.3 | H 5.6 | N 9.2 |
| $C_{23}H_{25}N_3O_5S$ | Requires | C 60.6 | H 5.5 | N 9.2% |

The starting material was prepared as follows:

A solution of ethylthiopropanol (1.2 g, 10 mmol) in methylene chloride (30 ml) containing 3-chloroperoxybenzoic acid (5 g, 20 mmol) was stirred at ambient temperature for 30 minutes. The precipitate was filtered, washed with methylene chloride and the filtrate was poured onto a column of aluminium oxide and eluted with methylene chloride, followed by methylene chloride/methanol (95/5 and 90/10). After removal of the solvent, the residue was dissolved in methylene chloride, dried (MgSO_4) and evaporated to give 3-(ethylsulphonyl)-1-propanol (1.05 g, 69 %).

^1H NMR Spectrum: (DMSO-d_6) 1.25 (t, 3H), 1.75-1.9 (m, 2H), 3.0-3.2 (m, 4H), 3.5 (q, 2H), 4.7 (t, 1H)

MS-ESI : 153 $[\text{MH}]^+$

Example 125

Using an analogous procedure to that described for Example 124, 4-(2,3-dimethylindol-5-yloxy)-7-hydroxy-6-methoxyquinazoline (120 mg, 0.36 mmol), (prepared as described for the starting material in Example 91), was reacted with 3-(ethylsulphonyl)-1-propanol (71 mg, 0.46 mmol), (prepared as described for the starting material in Example 124), to give 4-(2,3-dimethylindol-5-yloxy)-7-(3-ethylsulphonylpropoxy)-6-methoxyquinazoline (96 mg, 57 %).

^1H NMR Spectrum: (DMSO-d_6) 1.25 (t, 3H), 2.15 (s, 3H), 2.2-2.3 (m, 2H), 2.35 (s, 3H), 3.2 (q, 2H), 3.3 (t, 2H), 4.02 (s, 3H), 4.35 (t, 2H), 6.9 (dd, 1H), 7.22 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.63 (s, 1H), 8.51 (s, 1H)

MS-ESI : 470 $[\text{MH}]^+$

Elemental analysis	Found	C 60.6	H 6.0	N 8.8
$\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}_5\text{S} \cdot 0.4 \text{H}_2\text{O}$	Requires	C 60.5	H 5.9	N 8.8%

Example 126

Using an analogous procedure to that described for Example 124, 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (128 mg, 0.4 mmol), (prepared as described in Example 49), was reacted with 4-(2-hydroxyethyl)-(1-*tert*-butoxycarbonyl)piperidine (119 mg, 0.52 mmol) overnight to give 7-(2-(1-*tert*-butoxycarbonylpiperidin-4-yl)ethoxy)-6-

methoxy-4-(2-methylindol-5-yloxy)quinazoline (34 mg, 16 %).

¹H NMR Spectrum: (DMSO-d₆) 1.05-1.2 (m, 2H), 1.42 (s, 9H), 1.62-1.85 (m, 5H), 2.42 (s, 3H), 2.62-2.82 (m, 2H), 3.9-4.0 (m, 2H), 4.0 (s, 3H), 4.25 (t, 2H), 6.17 (s, 1H), 6.9 (dd, 1H), 7.3 (d, 1H), 7.32 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

5 MS-ESI : 533 [MH]⁺

Elemental analysis	Found	C 67.8	H 6.9	N 10.5
C ₃₀ H ₃₆ N ₄ O ₅	Requires	C 67.7	H 6.8	N 10.5%

The starting material was prepared as follows:

10 A solution of 4-(2-hydroxyethyl)pyridine (1.8 g, 14.6 mol) in acetic acid (15 ml) containing platinum oxide (200 mg) was hydrogenated for 20 hours at 3.3-4 atmospheres pressure. After filtration, the filtrate was evaporated and azeotroped twice with toluene. The residue was triturated with 2N sodium hydroxide and solid sodium hydroxide was added to adjust the pH to 13. The volatiles were removed under vacuum and the residue was triturated
15 with ether, filtered, washed with methylene chloride, and dried under vacuum to give 2-(piperidin-4-yl)-1-ethanol (860 mg, 46 %).

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.3-1.5 (m, 4H), 1.6-1.7 (m, 1H), 1.7-1.9 (d, 2H), 1.75 (t, 2H), 3.25 (d, 2H), 3.55 (t, 2H)

A solution of 2-(piperidin-4-yl)-1-ethanol (830 mg, 6.4 mmol) in DMF (5 ml)
20 containing tertbutyl dicarbonate anhydride (1.4 g, 6.4 mol) was stirred at ambient temperature for 48 hours. After removal of the volatiles under vacuum, the residue was partitioned between ether and water. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated to give 4-(2-hydroxyethyl)-(1-*tert*-butoxycarbonyl)piperidine (1 g, 68 %).

25 ¹H NMR Spectrum: (DMSO-d₆) 0.9-1.1 (m, 2H), 1.3-1.6 (m, 3H), 1.4 (s, 9H), 1.6 (d, 2H), 2.5-2.8 (br s, 2H), 3.45 (dd, 2H), 3.9 (d, 2H), 4.35 (t, 1H)

Example 127

Using an analogous procedure to that described for Example 121, 4-chloro-6-methoxy-
30 7-(3-morpholinopropoxy)quinazoline (160 mg, 0.47 mol), (prepared as described for the starting material in Example 1), was reacted with 6-hydroxy-2-methylindole (84 mg, 0.57 mol), (Eur. J. Med. Chem. 1975, 10, 187), to give **6-methoxy-4-(2-methylindol-6-yloxy)-7-**

(3-morpholinopropoxy)quinazoline (157 mg, 73 %).

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.25-2.35 (m, 2H), 2.38 (s, 3H), 3.15 (t, 2H), 3.35 (t, 2H), 3.5 (d, 2H), 3.68 (t, 2H), 4.0 (d, 2H), 4.05 (s, 3H), 4.35 (t, 2H), 6.18 (s, 1H), 6.9 (d, 1H), 7.22 (s, 1H), 7.45 (d, 1H), 7.52 (s, 1H), 7.8 (s, 1H), 9.05 (s, 1H)

5 MS-ESI : 449 [MH]⁺

Elemental analysis	Found	C 66.4	H 6.4	N 12.4
C ₂₅ H ₂₈ N ₄ O ₄ 0.2 H ₂ O	Requires	C 66.4	H 6.3	N 12.4%

Example 128

10 Using an analogous procedure to that described for the synthesis of 4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline, (prepared as described in Example 122), 7-(2-(1-*tert*-butoxycarbonylpiperidin-4-yl)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (400 mg, 0.75 mmol), (prepared as described in Example 126), was used to give 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(piperidin-4-yl)ethoxy)quinazoline (284
15 mg, 87 %).

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.3-1.5 (m, 2H), 1.8-2.0 (m, 5H), 2.4 (s, 3H), 2.9 (t, 2H), 3.3 (d, 2H), 4.05 (s, 3H), 4.35 (t, 2H), 6.2 (s, 1H), 6.95 (dd, 1H), 7.35 (s, 1H), 7.37 (d, 1H), 7.52 (s, 1H), 7.8 (s, 1H), 9.1 (s, 1H)

MS-ESI : 433 [MH]⁺

20

Example 129

Diethyl azodicarboxylate (65 µl, 0.4 mmol) was added in portions to a suspension of 4-(2,3-dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol), triphenylphosphine (107 mg, 0.4 mmol), (E)-4-(pyrrolidin-1-yl)but-2-en-1-ol (40 mg, 0.28
25 mmol) in DMF (0.4 ml) and dichloromethane (1.5 ml) cooled at 0°C. The reaction mixture was left to warm up to ambient temperature and was stirred overnight. The mixture was poured onto a column of silica and was eluted with methylene chloride followed by methylene chloride/methanol (98/2), followed by methylene chloride/3N ammonia in methanol (95/5 and 90/10) to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-((E)-4-(pyrrolidin-1-yl)but-2-
30 en-1-yloxy)quinazoline (51 mg, 55 %).

¹H NMR Spectrum: (DMSO-d₆) 1.6-1.7 (m, 4H), 2.15 (s, 3H), 2.3 (s, 3H), 2.4 (br s, 4H), 3.1 (d, 2H), 3.97 (s, 3H), 4.7 (d, 2H), 5.8-6.0 (m, 2H), 7.15 (s, 1H), 7.22 (d, 1H), 7.3 (d, 1H), 7.55

- 170 -

(s, 1H), 7.87 (s, 1H), 8.3 (s, 1H), 9.4 (s, 1H), 10.62 (s, 1H)

MS-ESI : 458 [MH]⁺

The starting material was prepared as follows:

- 5 Thionyl chloride (9.3ml, 128mmol) was added in portions to a stirred solution of 2-butyne-1,4-diol (10g, 116mmol) in toluene (15ml) and pyridine (10.3ml) cooled at 0°C. The mixture was stirred for 3.5 hours at ambient temperature and then poured onto ice water. The mixture was extracted with ether, the organic layer was washed with saturated aqueous sodium hydrogen carbonate solution and then brine, dried (MgSO₄) and the volatiles removed
10 by evaporation. The residue was purified by column chromatography eluting with petroleum ether/ether (7/3) to give 4-chlorobut-2-yn-1-ol (4.74g, 39%).

¹H NMR Spectrum: (CDCl₃) 1.68(t, 1H); 4.18(d, 2H); 4.33(d, 2H)

- Pyrrolidine (7.8ml, 94mmol) was added dropwise to a solution of 4-chlorobut-2-yn-1-ol (4.74g, 45mmol) in toluene (40ml) and the mixture stirred and heated at 60°C for 1 hour.
15 The volatiles were removed by evaporation and the residue was purified by chromatography eluting with methylene chloride/methanol (96/4) to give 4-(pyrrolidin-1-yl)but-2-yn-1-ol (4.3g, 69%).

¹H NMR Spectrum: (CDCl₃) 1.82(t, 4H); 2.63(t, 4H); 3.44(t, 2H), 4.29(t, 2H)

- A solution of 4-(pyrrolidin-1-yl)but-2-yn-1-ol (4.3g, 31mmol) in THF (20ml) was
20 added dropwise to a suspension of lithium aluminium hydride (2.35g, 62mmol) in anhydrous THF (8ml) and the mixture stirred and heated at 60°C for 2 hours. The mixture was cooled to 5°C and 2M aqueous sodium hydroxide solution (28ml) was added dropwise. The resulting suspension was filtered and the volatiles removed from the filtrate by evaporation. The residue was dissolved in a mixture of methylene chloride/ethyl acetate, dried (MgSO₄) and the
25 solvent removed by evaporation. The residue was purified by column chromatography on aluminum oxide eluting with methylene chloride/methanol (97/3) to give (E)-4-(pyrrolidin-1-yl)but-2-en-1-ol (3.09g, 70%).

¹H NMR Spectrum: (CDCl₃) 1.82(m, 4H); 2.61(m, 4H); 3.17(m, 2H); 4.13(s, 2H); 5.84(m, 2H)

- 30 A solution of 4-chloro-6-methoxy-7-benzyloxyquinazoline (7g, 23 mmol), (prepared as described for the starting material in Example 1), and 5-amino-2,3-dimethylindole (4.5g, 28 mmol) in isopropanol (90ml) containing 6.2 N hydrogen chloride in isopropanol (380μl) was

heated at reflux for 3 hours and stirred overnight at ambient temperature. The mixture was triturated with ether and the solid was filtered, washed with ether and dried under vacuum to give 7-benzyloxy-4-(2,3-dimethylindol-5-ylamino)-6-methoxyquinazoline (10.5 g, quant.).

¹H NMR Spectrum: (DMSO-d₆) 2.16 (s, 3H), 2.33 (s, 3H), 4.0 (s, 3H), 5.34 (s, 2H), 7.2 (d, 1H), 7.32 (d, 1H), 7.35-7.55 (m, 7H), 8.2 (s, 1H), 8.7 (s, 1H), 10.9 (s, 1H), 11.15 (s, 1H)

MS-ESI : 425 [MH]⁺

Ammonium formate (20g, 326 mmol) and 10% palladium on carbon (1g) were added to a solution of 7-benzyloxy-4-(2,3-dimethylindol-5-ylamino)-6-methoxyquinazoline (10g, 22 mmol) in DMF (100ml) and methanol (300ml). After stirring for 3 hours at ambient temperature, aqueous ammonia (120ml) was added. The precipitate was filtered, washed with water and dried under vacuum. The residue was triturated with ethyl acetate and ether and was filtered, dried under vacuum and purified by column chromatography eluting with methanol/methylene chloride (5/95 followed by 10/90) to give 4-(2,3-dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (5.5g, 75%).


¹H NMR Spectrum: (DMSO-d₆) 2.2 (s, 3H), 2.35 (s, 3H), 3.97 (s, 3H), 7.0 (s, 1H), 7.22 (d, 1H), 7.3 (d, 1H), 7.55 (s, 1H), 7.85 (s, 1H), 8.28 (s, 1H), 9.35 (s, 1H), 10.2 (br s, 1H), 10.62 (s, 1H)

MS-ESI : 335 [MH]⁺

Examples 130-145














Using an analogous procedure to that described in Example 129, 4-(2,3-dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol), (prepared as described for the starting material in Example 129), was reacted with the appropriate alcohol to give the compounds described in Table IX.




Table IX



Example	Weight (mg)	Yield %	MS-ESI	R	Note
---------	-------------	---------	--------	---	------



number			[MH] ⁺		
130	10	11	458		a
131	63	69	450		b
132	5	6	443		c
133	35	36	475		d
134	53	51	510		e
135	56	58	469		f
136	4	4.6	415		g
137	29	35	406		h
138	49	56	432		i
139	8	8.6	481		j
140	15	15	477		k
141	38	42	446		l
142	69	72	470		m

143	21	21	492		n
144	36	40	440		o
145	31	33	460		p

a) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 3-(5-methyl-[1,2,4]-triazol-1-yl)propan-1-ol (40 mg) to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(3-(5-methyl-1H-[1,2,4]-triazol-1-yl)propoxy)quinazoline.

The starting material was prepared as follows:

Under argon, 1,2,4-triazole (13.8g, 200mmol) was added to a solution of sodium ethoxide (freshly prepared from sodium (4.6g) and ethanol (250ml)). After complete dissolution, 3-bromopropan-1-ol (18ml, 200 mmol) was added dropwise. The mixture was refluxed for 18 hours and the solid was filtered and washed with ethanol. The filtrate was evaporated and the residue was purified by column chromatography eluting with methylene chloride/methanol (9/1) to give 3-(1,2,4-triazol-1-yl)propan-1-ol (22.8 g, 90%).

¹H NMR Spectrum: (CDCl₃): 2.12 (m, 2H) ; 2.6 (br s, 1H) ; 3.65 (t, 2H) ; 4.35 (t, 2H) ; 7.95 (s, 1H) ; 8.1 (s, 1H)

To a solution of 3-(1,2,4-triazol-1-yl)propan-1-ol (7 g, 55 mmol) in DMF (70ml) was added *tert*butyldimethylsilyl chloride (9.1g, 60 mmol) followed by DMAP (336mg, 2.7 mmol) followed by imidazole (4.5gr, 66 mmol). After stirring overnight at ambient temperature, the volatiles were removed under vacuum and the residue was partitioned between water and ethyl acetate/ether. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with methylene chloride/ether (6/4) to give 3-(*tert*butyldimethylsilyloxy)-1-(1,2,4-triazol-1-yl)propane (11.1 gr, 84%).

MS-EI : 242 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 0.25 (s, 6H) ; 0.9 (s, 9H) ; 2.05 (m, 2H) ; 3.52 (t, 2H) ; 4.25 (t,

2H) ; 7.9 (s, 1H) ; 8.02 (s, 1H)

To a solution of 3-(*tert*butyldimethylsilyloxy)-1-(1,2,4-triazol-1-yl)propane (7 g, 29 mmol) in DMF (100ml) cooled at -70°C was added 2.5M *n*-butyllithium (17.4 ml) over 45 minutes. After stirring for 90 minutes at -70°C, methyl iodide (3.6ml, 58 mmol) was added.

- 5 After stirring for 2 hours at ambient temperature, the mixture was poured onto saturated ammonium chloride. The mixture was then diluted with ether and ethyl acetate. The organic layer was separated, washed with aqueous sodium thiosulphate followed by brine, dried (MgSO₄) and evaporated to give 3-(*tert*butyldimethylsilyloxy)-1-(5-methyl-[1,2,4]-triazol-1-yl)propane (7.3 g, 98%).

10 MS-EI : 256 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 0.25 (s, 6H) ; 0.85 (s, 9H) ; 2.0 (s, 2H) ; 2.4 (s, 3H) ; 3.52 (t, 2H) ; 4.15 (t, 2H) ; 7.72 (s, 1H)

To a solution of ammonium fluoride (10.4 g, 280 mmol) in methanol (110ml) was added a solution of 3-(*tert*butyldimethylsilyloxy)-1-(5-methyl-[1,2,4]-triazol-1-yl)propane (7.2 g, 28 mmol) in methanol (30ml). The mixture was refluxed for 4.5 hours. After cooling, silica (100g) was added and the volatiles were removed under vacuum. The residue was added onto a column of silica and eluted with a mixture of methylene chloride/ethyl acetate (1/1) followed by methylene chloride/methanol (9/1) to give 3-(5-methyl-[1,2,4]-triazol-1-yl)propan-1-ol (3.65 g, 92%).

20 MS-ESI : 142 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 2.05 (m, 2H) ; 2.5 (s, 3H) ; 3.62 (t, 2H) ; 4.25 (t, 2H) ; 7.8 (s, 1H)

b) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol)

- 25 was reacted with 2-(*N*-(2-methoxyethyl)-*N*-methylamino)ethanol (38 mg), (prepared as described for the starting material in Example 59), to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(*N*-(2-methoxyethyl)-*N*-methylamino)ethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.35 (s, 6H), 2.65 (t, 2H), 2.85 (t, 2H), 3.25 (s, 3H), 3.45 (t, 2H), 3.95 (s, 3H), 4.2 (t, 2H), 7.15 (s, 1H), 7.22 (s, 1H), 7.3 (dd, 1H), 7.55 (s, 30 1H), 7.85 (s, 1H), 8.3 (s, 1H), 9.4 (s, 1H), 10.62 (s, 1H)

c) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was

reacted with 2-(1-methylimidazol-2-yl)ethanol (36 mg), (EP 06751112 A1), to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(1-methylimidazol-2-yl)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.32 (s, 3H), 3.2 (t, 2H), 3.7 (s, 3H), 3.95 (s, 3H), 4.45 (t, 2H), 6.8 (s, 1H), 7.05 (s, 1H), 7.15 (s, 1H), 7.22 (d, 1H), 7.3 (dd, 1H), 7.55 (s, 1H),
5 7.88 (s, 1H), 8.32 (s, 1H), 9.4 (s, 1H), 10.62 (s, 1H)

d) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 1-(3-hydroxypropyl)-4-methylpiperazine (45 mg) to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline**.

10 ¹H NMR Spectrum: (DMSO-d₆) 1.9-2.0 (m, 2H), 2.15 (2s, 6H), 2.0-2.9 (m, 8H), 2.32 (s, 3H), 2.45 (t, 2H), 3.95 (s, 3H), 4.2 (t, 2H), 7.1 (s, 1H), 7.22 (d, 1H), 7.3 (dd, 1H), 7.55 (s, 1H), 7.85 (s, 1H), 8.3 (s, 1H), 9.4 (s, 1H), 10.62 (s, 1H)

The starting material was prepared as follows:

15 3-Bromopropan-1-ol (20ml, 20mmol) was added dropwise to a solution of 1-methylpiperazine (29ml, 26 mmol) in ethanol (200ml). Potassium carbonate (83 gr, 60 mmol) was added and the mixture was refluxed for 20 hours. After cooling, the solid was filtered and the filtrate was evaporated. The residue was triturated with ether, filtrate and evaporated. The residue was distilled at about 60-70°C under about 0.2 mm Hg to give 1-(3-
20 hydroxypropyl)-4-methylpiperazine (17g, 53%).

¹H NMR Spectrum: (CDCl₃) 1.72 (m, 2H) ; 2.3 (s, 3H) ; 2.2-2.8 (m, 8H) ; 2.6 (t, 2H) ; 3.8 (t, 2H) ; 5.3 (br s, 1H)

e) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was
25 reacted with 3-(1,1-dioxothiomorpholino)-1-propanol (55 mg), (prepared as described for the starting material in Example 5), to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(3-(1,1-dioxothiomorpholino)propoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.9-2.0 (m, 2H), 2.5 (s, 9H), 2.65 (t, 2H), 2.9 (br s, 4H), 3.15 (br s, 4H), 3.95 (s, 3H), 4.25 (t, 2H), 7.2 (s, 1H), 7.85 (s, 1H), 8.0 (dd, 1H), 8.15 (d, 1H), 8.2
30 (s, 1H), 8.45 (s, 1H), 9.6 (s, 1H), 10.95 (s, 1H)

f) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was

reacted with 2-(N-methyl-N-(4-pyridyl)amino)ethanol (43mg), (EP 0359389), to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(N-methyl-N-(4-pyridyl)amino)ethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.35 (s, 3H), 3.07 (s, 3H), 3.85 (t, 2H), 3.95 (s, 3H), 4.3 (t, 2H), 6.7 (d, 2H), 7.15 (s, 1H), 7.22 (d, 1H), 7.3 (dd, 1H), 7.55 (s, 1H), 7.85 (s, 1H), 8.15 (d, 2H), 8.3 (s, 1H), 9.4 (s, 1H), 10.65 (s, 1H)

g) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 2-furanmethanol (28 mg) to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-furylmethoxy)quinazoline.

h) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 2-N,N-dimethylethanolamine (25 mg) to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(N,N-dimethylamino)ethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.25 (s, 6H), 2.32 (s, 3H), 2.72 (t, 2H), 3.95 (s, 3H), 4.2 (t, 2H), 7.15 (s, 1H), 7.22 (d, 1H), 7.3 (dd, 1H), 7.55 (s, 1H), 7.85 (s, 1H), 8.32 (s, 1H), 9.4 (s, 1H), 10.6 (s, 1H)

i) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 1-(2-hydroxyethyl)pyrrolidine (33 mg) to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline.

¹H NMR Spectrum: (DMSO-d₆) 1.65-1.75 (m, 4H), 2.15 (s, 3H), 2.35 (s, 3H), 2.55-2.65 (m, 4H), 2.9 (t, 2H), 3.95 (s, 3H), 4.25 (t, 2H), 7.15 (s, 1H), 7.22 (d, 1H), 7.3 (dd, 1H), 7.55 (s, 1H), 7.85 (s, 1H), 8.32 (s, 1H), 9.4 (s, 1H), 10.62 (s, 1H)

j) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with triethylene glycol monomethyl ether (47 mg) to give 4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(2-(2-methoxyethoxy)ethoxy)ethoxy)quinazoline.

k) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 5,5-dimethyl-1,3-dioxane-2-ethanol (46 mg) to give 7-(2-(5,5-dimethyl-1,3-dioxan-2-yl)ethoxy)-4-(2,3-dimethylindol-5-ylamino)-6-methoxyquinazoline.

¹H NMR Spectrum: (DMSO-d₆) 0.7 (s, 3H), 1.15 (s, 3H), 2.05-2.1 (m, 2H), 2.1 (s, 3H), 2.6 (s, 3H), 3.42 (d, 2H), 3.57 (d, 2H), 4.0 (s, 3H), 4.22 (t, 2H), 4.7 (t, 1H), 7.2 (s, 1H), 7.82 (s, 1H), 8.0 (dd, 1H), 8.17 (d, 1H), 8.3 (s, 1H), 8.45 (s, 1H), 9.6 (s, 1H), 10.95 (s, 1H)

- 5 l) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 1-(2-hydroxyethyl)piperidine (37 mg) to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-piperidinoethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.45 (m, 2H), 1.45-1.6 (m, 4H), 2.15 (s, 3H), 2.35 (s, 3H), 2.45 (br s, 4H), 2.75 (t, 2H), 3.95 (s, 3H), 4.25 (t, 2H), 7.15 (s, 1H), 7.22 (d, 1H), 7.3 (dd, 1H),
10 7.55 (s, 1H), 7.85 (s, 1H), 8.3 (s, 1H), 9.4 (s, 1H), 10.62 (s, 1H)

- m) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 2-(N-methyl-N-(pyridazin-4-yl)amino)ethanol (44 mg) to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(N-methyl-N-(pyridazin-4-yl)amino)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.32 (s, 3H), 3.1 (s, 3H), 3.9 (s, 3H), 3.95 (t, 2H), 4.35 (t, 2H), 6.85 (dd, 1H), 7.15 (s, 1H), 7.20 (d, 1H), 7.28 (dd, 1H), 7.55 (s, 1H), 7.85 (s, 1H), 8.3 (s, 1H), 8.58 (d, 1H), 8.9 (d, 1H), 9.4 (s, 1H), 10.62 (s, 1H)

- 20 The starting material was prepared as follows:

A solution of 4-bromo-3,6-dichloro-pyridazine (1.11 g, 5mmol), (J.Chem. Soc., Perkin Trans I, 1974, 696), and 2-(methylamino)ethanol (0.75g, 10mmol) in isopropanol (10ml) was heated at reflux for 30 minutes. The solvent was removed by evaporation, the residue was partitioned between methylene chloride and water and the aqueous layer was
25 adjusted to pH9 with solid potassium carbonate. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was triturated with ether, collected by filtration and dried under vacuum to give 2-(N-(3,6-dichloropyridazin-4-yl)-N-methylamino)ethanol (1g, 90%).

¹H NMR Spectrum: (CDCl₃) 2.1(br s, 1H); 3.09(s, 3H); 3.71(t, 2H); 3.93(t, 2H); 6.8(s, 1H)

- 30 MS - ESI: 221 [MH]⁺

A mixture of 2-(N-(3,6-dichloropyridazin-4-yl)-N-methylamino)ethanol (444mg,

2mmol) and 10% palladium-on-charcoal catalyst (150mg) in ethanol (15ml), methanol (5ml) and aqueous ammonia (15ml) was stirred under hydrogen at 3 atmospheres pressure for 4 hours. The catalyst was removed by filtration and the solvent removed from the filtrate by evaporation. The residue was dissolved in methylene chloride, the insoluble material was removed by filtration and the solvent was removed from the filtrate by evaporation. The residue was purified by column chromatography on neutral aluminum oxide eluting with methylene chloride/methanol (95/5 followed by 90/10). The purified product was triturated with petroleum ether, the solid product was collected by filtration and dried under vacuum to give 2-(N-methyl-N-(pyridazin-4-yl)amino)ethanol (275mg, 91%).

¹H NMR Spectrum: (CDCl₃) 3.06(s, 3H); 3.57(t, 2H); 3.89(t, 2H); 6.52(dd, 1H); 8.48(d, 1H); 8.54 (d, 1H)

MS - ESI: 153 [MH]⁺

n) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 2-(2-morpholinoethoxy)ethanol (50 mg) to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(2-morpholinoethoxy)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.18 (s, 3H), 2.35 (s, 3H), 2.35-2.45 (m, 4H), 2.45-2.5 (m, 2H), 3.5-3.55 (m, 4H), 3.65 (t, 2H), 3.8-3.85 (m, 2H), 3.95 (s, 1H), 4.25 (m, 2H), 7.15 (s, 1H), 7.22 (d, 1H), 7.3 (dd, 1H), 7.55 (s, 1H), 7.85 (s, 1H), 8.3 (s, 1H), 9.4 (s, 1H), 10.62 (s, 1H)

The starting material was prepared as follows :

2-(2-Chloroethoxy)ethanol (1.25g, 10mmol) was added to a mixture of morpholine (2.58g, 30mmol) and potassium carbonate (5.5g, 40mmol) in acetonitrile (50ml). The mixture was heated at reflux for 6 hours and then stirred for 18 hours at ambient temperature. The insolubles were removed by filtration and the volatiles were removed from the filtrate by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (95/5 followed by 90/10 and then 80/20) to give 2-(2-morpholinoethoxy)ethanol (600mg, 34%).

¹H NMR Spectrum: (CDCl₃) 2.5(br s, 4H); 2.59(t, 2H); 3.6-3.85(m, 10H)

MS - (EI): 175 [M]⁺

o) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 3-(2-hydroxyethyl)pyridine (35 mg) to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(2-(3-pyridyl)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H), 2.32 (s, 3H), 3.15 (t, 2H), 3.95 (s, 3H), 4.4 (t, 2H), 7.2 (s, 1H), 7.22 (d, 1H), 7.3 (dd, 1H), 7.35 (dd, 1H), 7.55 (s, 1H), 7.8 (d, 1H), 7.85 (s, 1H), 8.32 (s, 1H), 8.45 (dd, 1H), 8.6 (s, 1H), 9.4 (s, 1H), 10.68 (s, 1H)

p) 4-(2,3-Dimethylindol-5-ylamino)-7-hydroxy-6-methoxyquinazoline (68 mg, 0.2 mmol) was reacted with 1-(3-hydroxypropyl)pyrrolidin-2-one (41 mg) to give **4-(2,3-dimethylindol-5-ylamino)-6-methoxy-7-(3-(2-oxopyrrolidin-1-yl)propoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.9-2.05 (m, 4H), 2.12 (s, 3H), 2.15-2.3 (m, 2H), 2.6 (s, 3H), 3.3-3.45 (m, 4H), 4.0 (s, 3H), 4.15 (t, 2H), 7.15 (s, 1H), 7.82 (s, 1H), 8.0 (dd, 1H), 8.17 (d, 1H), 8.3 (s, 1H), 8.45 (s, 1H), 9.6 (s, 1H), 10.95 (s, 1H)

Example 146

Using an analogous procedure to that described for Example 121, 4-chloro-6-methoxy-7-(3-pyrrolidinopropoxy)quinazoline (150 mg, 0.47 mmol), (prepared as described for the starting material in Example 9), was reacted with 6-hydroxy-2-methylindole (83 mg, 0.56 mol), (Eur. J. Med. Chem. 1975, 10, 187), to give **6-methoxy-4-(2-methylindol-6-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (170 mg, 85 %).

¹H NMR Spectrum: (DMSO-d₆) 1.65-1.8 (m, 4H), 1.95-2.05 (m, 2H), 2.42 (s, 3H), 2.5 (br s, 1H), 2.6 (t, 2H), 4.0 (s, 3H), 4.27 (t, 2H), 6.2 (s, 1H), 6.85 (dd, 1H), 7.2 (s, 1H), 7.4 (s, 1H), 7.45 (d, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

MS-ESI : 433 [MH]⁺

Elemental analysis	Found	C 68.3	H 6.4	N 12.8
C ₂₅ H ₂₈ N ₄ O ₃ 0.4 H ₂ O	Requires	C 68.3	H 6.6	N 12.7%

Example 147

Using an analogous procedure to that described in Example 123, 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(piperidin-4-yl)ethoxy)quinazoline (120 mg, 0.28 mmol) was used to give **7-(2-(1-(2-methoxyethyl)piperidin-4-yl)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (55 mg, 40 %).

¹H NMR Spectrum: (DMSO-d₆) 1.15-1.3 (m, 2H), 1.4-1.55 (m, 1H), 1.65-1.8 (m, 4H), 1.95 (t, 2H), 2.4 (s, 3H), 2.42 (t, 2H), 2.85 (d, 2H), 3.25 (s, 3H), 3.42 (t, 2H), 4.0 (s, 3H), 4.22 (t, 2H), 6.15 (s, 1H), 6.85 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.38 (s, 1H), 7.59 (s, 1H), 8.5 (s, 1H).

MS-ESI : 491 [MH]⁺

5	Elemental analysis	Found	C 65.3	H 7.1	N 10.9
	C ₂₈ H ₃₄ N ₄ O ₄ 1.3 H ₂ O	Requires	C 65.4	H 7.2	N 10.9%

Example 148

Using an analogous procedure to that described in Example 120 OR 121 PER PP, 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (160 mg, 0.48 mmol), (prepared as described for the starting material in Example 1), was reacted with 1,2-dimethyl-5-hydroxyindole (92 mg, 0.57 mol), (Tetrahedron 1994, 50, 13433), to give 4-(1,2-dimethylindol-5-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline (163 mg, 74 %).

¹H NMR Spectrum: (DMSO-d₆) 1.95-2.1 (m, 2H), 2.4 (br s, 4H), 2.45 (s, 3H), 2.5 (t, 2H), 3.65 (t, 4H), 3.75 (s, 3H), 4.0 (s, 3H), 4.25 (t, 2H), 6.25 (s, 1H), 6.95 (dd, 1H), 7.3 (s, 1H), 7.38 (s, 1H), 7.45 (d, 1H), 7.6 (s, 1H), 8.5 (s, 1H)

MS-ESI : 463 [MH]⁺

Elemental analysis	Found	C 67.2	H 6.5	N 12.1
C ₂₆ H ₃₀ N ₄ O ₄	Requires	C 67.5	H 6.5	N 12.1%

Example 149

Using an analogous procedure to that described in Example 124, 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (2.3 g, 7.16 mmol), (prepared as described in Example 49), was reacted with (N-methyl-N-tert-butoxycarbonyl)ethanolamine (1.51 g, 8.6 mmol) to give 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-tert-butoxycarbonylamino)ethoxy)quinazoline (1.93 g, 56 %).

¹H NMR Spectrum: (DMSO-d₆) 1.4 (s, 9H), 2.4 (s, 3H), 2.90 (s, 3H), 3.65 (t, 2H), 4.0 (s, 3H), 4.35 (t, 2H), 6.15 (s, 1H), 6.8 (dd, 1H), 7.28 (s, 1H), 7.35 (d, 1H), 7.42 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H) ;

MS-ESI : 479 [MH]⁺

Elemental analysis	Found	C 65.0	H 6.4	N 11.7
C ₂₆ H ₃₀ N ₄ O ₅ S	Requires	C 65.3	H 6.3	N 11.7%

Example 150

A solution of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(*N*-methyl-*N*-*tert*-butoxycarbonylamino)ethoxy)quinazoline (550 mg, 1.15 mmol), (prepared as described in Example 149), in methylene chloride (10 ml) containing TFA (12 ml) was stirred for 3 hours at ambient temperature. After removal of the volatiles under vacuum, the residue was partitioned between methylene chloride and sodium hydrogen carbonate. The pH of the aqueous layer was adjusted to 11 with 2N sodium hydroxide. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was triturated with ether, filtered and dried under vacuum to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(*N*-methylamino)ethoxy)quinazoline** (356 mg, 82 %).

¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H), 2.5 (s, 3H), 2.9 (t, 2H), 4.0 (s, 3H), 4.25 (t, 2H), 6.25 (s, 1H), 6.9 (dd, 1H), 7.25 (s, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 8.5 (s, 1H), 11.0 (s, 1H)

MS-ESI : 379 [MH]⁺

Elemental analysis	Found	C 64.6	H 5.8	N 14.2
C ₂₁ H ₂₂ N ₄ O ₃ 0.7 H ₂ O	Requires	C 64.5	H 6.0	N 14.3%

Example 151

A mixture of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline (419 mg, 1 mmol), (prepared as described in Example 70), in DMF (6 ml) containing chloroacetonitrile (114 mg, 1.5 mmol), potassium carbonate (346 mg, 2.5 mmol) and potassium iodide (50 mg, 0.3 mmol) was stirred at ambient temperature overnight. The mixture was poured into water and the precipitate was filtered, washed with water and dried under vacuum. The residue was purified by column chromatography, eluting with methylene chloride, followed by methylene chloride/methanol (98/2 and 95/5). After removal of the solvent under vacuum, the residue was triturated with ether, filtered, washed with ether and dried under vacuum to give **7-((1-cyanomethyl)piperidin-4-ylmethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (304 mg, 66 %).

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.6-1.8 (m, 2H), 2.05-2.2 (d, 2H), 2.2-2.3 (m, 1H), 2.45 (s, 3H), 3.2 (t, 2H), 3.65 (d, 2H), 4.1 (s, 3H), 4.22 (d, 2H), 4.6 (s, 2H), 6.2 (s, 0.5H, partially exchanged), 6.9 (dd, 1H), 7.35 (s, 1H), 7.4 (d, 1H), 7.55 (s, 1H), 7.8 (s, 1H), 9.1 (s,

1H)

MS-ESI : 458 [MH]⁺

Elemental analysis

Found C 67.6 H 6.1 N 15.2

C₂₆H₂₇N₅O₃ 0.2 H₂O

Requires C 67.7 H 6.0 N 15.2%

5

Example 152

A mixture of 4-chloro-6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline (360 mg, 1.00 mmol), potassium carbonate (215 mg, 1.56 mmol) and 5-hydroxyindole (147 mg, 1.10 mmol) in DMF (8.0 ml) was stirred at 100 °C for 5 hours and allowed to cool to ambient temperature. The solvent was removed by evaporation and the residue purified by silica column chromatography eluting with methanol (2.5 to 5%) in dichloromethane. The resulting solid was recrystallised from ethyl acetate, filtered and washed with diethyl ether to give 4-(indol-5-yloxy)-6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline (77mg, 17%).

¹H NMR Spectrum: (DMSO-d₆) 2.07 (m, 2H), 2.78 (s, 3H), 2.87 (s, 3H), 3.25 (t, 2H), 3.97 (s, 3H), 4.23 (t, 2H), 6.43 (br s, 1H), 6.96 (dd, 1H), 7.32 (s, 1H), 7.41 (m, 3H), 7.59 (d, 1H), 8.48 (s, 1H) and 11.17 (s, 1H)

MS (ESI) : 457 (MH)⁺

Elemental analysis Found C 57.5 H 5.3 N 12.0

C₂₂H₂₄N₄O₅S Requires C 57.9 H 5.3 N 12.3%

The starting material was prepared as follows:

Using an analogous procedure to that described for the synthesis of the starting material in Example 5, 4-(4-bromo-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline was made in a similar way to 4-(4-chloro-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline using 4-bromo-2-fluorophenol instead of 4-chloro-2-fluorophenol.

A mixture of 4-(4-bromo-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline (9.64g, 26.4 mmol) and triphenylphosphine (20.9g, 79.8 mmol) in dichloromethane (240ml) was stirred under nitrogen, at ambient temperature for 30 minutes. 3-(N-tertButoxycarbonyl)-propanolamine (6.26g, 35.8 mmol) was added followed by diethyl azodicarboxylate (12.4ml, 13.7g, 78.7 mmol). The reaction mixture was stirred for 2 hours. The solvent was then removed by evaporation and the residue taken up in acetonitrile (250ml). The solution was

concentrated to half the original volume and cooled. The resulting crystalline solid was filtered, washed with ether and dried to give 4-(4-bromo-2-fluorophenoxy)-7-(3-(*N*-*tert*butoxycarbonylamino)propoxy)-6-methoxyquinazoline (10.0g, 73%).

¹H NMR Spectrum: (DMSO-d₆) 1.37 (s, 9H), 1.94 (t, 2H), 3.13 (q, 2H), 3.97 (s, 3H), 4.21 (t, 2H), 6.89 (br s, 1H), 7.38 (s, 1H), 7.43 - 7.53 (m, 2H), 7.57 (s, 1H), 7.78 (dd, 1H) and 8.55 (s, 1H)

MS (ESI) : 522 (MH)⁺

Elemental analysis	Found	C 52.1	H 4.7	N 7.9
--------------------	-------	--------	-------	-------

C ₂₃ H ₂₅ N ₃ BrFO ₅	Requires	C 52.3	H 4.9	N 8.0%
--	----------	--------	-------	--------

4-(4-Bromo-2-fluorophenoxy)-7-(3-(*N*-*tert*butoxycarbonylamino)propoxy)-6-methoxyquinazoline (5.46g, 10.5mmol) was taken up in trifluoroacetic acid (75ml) and heated at 85°C for 1.5 hours. The solution was allowed to cool and the excess trifluoroacetic acid removed by evaporation. The residue was then treated with aqueous ammonia (0.88) solution, extracted with dichloromethane (3x150ml) and filtered through phase separating paper. The solvent was removed by evaporation to give 7-(3-aminopropoxy)-4-(4-bromo-2-fluorophenoxy)-6-methoxyquinazoline (4.42g, 100%).

¹H NMR Spectrum: (DMSO-d₆) 1.87 (m, 2H), 2.73 (t, 2H), 3.98 (s, 3H), 4.26 (t, 2H), 7.40 (s, 1H), 7.50 (m, 2H), 7.55 (s, 1H), 7.78 (dd, 1H) and 8.55 (s, 1H)

MS (ESI) : 422 (MH)⁺

A solution of 7-(3-aminopropoxy)-4-(4-bromo-2-fluorophenoxy)-6-methoxyquinazoline (2.71g, 6.4mmol) and triethylamine (1.1ml, 0.80g, 7.9mmol) in dichloromethane (15ml) was treated with a solution of methanesulphonyl chloride (0.53ml, 0.79g, 6.9mmol) in dichloromethane (10ml) and stirred at ambient temperature, under nitrogen for 18 hours. The dichloromethane was then removed by evaporation and THF (4ml) added. The resulting solution was treated with saturated aqueous sodium hydrogen carbonate solution (to pH 8), stirred vigorously for 30 minutes and the precipitate filtered, washed with water and dried to give 4-(4-bromo-2-fluorophenoxy)-6-methoxy-7-(3-(*N*-methylsulphonylamino)propoxy)quinazoline (2.98g, 93%).

¹H NMR Spectrum: (DMSO-d₆) 2.01 (m, 2H), 2.90 (s, 3H), 3.15 (t, 2H), 3.96 (s, 3H), 4.25 (t, 2H), 7.06 (s, 1H), 7.40 (s, 1H), 7.49 (m, 2H), 7.56 (s, 1H), 7.78 (dd, 1H) and 8.54 (s, 1H)

MS (ESI) : 500/502 (MH)⁺

4-(4-Bromo-2-fluorophenoxy)-6-methoxy-7-(3-(*N*-

methysulphonylamino)propoxy)quinazoline (1.0g, 2mmol) was taken up in DMF (10ml), treated with sodium hydride (60% dispersion in mineral oil, 0.11g, 2.7mmol) and stirred, under nitrogen for 30 minutes. Methyl iodide (0.16ml, 2.6mmol) was added and the mixture stirred for 18 hours. The solvent was removed by evaporation and the residue taken up in
5 water and extracted with dichloromethane (3x 30ml). The organic solution was then washed with water, brine, dried (MgSO₄) and evaporated to dryness. The crude product was purified by silica column chromatography eluting with methanol (2.5 to 5 %) in dichloromethane to give 4-(4-bromo-2-fluorophenoxy)-6-methoxy-7-(3-(N-methyl N-methylsulphonylamino)propoxy)quinazoline (0.86g, 83%).

10 ¹H NMR Spectrum: (DMSO-d₆) 2.06 (m, 2H), 2.78 (s, 3H), 2.87 (s, 3H), 3.24 (t, 2H), 3.97 (s, 3H), 4.23 (t, 2H), 7.39 (s, 1H), 7.48 (m, 2H), 7.55 (s, 1H), 7.78 (dd, 1H) and 8.54 (s, 1H)
MS (ESI) : 514/516 (MH)⁺

4-(4-Bromo-2-fluorophenoxy)-6-methoxy-7-(3-(N-methyl N-methylsulphonylamino)propoxy)quinazoline (4.70g, 9.1mmol) was dissolved in 2N aqueous hydrochloric acid
15 solution (85ml) and heated at reflux for 1 hour. After cooling, the solution was carefully poured into saturated aqueous sodium hydrogen carbonate solution (to pH8) and stirred vigorously for 30 minutes. The resulting precipitate was filtered and dried. The filter cake was then taken up as a suspension in acetone, filtered, washed with diethyl ether and dried to give 6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazolin-4-one (3.23g,
20 88%).

¹H NMR Spectrum: (DMSO-d₆) 2.02 (m, 2H), 2.77 (s, 3H), 2.86 (s, 3H), 3.22 (t, 2H), 3.86 (s, 3H), 4.13 (t, 2H), 7.09 (s, 1H), 7.42 (s, 1H), 7.95 (s, 1H) and 12.02 (s, 1H)
MS (ESI) : 342 (MH)⁺

6-Methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazolin-4-one
25 (2.24g, 6.6mmol) was taken up in thionyl chloride (25ml) and treated with DMF (5 drops). The resulting solution was then heated at reflux for 1 hour followed by cooling to ambient temperature. The excess thionyl chloride was removed by evaporation followed by azeotroping with toluene (3x). The residue was basified with saturated aqueous sodium hydrogen carbonate solution (to pH8) and extracted twice with ethyl acetate. The organic
30 solution was washed with water, brine, dried (MgSO₄) and evaporated to dryness to give 4-chloro-6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline (1.90g, 80%).

¹H NMR Spectrum: (DMSO-d₆) 2.08 (m, 2H), 2.78 (s, 3H), 2.88 (s, 3H), 3.24 (t, 2H), 3.98 (s, 3H), 4.26 (t, 2H), 7.37 (s, 1H), 7.42 (s, 1H) and 8.86 (s, 1H)

MS (ESI) : 360(MH)⁺

5 Example 153

A mixture of 4-chloro-6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline (360 mg, 1.00 mmol), (prepared as described for the starting material in Example 152), potassium carbonate (215 mg, 1.56 mmol) and 5-hydroxy-2-methylindole (162 mg, 1.10 mmol) in DMF (8.0 ml) was stirred at 100 °C for 5
10 hours and allowed to cool to ambient temperature. The solvent was removed by evaporation and the residue was purified by silica column chromatography eluting with methanol (2.5 to 5%) in dichloromethane. The resulting solid was recrystallised from ethyl acetate, filtered and washed with diethyl ether to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline** (166mg, 35%).

15 ¹H NMR Spectrum: (DMSO-d₆) 2.06 (m, 2H), 2.38 (s, 3H), 2.79 (s, 3H), 2.89 (s, 3H), 3.24 (t, 2H), 3.96 (s, 3H), 4.21 (t, 2H), 6.11 (br s, 1H), 6.87 (dd, 1H), 7.23 (d, 1H), 7.30 (d, 1H), 7.35 (s, 1H), 7.57 (s, 1H), 8.46 (s, 1H) and 10.98 (s, 1H)

MS (ESI) : 471 (MH)⁺

Elemental analysis Found C 58.3 H 5.6 N 11.7

20 C₂₃H₂₆N₄O₅S Requires C 58.7 H 5.6 N 11.9%

Example 154

A mixture of 4-chloro-6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline (150 mg, 0.42 mmol), (prepared as described for
25 the starting material in Example 152), potassium carbonate (90 mg, 0.63 mmol) and 7-hydroxyquinoline (67 mg, 0.46 mmol) in DMF (5.0 ml) was stirred at 100 °C for 2 hours and allowed to cool to ambient temperature. The solvent was removed by evaporation and the residue taken up in 2N. aqueous sodium hydroxide solution. The precipitate was filtered off, dried, taken up in dichloromethane and the solution filtered through phase separating paper.
30 The filtrate was then evaporated to dryness. The resulting solid was recrystallised from acetonitrile, filtered and washed with diethyl ether to give **6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)-4-(quinolin-7-yloxy)quinazoline** (122mg, 63%).

¹H NMR Spectrum: (DMSO-d₆) 2.09 (m, 2H), 2.79 (s, 3H), 2.90 (s, 3H), 3.26 (t, 2H), 3.99 (s, 3H), 4.26 (t, 2H), 7.39 (s, 1H), 7.54 (dd, 1H), 7.56 (dd, 1H), 7.60 (s, 1H), 7.91 (d, 1H), 8.09 (d, 1H), 8.44 (d, 1H), 8.55 (s, 1H) and 8.93 (dd, 1H)

MS (ESI) : 469 (MH)⁺

5	Elemental analysis	Found	C 58.6	H 5.1	N 11.9
	C ₂₃ H ₂₄ N ₄ O ₅ S	Requires	C 59.0	H 5.2	N 12.0%

Example 155

A mixture of 4-chloro-6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline (150 mg, 0.42 mmol), (prepared as described for the starting material in Example 152), potassium carbonate (90 mg, 0.63 mmol) and 7-hydroxy-4-methylquinoline (71 mg, 0.46 mmol), (Chem. Berich. 1967, 100, 2077), in DMF (5.0 ml) was stirred at 100 °C for 2 hours and allowed to cool to ambient temperature. The DMF solvent was removed by evaporation and the residue was taken up in 2N aqueous sodium hydroxide solution. The precipitate was filtered off, dried, taken up in dichloromethane and then filtered through phase separating paper. The solution was then evaporated to dryness. The resulting solid was recrystallised from acetonitrile, filtered and washed with diethyl ether to give 6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)-4-(4-methylquinolin-7-yloxy)quinazoline (84mg, 42%).

¹H NMR Spectrum: (DMSO-d₆) 2.09 (m, 2H), 2.71 (s, 3H), 2.79 (s, 3H), 2.89 (s, 3H), 3.25 (t, 2H), 3.98 (s, 3H), 4.25 (t, 2H), 7.37 (s, 1H), 7.38 (d, 1H), 7.61 (dd, 1H), 7.63 (s, 1H), 7.89 (d, 1H), 8.20 (d, 1H), 8.54 (s, 1H) and 8.76 (d, 1H)

MS (ESI) : 483 (MH)⁺

	Elemental analysis	Found	C 59.1	H 5.3	N 11.5
25	C ₂₄ H ₂₆ N ₄ O ₅ S	Requires	C 59.1	H 5.0	N 12.0%

Example 156

A mixture of (R,S)-4-chloro-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline (90 mg, 0.28 mmol), (prepared as described for the starting material in Example 7), potassium carbonate (60 mg, 0.44 mmol) and 7-hydroxy-4-trifluoromethylquinoline (65 mg, 0.31 mmol), (prepared as in Ukr. Khim. Zh. (Russ. Ed) Vol. 59, No. 4, pp. 408-411, 1993), in DMF (2 ml) was stirred at 100 °C for 6 hours and then

allowed to cool to ambient temperature. The DMF solvent was removed by evaporation, the residue was taken up in methanol/dichloromethane (1/1) and pre-absorbed onto silica. The crude mixture was purified by silica column chromatography eluting with dichloromethane/methanol/0.880 aqueous ammonia (95/5/1) and the product recrystallised from acetonitrile to give **(*R,S*)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(4-trifluoromethylquinolin-7-yloxy)quinazoline** (58mg, 42%).

¹H NMR Spectrum: (DMSO-d₆, 100°C) 1.24 (m, 1H), 1.59 (m, 1H), 1.70 (m, 1H), 1.83 (m, 1H), 2.05 (m, 2H), 2.17 (m, 1H), 2.24 (s, 3H), 2.64 (dt, 1H), 2.84 (dd, 1H), 4.05 (s, 3H), 4.18 (d, 2H), 7.43 (s, 1H), 7.69 (s, 1H), 7.87 (dd, 1H), 7.96 (d, 1H), 8.18 (s, 1H), 8.25 (dd, 1H), 8.59 (s, 1H) and 9.16 (d, 1H)

MS (ESI) : 499 (MH)⁺

Elemental analysis	Found	C 62.2	H 5.1	N 11.0
--------------------	-------	--------	-------	--------

C ₂₆ H ₂₅ N ₄ F ₃ O ₃	Requires	C 62.6	H 5.1	N 11.2%
--	----------	--------	-------	---------

15 **Example 157**

A mixture of **(*R,S*)-4-chloro-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline** (150 mg, 0.46 mmol), (prepared as described for the starting material in Example 7), potassium carbonate (106 mg, 0.77 mmol) and 3-fluoro-7-hydroxyquinoline (119 mg, 0.73 mmol) in DMF (5 ml) was stirred at 100 °C for 2 hours and then allowed to cool to ambient temperature. The solvent was removed by evaporation and the residue treated with 1.0 N aqueous sodium hydroxide solution (30 ml) then allowed to stir for 30 minutes. The crude solid was collected by filtration and washed with water. The resultant solid was dissolved in dichloromethane and filtered through phase separating paper. The solvent was removed by evaporation and the solid residue was recrystallised from acetonitrile to give **(*R,S*)-4-(3-fluoroquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline** (83mg, 40%).

¹H NMR Spectrum: (DMSO-d₆) 1.11 (m, 1H), 1.50 (m, 1H), 1.64 (m, 1H), 1.84 (m, 3H), 2.10 (m, 1H), 2.15 (s, 3H), 2.62 (d, 1H), 2.83 (d, 1H), 4.00 (s, 3H), 4.08 (d, 2H), 7.38 (s, 1H), 7.62 (s, 1H), 7.68 (dd, 1H), 7.97 (d, 1H), 8.10 (d, 1H), 8.34 (dd, 1H), 8.54 (s, 1H) and 8.97 (d, 1H)

MS (ESI) : 449 (MH)⁺

Elemental analysis	Found	C 66.2	H 5.6	N 12.3
--------------------	-------	--------	-------	--------

C ₂₅ H ₂₅ N ₄ FO ₃ 0.2 H ₂ O	Requires	C 66.4	H 5.7	N 12.4%
---	----------	--------	-------	---------

The starting material, 3-fluoro-7-hydroxyquinoline was prepared as follows:

3-Fluoro-7-methoxyquinol-2(1*H*)-one (300mg, 1.55mmol), (prepared as in Tetrahedron, Vol. 52, No. 9, pp. 3223-3228, 1996), was dissolved in thionyl chloride (3ml),
5 treated with DMF (1 drop) and heated at reflux for 1 hour. The excess thionyl chloride was removed by evaporation and the residue azeotroped with toluene (3x). The residue was basified to pH8 with saturated aqueous sodium hydrogen carbonate solution and extracted with ethyl acetate (3 x 20ml). The organic solution was washed with water and brine then dried (MgSO₄) and evaporated to dryness to give 2-chloro-3-fluoro-7-methoxyquinoline
10 (320mg, 97%).

¹H NMR Spectrum: (CDCl₃) 3.95 (s, 3H), 7.25 (dd, 1H), 7.37 (d, 1H), 7.67 (d, 1H) and 7.78 (d, 1H)

MS (ESI) : 212 (MH)⁺

A mixture of 2-chloro-3-fluoro-7-methoxyquinoline (310mg, 1.47mmol),
15 triethylamine (310mg, 0.4ml, 3.07mmol) and 10% palladium on activated charcoal (50mg) in dry ethanol (5ml) was stirred under hydrogen gas at ambient temperature for 24 hours. The mixture was then filtered through celite. The celite was washed with methanol and the solvent was removed by evaporation from the combined filtrates. The crude material was purified by chromatography on silica, eluting with 10% ethyl acetate in isohexane to give 3-
20 fluoro-7-methoxyquinoline (130mg, 54%).

¹H NMR Spectrum: (CDCl₃) 3.96 (s, 3H), 7.24 (dd, 1H), 7.44 (d, 1H), 7.66 (d, 1H) and 7.73 (dd, 1H) and 8.76 (d, 1H)

MS (ESI) : 178 (MH)⁺

3-Fluoro-7-methoxyquinoline (130mg, 0.74mmol) was taken up in dichloromethane
25 (2ml) under nitrogen and treated with boron tribromide (4ml of a 1.0M solution of in dichloromethane). The reaction mixture was stirred for 24 hours at ambient temperature followed by quenching the reaction by the slow addition of excess methanol. The solution was stirred for a further 2 hours and evaporated to dryness to give 3-fluoro-7-hydroxyquinoline which was used without further purification.

30 MS (ESI) : 164 (MH)⁺

Example 158

A mixture of (*R,S*)-4-chloro-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline (240 mg, 0.75 mmol), (prepared as described for the starting material in Example 7), potassium carbonate (160 mg, 1.16 mmol) and 3-fluoro-7-hydroxy-2-methylquinoline (150 mg, 0.85 mmol) in DMF (6 ml) was stirred at 100 °C for 5 hours and then allowed to cool to ambient temperature. The solvent was removed by evaporation, then the residue was treated with water and 1.0 N aqueous sodium hydroxide solution (30 ml) then allowed to stir for 30 minutes. The crude solid was collected by filtration and washed with water. The resulting solid was dissolved in dichloromethane and filtered through phase separating paper. The solvent was removed by evaporation to give a solid residue which was recrystallised from acetonitrile to give 4-(3-fluoro-2-methylquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline (71mg, 21%).

¹H NMR Spectrum: (DMSO-d₆) 1.11 (m, 1H), 1.68 (m, 5H), 2.10 (m, 1H), 2.20 (s, 3H), 2.64 (m, 4H), 2.87 (d, 1H), 3.98 (s, 3H), 4.09 (d, 2H), 7.37 (s, 1H), 7.57 (dd, 1H), 7.60 (s, 1H), 7.86 (d, 1H), 8.02 (d, 1H), 8.20 (d, 1H) and 8.53 (s, 1H)

MS (ESI) : 463 (MH)⁺

Elemental analysis	Found	C 66.4	H 6.1	N 11.8
C ₂₆ H ₂₇ N ₄ FO ₃ 0.4 H ₂ O	Requires	C 66.5	H 6.0	N 11.9%

The starting material was prepared as follows:

2-Chloro-3-fluoro-7-methoxyquinoline (210g, 1mmol), (prepared as described for the starting material in Example 157), in anhydrous THF (1ml) was added to a mixture of copper(I)bromide (570mg, 4.0mmol) and methylmagnesium bromide (3.0M solution in diethyl ether, 2.7ml, 8mmol) in anhydrous THF (20ml) at -78°C. The mixture was stirred for 1 hour at -78°C, allowed to warm to ambient temperature and then stirred for a further 18 hours. Saturated aqueous ammonium chloride solution and 5N aqueous sodium hydroxide solution (pH 12) were added and the product extracted with ethyl acetate (3x). The organic solution was washed with water, brine, dried (MgSO₄) and evaporated to dryness to yield 3-fluoro-7-methoxy-2-methylquinoline (0.17g, 91%).

¹H NMR Spectrum: (CDCl₃) 2.70 (d, 3H), 3.94 (s, 3H), 7.17 (dd, 1H), 7.37 (d, 1H) and 7.61 (m, 2H)

MS (ESI) : 192 (MH)⁺

3-Fluoro-7-methoxy-2-methylquinoline (0.16g, 0.85mmol) was taken up in

- 190 -

dichloromethane (4ml) under nitrogen and treated with boron tribromide solution (4ml of a 1.0M solution in dichloromethane, 4.0mmol). The reaction was stirred for 24 hours at ambient temperature followed by the slow addition of excess methanol. The solution was stirred for a further 2 hours and then evaporated to dryness to give 3-fluoro-7-hydroxy-2-methylquinoline which was used without further purification.

MS (ESI) : 178 (MH)⁺

Example 159

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (400mg, 1.19mmol), (prepared as described for the starting material in Example 67), potassium carbonate (255 mg, 1.84 mmol) and 7-hydroxyquinoline (180mg, 1.32 mmol) in DMF (10 ml) was stirred at 100 °C for 4 hours and then allowed to cool to ambient temperature. The resulting mixture was treated with 1.0 N aqueous sodium hydroxide solution (30 ml) and allowed to stir for 1 hour. The crude solid was collected by filtration and washed with water.

The resulting solid was dissolved in dichloromethane and filtered through phase separating paper. The solvent was removed by evaporation to give a solid residue which was recrystallised from acetonitrile to give **6-methoxy-7-(3-piperidinopropoxy)-4-(quinolin-7-yloxy)quinazoline** (0.27 g, 52%).

¹H NMR Spectrum: (DMSO-d₆) 1.37 (m, 2H), 1.51 (m, 4H), 1.95 (m, 2H), 2.32 (m, 4H), 2.42 (t, 2H), 3.98 (s, 3H), 4.23 (t, 2H), 7.38 (s, 1H), 7.56 (m, 2H), 7.62 (s, 1H), 7.91 (d, 1H), 8.09 (d, 1H), 8.44 (d, 1H), 8.54 (s, 1H) and 8.91 (dd, 1H)

MS (ESI) : 445 (MH)⁺

Elemental analysis	Found	C 70.9	H 6.3	N 12.7
--------------------	-------	--------	-------	--------

C ₂₆ H ₂₈ N ₄ O ₃	Requires	C 70.3	H 6.3	N 12.6%
---	----------	--------	-------	---------

Example 160

A mixture of 4-chloro-6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline (360 mg, 1.00 mmol), (prepared as described for the starting material in Example 152), potassium carbonate (215 mg, 1.56 mmol) and 2,3-dimethyl-5-hydroxyindole (177 mg, 1.10 mmol), (Arch. Pharm. 1972, 305, 159), in DMF (8.0 ml) was stirred at 100 °C for 5 hours and allowed to cool to ambient temperature. The solvent was removed by evaporation and the residue purified by silica column chromatography

- 191 -

eluting with methanol (2.5%) in dichloromethane. The resulting solid was recrystallised from *tert*butyl methyl ether/acetonitrile, filtered and washed with diethyl ether to give 4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(3-(*N*-methyl-*N*-methylsulphonylamino)propoxy)quinazoline (201mg, 42%).

5 ¹H NMR Spectrum: (DMSO-d₆) 2.07 (m, 2H), 2.12 (s, 3H), 2.31 (s, 3H), 2.79 (s, 3H), 2.89 (s, 3H), 3.25 (t, 2H), 3.97 (s, 3H), 4.23 (t, 2H), 6.86 (dd, 1H), 7.20 (d, 1H), 7.25 (d, 1H), 7.35 (s, 1H), 7.58 (s, 1H), 8.46 (s, 1H) and 11.17 (s, 1H)

MS (ESI) : 485 (MH)⁺

Elemental analysis	Found	C 59.5	H 5.8	N 11.4
--------------------	-------	--------	-------	--------

10	C ₂₄ H ₂₈ N ₄ O ₅ S	Requires	C 59.5	H 5.8	N 11.6%
----	---	----------	--------	-------	---------

Example 161

A mixture of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (322 mg, 1.00 mmol), (prepared as described in Example 49), potassium carbonate (414 mg, 3.00 mmol) and epibromohydrin (274 mg, 2.00 mmol) in DMF (7.0 ml) was stirred at 60 °C for 2 hours and allowed to cool to ambient temperature. The solvent was removed by evaporation and the residue taken up in dichloromethane (10ml). An aliquot (5ml) of this solution was treated with morpholine (48ul, 0.6 mmol) and stirred for 24 hours at ambient temperature. The solvent was removed by evaporation, treated with water and stirred vigorously for 30 minutes.

20 The precipitate was filtered, washed with water and dried. The resultant solid was stirred as a suspension in acetone, filtered, washed with diethyl ether and dried to give 7-(2-hydroxy-3-morpholinopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (127mg, 27%).

¹H NMR Spectrum: (DMSO-d₆) 2.38 (s, 3H), 2.45 (m, 6H), 3.57 (t, 4H), 3.95 (s, 3H), 4.03 - 4.14 (m, 2H), 4.23 (m, 1H), 4.95 (s, 1H), 6.12 (s, 1H), 6.86 (dd, 1H), 7.23 (d, 1H), 7.29 (d, 1H), 7.37 (s, 1H), 7.57 (s, 1H), 8.47 (s, 1H) and 10.98 (s, 1H)

MS (ESI) : 465 (MH)⁺

Elemental analysis	Found	C 62.7	H 5.9	N 11.5
--------------------	-------	--------	-------	--------

C ₂₅ H ₂₈ N ₄ O ₅ ·0.7H ₂ O	Requires	C 62.9	H 6.2	N 11.7%
--	----------	--------	-------	---------

30 Example 162

A mixture of 7-(2,3-epoxypropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (100mg, 0.27 mmol) and piperidine (79ul, 0.8mmol) in DMF (4ml) was heated at 70°C for 24

hours. The solvent was removed by evaporation and the residue was recrystallised from acetonitrile. The solid was filtered, washed with diethyl ether and dried to give 7-(2-hydroxy-3-piperidinopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (80mg, 65%).

- 5 ¹H NMR Spectrum: (DMSO-d₆) 1.35 (m, 2H), 1.51 (m, 4H), 2.39 (m, 9H), 3.96 (s, 3H), 4.08 (m, 2H), 4.21 (dd, 1H), 4.86 (br s, 1H), 6.11 (s, 1H), 6.87 (dd, 1H), 7.23 (d, 1H), 7.29 (d, 1H), 7.37 (s, 1H), 7.56 (s, 1H), 8.45 (s, 1H) and 10.98 (s, 1H)

MS (ESI) : 464 (MH)⁺

Elemental analysis Found C 66.2 H 6.4 N 11.9

- 10 C₂₆H₃₀N₄O₄·0.4H₂O Requires C 66.5 H 6.6 N 11.9%

The starting material was prepared as follows:

- A mixture of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (1.89 g, 5.90 mmol), (prepared as described in Example 49), potassium carbonate (2.43 g, 17.6 mmol) and epibromohydrin (1.61 g, 11.7 mmol) in DMF (40 ml) was stirred at 60 °C for 2 hours and allowed to cool to ambient temperature. The insoluble inorganic material was removed by filtration and the solvent was removed by evaporation. The residue was triturated with diethyl ether, filtered, washed with further diethyl ether and dried to give 7-(2,3-epoxypropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (1.97g, 89%).

- 20 ¹H NMR Spectrum: (DMSO-d₆) 2.38 (s, 3H), 2.76 (m, 1H), 2.90 (t, 1H), 3.43 (m, 1H), 3.97 (s, 3H), 4.04 (m, 1H), 4.57 (dd, 1H), 6.11 (s, 1H), 6.86 (dd, 1H), 7.27 (m, 2H), 7.38 (s, 1H), 7.59 (s, 1H), 8.46 (s, 1H) and 10.92 (s, 1H)

MS (ESI) : 378 (MH)⁺

25 Example 163

- A mixture of 7-(2,3-epoxypropoxy)-6-methoxy-4-(2-methylindol-5-yloxy) quinazoline (100mg, 0.27 mmol), (prepared as described for the starting material in Example 162), and pyrrolidine (67ul, 0.8mmol) in DMF (4ml) was heated at 70°C for 24 hours. The solvent was removed by evaporation and the residue purified by silica column chromatography eluting with dichloromethane/methanol/0.880 aqueous ammonia (100/8/1). The relevant fractions were evaporated to dryness then the residue treated with a little dichloromethane and dried under high vacuum to give 7-(2-hydroxy-3-pyrrolidin-1-ylpropoxy)-6-methoxy-4-(2-

methyldol-5-yloxy)quinazoline (44mg, 37%) as a white foam.

¹H NMR Spectrum: (DMSO-d₆) 1.69 (br s, 4H), 2.38 (s, 3H), 2.50 (m, 6H), 3.97 (s, 3H), 4.07 (m, 2H), 4.21 (dd, 1H), 4.96 (br s, 1H), 6.11 (s, 1H), 6.86 (dd, 1H), 7.23 (d, 1H), 7.29 (d, 1H), 7.35 (s, 1H), 7.56 (s, 1H), 8.46 (s, 1H) and 10.98 (s, 1H)

5 MS (ESI) : 450 (MH)⁺

Elemental analysis	Found	C 65.5	H 6.3	N 11.8
--------------------	-------	--------	-------	--------

C ₂₅ H ₂₈ N ₄ O ₄ ·0.4H ₂ O	Requires	C 65.9	H 6.4	N 12.3%
--	----------	--------	-------	---------

Example 164

10 A mixture of 7-(2,3-epoxypropoxy)-6-methoxy-4-(2-methyldol-5-yloxy) quinazoline (100mg, 0.27 mmol), (prepared as described for the starting material in Example 162), and diethylamine (100ul, 0.8mmol) in DMF (4ml) was heated at 70°C for 24 hours. The solvent was removed by evaporation and the residue was purified by silica column chromatography eluting with dichloromethane/methanol/0.880 aqueous ammonia (100/8/1). The relevant
15 fractions were evaporated to dryness then the residue treated with a little dichloromethane and dried under high vacuum to give **7-(3-(N,N-diethylamino)-2-hydroxypropoxy)-6-methoxy-4-(2-methyldol-5-yloxy)quinazoline** (55mg, 46%) as a white foam.

¹H NMR Spectrum: (DMSO-d₆) 0.96 (t, 6H), 2.38 (s, 3H), 2.52 (m, 6H), 3.96 (s, 3H), 3.97 (m, 1H), 4.09 (m, 1H), 4.23 (dd, 1H), 4.84 (br s, 1H), 6.12 (s, 1H), 6.88 (dd, 1H), 7.24 (d, 1H),
20 7.29 (d, 1H), 7.36 (s, 1H), 7.56 (s, 1H), 8.45 (s, 1H) and 10.98 (s, 1H)

MS (ESI) : 452 (MH)⁺

Elemental analysis	Found	C 66.2	H 6.7	N 12.4
--------------------	-------	--------	-------	--------

C ₂₅ H ₃₀ N ₄ O ₄	Requires	C 66.6	H 6.7	N 12.4%
---	----------	--------	-------	---------

25 Example 165

A mixture of 7-(2,3-epoxypropoxy)-6-methoxy-4-(2-methyldol-5-yloxy) quinazoline (100mg, 0.27 mmol), (prepared as described for the starting material in Example 162), and N-methylpiperazine (200ul, 1.8mmol) in DMF (4ml) was heated at 70°C for 24 hours. The solvent was removed by evaporation and the residue was recrystallised from acetonitrile. The
30 solid was filtered, washed with diethyl ether and dried to give **7-(2-hydroxy-3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-4-(2-methyldol-5-yloxy)quinazoline** (41mg, 32%).

- 194 -

¹H NMR Spectrum: (DMSO-d₆) : 2.11 (s, 3H), 2.29 (m, 4H), 2.40 (s, 3H), 2.47 (m, 6H), 3.96 (s, 3H), 4.07 (m, 2H), 4.20 (dd, 1H), 4.89 (d, 1H), 6.11 (s, 1H), 6.87 (dd, 1H), 7.23 (d, 1H), 7.29 (d, 1H), 7.35 (s, 1H), 7.58 (s, 1H), 8.46 (s, 1H) and 10.98 (s, 1H)

MS (ESI) : 479 (MH)⁺

5	Elemental analysis	Found	C 64.4	H 6.5	N 14.4
	C ₂₆ H ₃₁ N ₅ O ₄ ·0.3H ₂ O	Requires	C 64.7	H 6.6	N 14.5%

Example 166

A mixture of 7-(2,3-epoxypropoxy)-6-methoxy-4-(2-methylindol-5-yloxy) quinazoline (100mg, 0.27 mmol), (prepared as described for the starting material in Example 162), and isopropylamine (100ul, 0.8mmol) in DMF (4ml) was heated at 70°C for 24 hours. The solvent was removed by evaporation and the residue was purified by silica column chromatography eluting with dichloromethane/methanol/0.880 aqueous ammonia (100/8/1) to give 7-(2-hydroxy-3-(isopropylamino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (18mg, 16%).

¹H NMR Spectrum: (DMSO-d₆) 1.00 (d, 6H), 2.40 (s, 3H), 2.56 - 2.78 (m, 3H), 3.97 (m, 4H), 4.07 - 4.28 (m, 2H), 5.04 (m, 1H), 6.12 (s, 1H), 6.88 (dd, 1H), 7.22 - 7.33 (m, 2H), 7.38 (s, 1H), 7.58 (s, 1H), 8.48 (s, 1H) and 10.98 (s, 1H)

MS (ESI) : 437 (MH)⁺

Example 167

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (168 mg, 0.5 mmol), (prepared as described for the starting material in Example 67), potassium carbonate (276 mg, 2.0 mmol) and 5-hydroxy-6-trifluoromethylindole (110 mg, 0.55 mmol) and DMA (4.0 ml) were stirred at 95°C for 1.5 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated under vacuum. The residue was purified by silica column chromatography eluting with dichloromethane/methanol/0.880 aqueous ammonia (89/10/1) to give a partially purified oil. This oil was further purified by high performance column chromatography on octadecylsilane reverse phase silica eluting with acetonitrile/water/trifluoroacetic acid (60/39.8/0.2) to give an oil which was dissolved in dichloromethane and washed with saturated aqueous sodium hydrogen carbonate solution. The dichloromethane layer was evaporated to give 6-methoxy-7-(3-piperidinopropoxy)-4-

- 195 -

(6-trifluoromethylindol-5-yloxy)quinazoline (62 mg, 25%).

¹H NMR Spectrum: (DMSO-d₆) 1.45 (m, 2H), 1.60 (m, 4H), 2.13 (m, 2H), 2.44 (m, 4H), 2.56 (m, 2H), 4.04 (s, 3H), 4.27 (t, 2H), 6.63 (br s, 1H), 7.33 (s, 1H), 7.40 (t, 1H), 7.61 (s, 1H), 7.67 (s, 1H), 7.75 (s, 1H) and 8.60 (m, 2H)

5 MS (ESI) : 501 (MH)⁺

Elemental analysis	Found	C 62.0 H 5.6 N 10.6
--------------------	-------	---------------------

C ₂₆ H ₂₇ F ₃ N ₄ O ₃ 0.35 H ₂ O,	Requires	C 61.6 H 5.5 N 11.0%
---	----------	----------------------

The starting material was prepared as follows:

10 Sodium hydride (1.8g, of a 60% dispersion in oil, 45 mmol) was added in portions to a stirred solution of benzyl alcohol (10.8g, 100 mmol) in DMA (100ml) with vigorous stirring under an atmosphere of nitrogen at ambient temperature. After warming to 45°C for 30 minutes the mixture was cooled to ambient temperature and added dropwise to a stirred solution of 2-chloro-5-nitro-trifluoromethylbenzene (11.3g, 50 mmol) in DMA (30ml),
15 keeping the temperature below 10°C. The mixture was stirred at 25°C for 1 hour, then acidified with acetic acid and evaporated to give a yellow solid. The residue was dissolved in dichloromethane, washed with water then dried (MgSO₄), and evaporated. The residue was suspended in a mixture of hexane (70 ml) and diethyl ether (10 ml) and the resulting solid filtered off to give 2-benzyloxy-5-nitro-trifluoromethylbenzene (6.6g, 49%).

20 ¹H NMR Spectrum: (CDCl₃) 5.33 (s, 2H), 7.13 (d, 1H), 7.31-7.43 (m, 5H), 8.35 (dd, 1H), 8.52 (d, 1H)

Potassium *tert*-butoxide (3.94g, 35.4mmol) was dissolved in anhydrous DMF (15ml) and a mixture of 2-benzyloxy-5-nitro-trifluoromethylbenzene (3.5g, 16.1 mmol) and 4-chlorophenylacetonitrile (2.96g, 17.7 mmol) in DMF (20 ml) was added over 30 minutes
25 keeping the temperature at -15°C. The mixture was stirred at -10°C for 1 hour, then poured into 1M hydrochloric acid (150ml) and the product extracted with dichloromethane (2x100ml). The organic extracts were dried (MgSO₄) and purified by silica column chromatography eluting with dichloromethane/hexane (1/1) to give 5-benzyloxy-2-nitro-4-(trifluoromethyl)phenylacetonitrile (5.2g, 77%).

30 ¹H NMR Spectrum: (CDCl₃) 4.30 (s, 2H), 5.38 (s, 2H), 7.25 (s, 1H), 7.33-7.50 (m, 5H) and 8.51 (s, 1H)

MS (ESI) : 335 (M-H)⁻

5-Benzyloxy-2-nitro-4-(trifluoromethyl)phenylacetonitrile (2.22g, 6.6mmol) was dissolved in ethanol (45 ml), water (5ml) and acetic acid (0.32 ml) then hydrogenated with 10% palladium on carbon at 1 atmosphere pressure for 2 hours. The catalyst was filtered off and filtrate evaporated to give 5-hydroxy-6-trifluoromethylindole (1.12g, 84%).

5 ¹H NMR Spectrum: (CDCl₃) 4.48 (s, 1H), 6.48 (m, 1H), 7.14 (s, 1H), 7.32 (t, 1H), 7.57 (s, 1H) and 8.20 (br s, 1H)

MS (ESI) : 200 (M-H)⁻

Example 168

10 A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (200mg, 0.6 mmol), (prepared as described for the starting material in Example 67), potassium carbonate (248 mg, 1.8 mmol) and 5-hydroxy-6-methoxyindole (127 mg, 0.78 mmol) in DMA (4.0 ml) was stirred at 95°C for 2.5 hours. The reaction mixture was allowed to cool to ambient temperature, filtered and the filtrate evaporated under vacuum. The residue was purified by
15 silica column chromatography eluting with dichloromethane /methanol/0.880 aqueous ammonia (89/10/1) and the resulting oil triturated with diethyl ether to give 4-(6-methoxyindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline (106 mg, 38%).

¹H NMR Spectrum: (DMSO-d₆) 1.38 (m, 2H), 1.47 (m, 4H), 1.95 (m, 2H), 2.32 (m, 4H), 2.40 (m, 2H), 3.66 (3H, s), 3.97 (s, 3H), 4.28 (t, 2H), 6.35 (br s, 1H), 7.06 (s, 1H), 7.24 (t, 1H),
20 7.34 (s, 1H), 7.36 (s, 1H), 7.55 (s, 1H) and 8.41 (s, 1H)

MS (ESI) : 463 (MH)⁺

Elemental analysis	Found	C 65.2 H 6.8 N 11.2
C ₂₆ H ₃₀ N ₄ O ₄ 1.0 H ₂ O, 0.3 diethyl ether	Requires	C 64.9 H 7.0 N 11.1%

25 The 5-hydroxy-6-methoxyindole starting material was made as follows:

5-Benzyloxy-6-methoxyindole (253mg, 1.0mmol) was hydrogenated at 1 atmosphere pressure in methanol (10 ml) with 10% palladium on carbon (50 mg) for 2 hours at 25°C. The catalyst was filtered off and the filtrate evaporated to give 5-hydroxy-6-methoxyindole (141mg, 87%).

30 ¹H NMR Spectrum: (CDCl₃) 3.92 (s, 3H), 5.40 (s, 1H), 6.42 (br s, 1H), 6.87 (s, 1H), 7.07 (m, 1H), 7.13 (s, 1H), 7.93 (br s, 1H)

MS (ESI) : 162 (M-H)⁻

Example 169

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (200mg, 0.595 mmol), (prepared as described for the starting material in Example 67), potassium carbonate (411 mg, 2.98 mmol) and 4-hydroxyindole (103 mg, 0.774 mmol) in DMA (2.0 ml) was stirred at 85 °C for 3 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated to give a solid residue. The residue was purified by silica column chromatography, with gradient elution using dichloromethane with 0%, 2%, 4%, 10% methanolic ammonia to give **4-(indol-4-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline** (131 mg, 51 %).

¹H NMR Spectrum: (DMSO-d₆) 1.39 (m, 2H), 1.50 (m, 4H), 1.98 (t, 2H), 2.35 (m, 4H), 2.40 (t, 2H), 3.98 (s, 3H), 4.25 (t, 2H), 6.10 (t, 1H), 6.90 (d, 1H), 7.15 (t, 1H), 7.30 (t, 1H), 7.35 (d, 1H), 7.38 (s, 1H), 7.62 (s, 1H), 8.45 (s, 1H) and 11.29 (s, 1H)

MS (ESI) : 433 (MH)⁺

m.p. 80 - 82 °C

Example 170

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (200mg, 0.595 mmol), (prepared as described for the starting material in Example 67), potassium carbonate (411 mg, 2.98 mmol) and 3-hydroxycarbazole (142 mg, 0.774 mmol) in DMA (2.0 ml) was stirred at 85 °C for 3 hours then allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated to give a solid residue. The residue was purified by silica column chromatography with gradient elution using dichloromethane with 0%, 2%, 4%, 10% methanolic ammonia to give **4-(9H-carbazol-3-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline** (212 mg, 74 %).

¹H NMR Spectrum: (DMSO-d₆) 1.39 (m, 2H), 1.50 (m, 4H), 2.35 (m, 4H), 2.40 (t, 2H), 3.98 (s, 3H), 4.25 (t, 2H), 7.05 (dd, 1H), 7.15 (t, 1H), 7.35 (t, 1H), 7.38 (s, 1H), 7.40 (s, 1H), 7.50 (d, 1H), 7.60 (s, 1H), 8.10 (d, 1H), 8.15 (d, 1H), 8.55 (s, 1H) and 11.33 (s, 1H)

MS (ESI) : 483 (MH)⁺

Example 171

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (84 mg, 0.24 mmol), (prepared as described for the starting material in Example 67), potassium carbonate (162 mg, 1.18 mmol) and ethyl 7-chloro-5-hydroxyindole-2-carboxylate (62 mg, 0.26 mmol) in DMA (2.0 ml) was stirred at 100 °C for 2 hours and allowed to cool to ambient

temperature. The reaction mixture was filtered and the filtrate evaporated. The residue was purified by silica column chromatography using gradient elution dichloromethane with 2.5%, 5%, 10% methanol, then dichloromethane with 2% ammonia) to give **4-(7-chloro-2-(ethoxycarbonyl)indol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline** (78 mg, 63 %).

¹H NMR Spectrum: (DMSO-d₆) 1.30 (t, 3H), 1.40 (m, 2H), 1.50 (m, 4H), 1.98 (t, 2H), 2.35 (m, 4H), 2.40 (t, 2H), 3.98 (s, 3H), 4.25 (t, 2H), 4.30 (q, 2H), 7.15 (m, 1H), 7.18 (s, 1H), 7.60 (s, 1H), 8.40 (s, 1H) and 12.60 (s, 1H)

MS (ESI) : 539 (MH)⁺

Elemental analysis	Found	C 61.2	H 5.9	N 10.3
--------------------	-------	--------	-------	--------

C ₂₈ H ₃₁ ClN ₄ O ₅ · 0.5 H ₂ O	Requires	C 61.4	H 5.9	N 10.2%
--	----------	--------	-------	---------

The starting material was prepared as follows:

2-Chloro-4-methoxyaniline (2.719g, 14 mmol) was added to 8.0M aqueous hydrochloric acid (15 ml) and the suspension cooled to -5 °C. Sodium nitrite (1.063g, 15.4 mmol) was added as a solution in water (3 ml). After addition the pH was brought to pH 4-5 by addition of sodium acetate. In a separate flask, ethyl-α-ethyl acetoacetate (2.18 ml, 15.4 mmol) in ethanol (15 ml) at -5 °C was treated with potassium hydroxide (864 mg, 15.4 mmol) in water (3 ml) followed by ice (4 g). The diazonium salt prepared initially was then added rapidly to the second solution and stirred at -5 °C for 4 hours then allowed to warm to ambient

temperature overnight. The mixture was extracted with ethyl acetate (3 x 100 ml) and the organic solutions dried (MgSO₄), filtered and solvent removed *in vacuo* to give an orange oil. This oil was dissolved in ethanol (35 ml) and the flask fitted with a reflux condenser.

Concentrated sulphuric acid (35 ml) was then added dropwise, this caused the reaction to reflux with no external heating. The solution was stirred for 1 hour then the solvent removed by evaporation. The residue was taken up in water then extracted with ethyl acetate (3 x 100 ml). The organic solution was washed with brine, dried (MgSO₄), filtered and evaporated to give a brown oil. The crude oil was purified by silica column chromatography, eluting with

dichloromethane to give ethyl 7-chloro-5-methoxyindole-2-carboxylate (125 mg, 4%).

¹H NMR Spectrum: (CDCl₃) 1.40 (t, 3H), 3.98 (s, 3H), 4.40 (q, 2H), 6.60 (d, 1H), 7.05 (d, 1H), 7.15 (s, 1H) and 9.10 (s, 1H)

MS (ESI) : 254 (MH)⁺

5 To a solution of ethyl 7-chloro-5-methoxyindole-2-carboxylate (82 mg, 0.323 mmol) in dichloromethane (5 ml) at -78 °C was added boron tribromide (1.07 ml of a 1.0M solution in DCM, 1.07 mmol) and the reaction stirred at -78 °C for 30 minutes then allowed to warm to ambient temperature overnight. Water was carefully added and the pH adjusted to pH 6-7 by addition of 2M sodium hydroxide. The mixture was extracted with ethyl acetate (2 x 50
10 ml), and the organic solution washed with brine, dried (MgSO₄), filtered and evaporated to give ethyl 7-chloro-5-hydroxyindole-2-carboxylate (55 mg, 71%) as an orange solid.

¹H NMR Spectrum: (DMSO-d₆) 1.38 (t, 3H), 4.35 (q, 2H), 6.60 (d, 1H), 6.95 (d, 1H), 7.10 (d, 1H), 9.80 (s, 1H) and 11.80 (s, 1H)

MS (ESI) : 238 (MH)⁻

15

Example 172

A mixture of 7-benzyloxy-4-chloro-6-methoxyquinazoline (1.5 g, 4.99 mmol), (prepared as described for the starting material in Example 1), potassium carbonate (2.07 g, 15 mmol) and 2,3-dimethyl-5-hydroxyindole (1.21 g, 7.5 mmol), (Arch. Pharm. 1972, 305, 159),
20 in DMF (75 ml) was stirred at 100 °C for 2 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated. The solid residue was purified by silica column chromatography, eluting with 2.5% methanol in dichloromethane to give 7-**benzyloxy-4-(2,3-dimethylindol-5-yloxy)-6-methoxyquinazoline** (976 mg, 46 %).

¹H NMR Spectrum: (CDCl₃) 2.10 (s, 3H), 2.30 (s, 3H), 3.98 (s, 3H), 5.30 (s, 2H), 6.85 (dd, 1H), 7.20 (d, 1H), 7.25 (d, 1H), 7.40 (m, 6H), 7.60 (s, 1H), 8.40 (s, 1H) and 10.74 (s, 1H)

25 MS (ESI) : 426 (MH)⁺

Example 173

A mixture of 7-benzyloxy-4-(2,3-dimethylindol-5-yloxy)-6-methoxyquinazoline (912 mg, 2.14 mmol), (prepared as described in Example 172), di-*tert*-butyl dicarbonate (1.871 g, 8.56 mmol) and 4-dimethylaminopyridine (70 mg, 5 mol%) in acetonitrile (40 ml) was stirred at ambient temperature overnight. The solvent was then evaporated and the residue dissolved

- 200 -

in ethyl acetate. The organic solution was washed with 2N hydrochloric acid twice and then with brine. The organic layer was then dried (MgSO₄), filtered and evaporated to give **7-benzyloxy-4-(1-*tert*-butoxycarbonyl-2,3-dimethylindol-5-yloxy)-6-methoxyquinazoline** (1.108 g, 99%) as a yellow solid.

5 ¹H NMR Spectrum: (CDCl₃) 1.70 (s, 9H), 2.08 (s, 3H), 2.50 (s, 3H), 4.10 (s, 3H), 5.35 (s, 2H), 7.15 (dd, 1H), 7.38 (m, 6H), 7.60 (s, 1H), 8.20 (d, 1H) and 8.60 (s, 1H)
MS (ESI) : 526 (MH)⁺

Example 174

10 A mixture of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (225 mg, 0.67 mmol), (prepared as described for the starting material in Example 1), potassium carbonate (106 mg, 0.77 mmol) and 2-hydroxyquinoline (111 mg, 0.76 mmol) in DMF (7.5 ml) was stirred at 100 °C for 5 hours and allowed to cool to ambient temperature. The reaction mixture was treated with 1.0 N aqueous sodium hydroxide solution (40 ml) and
15 allowed to stir at ambient temperature for a few minutes. The reaction mixture was extracted 3 times with ethyl acetate and the extracts washed with water and brine. The organic extracts were dried over magnesium sulphate, filtered and the solvent removed by evaporation. The residue was purified by silica column chromatography eluting with dichloromethane/methanol (95/5) to give a solid which was triturated with ether, filtered and dried to give **6-methoxy-7-(3-morpholinopropoxy)-4-(quinolin-2-yloxy)-quinazoline** (33 mg, 11%).
20

¹H NMR Spectrum: (DMSO-d₆) 1.98 (m, 2H), 2.38 (m, 4H), 2.48 (t, 2H), 3.58 (m, 4H), 3.98 (s, 3H), 4.26 (t, 2H), 7.41 (s, 1H), 7.52 (d, 1H), 7.58 (s, 1H), 7.64 (t, 1H), 7.78 (m, 1H), 7.88 (d, 1H), 8.06 (d, 1H), 8.56 (d, 1H) and 8.57 (s, 1H)

MS (ESI) : 447 (MH)⁺

25	Elemental analysis	Found	C 66.8	H 5.9	N 12.4
	C ₂₅ H ₂₆ N ₄ O ₄ 0.2 H ₂ O	Requires	C 66.7	H 5.9	N 12.4%

Example 175

30 A mixture of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (225 mg, 0.67 mmol), (prepared as described for the starting material in Example 1), potassium carbonate (106 mg, 0.77 mmol) and 5-hydroxyquinoline (111 mg, 0.77 mmol) in DMF (7.5 ml) was stirred at 100 °C for 5 hours and allowed to cool to ambient temperature. The reaction

- 201 -

mixture was treated with 1.0 N aqueous sodium hydroxide solution (40 ml) and allowed to stir at ambient temperature for a few minutes. The resulting precipitate was filtered off, washed with water and air dried for a short while. The damp solid was dissolved in dichloromethane, filtered through phase separating paper and the filtrate evaporated under vacuum. The residue
5 was triturated with ether, filtered and dried to give **6-methoxy-7-(3-morpholinopropoxy)-4-(quinolin-5-yloxy)-quinazoline** (178 mg, 59%).

¹H NMR Spectrum: (DMSO-d₆) 1.98 (m, 2H), 2.39 (m, 4H), 2.48 (t, 2H), 3.59 (t, 4H), 4.01 (s, 3H), 4.28 (t, 2H), 7.42, (s, 1H), 7.50 (m, 1H), 7.59 (d, 1H), 7.74 (s, 1H), 7.87 (t, 1H), 8.02 (d, 1H), 8.20 (m, 1H), 8.44 (s, 1H) and 8.96 (m, 1H)

10 MS (ESI) : 447 (MH)⁺

Elemental analysis	Found	C 66.2	H 5.7	N 12.4
C ₂₅ H ₂₆ N ₄ O ₄ 0.4 H ₂ O	Requires	C 66.2	H 6.0	N 12.4%

Example 176

A mixture of 4-chloro-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (200 mg, 0.57 mmol), potassium carbonate (106 mg, 0.77 mmol) and 7-hydroxyquinoline (111 mg, 0.76 mmol) in DMF (7 ml) was stirred at 100 °C for 5 hours and allowed to cool to ambient temperature. The reaction mixture was treated with 1.0 N aqueous sodium hydroxide solution (40 ml) and allowed to stir at ambient temperature for a few minutes. The reaction mixture was extracted 4 times with ethyl acetate and the organic extracts washed with water and brine. The organic extracts were dried over magnesium sulphate, filtered and the solvent removed by evaporation. The residue was triturated with ether/isohexane, filtered and dried to give **6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)-4-(quinolin-7-yloxy)quinazoline** (102 mg, 39%).

¹H NMR Spectrum: (DMSO-d₆) 1.96 (m, 2H), 2.15 (s, 3H), 2.35 (m, 8H), 2.46 (t, 2H), 3.99 (s, 3H), 4.24 (t, 2H), 7.39 (s, 1H), 7.56 (m, 1H), 7.61 (m, 1H), 7.62 (s, 1H), 7.92 (d, 1H), 8.10 (d, 1H), 8.44 (d, 1H), 8.54 (s, 1H) and 8.92 (m, 1H)

MS (ESI) : 460 (MH)⁺

Elemental analysis	Found	C	67.2	H	6.2	N	15.0
--------------------	-------	---	------	---	-----	---	------

C ₂₆ H ₂₉ N ₅ O ₃ 0.3 H ₂ O	Requires	C	67.2	H	6.4	N	15.1%
--	----------	---	------	---	-----	---	-------

The starting material was prepared as follows:

A solution of 1-(3-hydroxypropyl)-4-methylpiperazine (2.4 g, 15 mmol), (prepared as described for the starting material in Example 133), in dichloromethane (60 ml) was treated with triethylamine (4.6 ml, 33 mmol) and p-toluenesulphonyl chloride (3.2 g, 17mmol) and stirred at ambient temperature for 2 hours. The solution was washed with saturated aqueous sodium hydrogen carbonate solution followed by water and filtered through phase separating paper. The filtrate was evaporated under vacuum to give 3-(4-methyl-piperazin-1-yl)propyl-4-toluene sulphonate as an oil which crystallised on standing (3.7 g, 78 %).

MS (ESI) : 313 (MH)⁺

A mixture of 2-amino-4-benzyloxy-5-methoxybenzamide (J. Med. Chem. 1977, vol 20, 146-149, 10g, 0.04mol) and Gold's reagent (7.4g, 0.05mol) in dioxane (100ml) was stirred and heated at reflux for 24 hours. Sodium acetate (3.02g, 0.037mol) and acetic acid (1.65ml, 0.029mol) were added to the reaction mixture and it was heated for a further 3 hours. The mixture was evaporated, water was added to the residue, the solid was filtered off, washed

with water and dried. Recrystallisation from acetic acid gave 7-benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (8.7g, 84%).

A mixture of 7-benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (2.82g, 0.01mol), thionyl chloride (40ml) and DMF (0.28ml) was stirred and heated at reflux for 1 hour. The mixture was evaporated and azeotroped with toluene to give 7-benzyloxy-4-chloro-6-methoxyquinazoline hydrochloride (3.45g).

4-Chloro-2-fluoro-phenol (264mg, 1.8mmol) was added to a solution of 7-benzyloxy-4-chloro-6-methoxyquinazoline hydrochloride (506mg, 1.5mmol) in pyridine (8ml) and the mixture heated at reflux for 45 minutes. The solvent was removed by evaporation and the residue partitioned between ethyl acetate and water. The organic layer was washed with 0.1M HCl, water and brine, dried (MgSO_4) and the solvent removed by evaporation. The solid residue was triturated with petroleum ether and the crude product collected by filtration and purified by flash chromatography eluting with methylene chloride/ether (9/1) to give 7-benzyloxy-4-(4-chloro-2-fluorophenoxy)-6-methoxyquinazoline (474mg, 77%) as a cream solid.

m.p. 179-180°C

^1H NMR Spectrum: (DMSO-d_6) 3.99(s, 3H); 5.36(s, 2H); 7.35-7.5(m, 4H); 7.55-7.65(m, 5H); 7.72(d, 1H); 8.6(s, 1H)

MS - ESI: 411 $[\text{MH}]^+$

Elemental analysis:	Found	C 63.38	H 4.07	N 6.78
$\text{C}_{22}\text{H}_{16}\text{ClFN}_2\text{O}_3 \cdot 0.06\text{H}_2\text{O} \cdot 0.05\text{CH}_2\text{Cl}_2$	Requires	C 63.64	H 3.93	N 6.73%

A solution of 7-benzyloxy-4-(4-chloro-2-fluorophenoxy)-6-methoxyquinazoline (451mg, 1.1mmol) in TFA (4.5ml) was heated at reflux for 3 hours. The mixture was diluted with toluene and the volatiles removed by evaporation. The residue was triturated with methylene chloride, collected by filtration, washed with ether and dried under vacuum to give 4-(4-chloro-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline (320mg, 90%).

^1H NMR Spectrum: (DMSO-d_6) 4.0(s, 3H); 7.27(s, 1H); 7.43(dd, 1H); 7.56(t, 1H); 7.57(s, 1H); 7.72(dd, 1H); 8.5(s, 1H)

MS - ESI: 321 $[\text{MH}]^+$

A mixture of the trifluoroacetic acid salt of 4-(4-chloro-2-fluorophenoxy)-7-hydroxy-6-methoxyquinazoline (3.2 g, 7.4 mmol), potassium carbonate (6.1 g, 44.2 mmol) and 3-(4-methyl-1-piperazinyl)propyl-4-toluene sulphonate (3.0 g, 9.6 mmol) in DMF (60 ml) was

stirred at 90 °C for 5 hours and allowed to cool to ambient temperature. The reaction mixture was poured into water (700 ml) and extracted 5 times with ethyl acetate. The combined extracts were washed with water, saturated aqueous sodium hydrogen carbonate, water and saturated brine. The ethyl acetate solution was dried over magnesium sulphate, filtered and the solvent removed under vacuum to give a residue which was purified by silica column chromatography, eluting with dichloromethane/methanol/0.880 aqueous ammonia (100/8/1). The relevant fractions were combined and evaporated under vacuum to give a residue which was triturated with ether, filtered and dried to give 4-(4-chloro-2-fluorophenoxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (1.64 g, 48 %).

¹H NMR Spectrum: (DMSO-d₆) 1.95 (m, 2H), 2.14 (s, 3H), 2.35 (m, 8H), 2.44 (t, 2H), 3.96 (s, 3H), 4.22 (t, 2H), 7.38 (s, 1H), 7.40 (m, 1H), 7.54 (m, 2H), 7.68 (m, 1H) and 8.55 (s, 1H)

MS (ESI) : 461 (MH)⁺

Elemental analysis	Found	C 59.6	H 5.7	N 12.2
--------------------	-------	--------	-------	--------

C ₂₃ H ₂₆ ClFN ₄ O ₃	Requires	C 59.9	H 5.7	N 12.2%
--	----------	--------	-------	---------

4-(4-Chloro-2-fluorophenoxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (2.6 g, 5.6 mmol) was treated with 2.0 N aqueous hydrochloric acid (45 ml) and the mixture stirred at 95 °C for 2 hours. The mixture was cooled, basified by the addition of solid sodium hydrogen carbonate and the water removed by azeotropeing with toluene. The residue was purified by silica column chromatography eluting with dichloromethane/methanol/0.880 aqueous ammonia (50/8/1) to give 6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)-3,4-dihydroquinazolin-4-one (1.8 g, 96%).

MS (ESI) : 333 (MH)⁺

6-Methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)-3,4-dihydroquinazolin-4-one (2.15 g, 6.48 mmol) was suspended in thionyl chloride (25 ml) and DMF (0.18 ml) and stirred under reflux for 2 hours. The thionyl chloride was evaporated under vacuum and the residue azeotroped twice with toluene. The residue was taken up in water, basified with saturated aqueous sodium hydrogen carbonate solution and the aqueous solution extracted 4 times with dichloromethane. The combined extracts were washed with water and brine then filtered through phase separating paper. The filtrate was evaporated under vacuum and the residue purified by silica column chromatography eluting with dichloromethane/methanol/0.880 aqueous ammonia (100/8/1) to give a solid which was triturated with a little acetone, filtered

and dried to give 4-chloro-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (1.2 g, 53 %). This was used without further purification.

MS (ESI) : 351 (MH)⁺

5 Example 177

A mixture of 4-chloro-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline (200 mg, 0.64 mmol), potassium carbonate (102 mg, 0.74 mmol) and 7-hydroxyquinoline (107 mg, 0.74 mmol) in DMSO (5 ml) was stirred at 100 °C for 5 hours and allowed to cool to ambient temperature. The mixture was poured into water, washed with dichloromethane and extracted
10 twice with a 10/1 mixture of dichloromethane/methanol. The extracts were washed with water and brine, dried over magnesium sulphate, filtered and the filtrate evaporated under vacuum. The residue was purified by silica column chromatography, eluting with dichloromethane/methanol/0.880 aqueous ammonia (100/8/1) to give an oil which crystallised on trituration with ether to give **6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(quinolin-7-
15 yloxy)quinazoline** (148 mg, 55 %).

¹H NMR Spectrum: (DMSO-d₆) 3.25 (s, 3H), 3.50 (t, 2H), 3.60 (t, 2H), 3.80 (t, 2H), 4.00 (s, 3H), 4.30 (t, 2H), 7.40 (s, 1H), 7.55 (m, 1H), 7.60 (m, 1H), 7.65 (s, 1H), 7.90 (d, 1H), 8.10 (d, 1H), 8.40 (m, 1H), 8.50 (s, 1H) and 8.90 (m, 1H)

MS (ESI) : 422 (MH)⁺

20	Elemental analysis	Found	C 65.8	H 5.2	N 10.0
	C ₂₃ H ₂₃ N ₃ O ₅	Requires	C 65.6	H 5.5	N 10.0%

The starting material was prepared as follows:

Diethyl azodicarboxylate (864 μl, 5.5 mmol) was added dropwise to a mixture of 7-
25 hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.2g, 3.9 mmol) (prepared as described for the starting material in Example 12), triphenylphosphine (1.44g, 5.5 mmol) and 2-(2-methoxyethoxy)ethanol (653 μl, 5.5 mmol) in methylene chloride (70 ml) cooled at 0 °C. The mixture was stirred for 1.5 hours at ambient temperature and the solvent was removed by evaporation. The residue was purified by column chromatography eluting
30 with a mixture of ethyl acetate/methylene chloride (50/50 followed by 80/20). The purified solid was suspended in ether, collected by filtration and dried under vacuum to give 6-

methoxy-7-(2-(2-methoxyethoxy)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.70g, 100%).

¹H NMR Spectrum: (DMSO-d₆) 1.13(s, 9 H); 3.26(s, 3H); 3.5(m, 2H); 3.65(m, 2H); 3.85(m, 2H); 3.91(s, 3H); 4.3(m, 2H); 5.9(s, 2H); 7.2(s, 1H); 7.5(s, 1H); 8.4(s, 1H)

5 Saturated methanolic ammonia (20ml) was added to a solution of 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (2.26g, 5.5mmol) in a mixture of ethanol (40ml) and methylene chloride (15ml). The mixture was stirred for 24 hours at ambient temperature, and further methanolic ammonia (20ml) was added. The mixture was stirred for a further 24 hours at ambient temperature and the volatiles
10 were removed by evaporation. The residue was triturated with ether, collected by filtration, dried under vacuum to give 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-3,4-dihydroquinazolin-4-one (975mg, 78%).

¹H NMR Spectrum: (DMSO-d₆) 3.25(s, 3H); 3.45(t, 2H); 3.6(t, 2H); 3.8(t, 2H); 3.9(s, 3H); 4.2(t, 2H); 7.15(s, 1H); 7.45(s, 1H); 8.0(s, 1H)

15 MS - EI: 294 [M]⁺

A solution of 6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-3,4-dihydroquinazolin-4-one (930mg, 3.16mmol) in thionyl chloride (15ml) and DMF (150μl) was heated at 60°C for 1.5 hours. The mixture was allowed to cool and the volatiles were removed by evaporation and by azeotroping with toluene. The residue was dissolved in methylene chloride and 5%
20 aqueous sodium hydrogen carbonate solution was added until the aqueous layer was at pH8. The organic layer was separated, washed with brine, dried (MgSO₄) and the solvent removed by evaporation. The residue was purified by flash chromatography eluting with ethyl acetate to give 4-chloro-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline (863mg, 87%).

¹H NMR Spectrum: (DMSO-d₆) 3.24(s, 3H); 3.47(m, 2H); 3.62(m, 2H); 3.84(t, 2H); 4.01(s, 3H); 4.25(t, 2H); 7.41(s, 1H); 7.49(s, 1H); 8.88(s, 1H)
25

Example 178

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (168 mg, 0.5 mmol), (prepared as described for the starting material in Example 67), potassium carbonate
30 (207 mg, 1.5 mmol), 3-methyl-5-hydroxyindole (88mg, 0.6 mmol), (Can. J. Chem. 1964, 42, 514), and DMA (2.0 ml) was purged with nitrogen for 5 minutes at 25°C. This mixture was then stirred at 100°C for 3 hours then allowed to cool to ambient temperature, was filtered and

the filtrate evaporated under vacuum. The residue was purified by silica column chromatography eluting with dichloromethane/methanolic ammonia (7M) (90/10) to give **6-methoxy-4-(3-methylindol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline** (155 mg, 69%).

¹H NMR Spectrum: (DMSO-d₆) 1.37 (m, 2H), 1.50 (m, 4H), 1.95 (m, 2H), 2.21 (s, 3H), 2.34 (m, 4H), 2.42 (t, 2H), 3.96 (s, 3H), 4.22 (t, 2H), 6.95 (dd, 1H), 7.16 (s, 1H), 7.35 (m, 3H), 7.58 (s, 1H), 8.48 (s, 1H) and 10.82 (s, 1H)

MS (ESI) : 447 (MH)⁺

Elemental analysis Found C 68.2 H 6.8 N 12.6

C₂₆H₃₀N₄O₃ 0.5 H₂O, Requires C 68.5 H 6.8 N 12.3%

Example 179

Using an analogous procedure to that described in Example 178, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline, (prepared as described for the starting material in Example 9), was used to give

6-methoxy-4-(3-methylindol-5-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (154 mg, 79%).

¹H NMR Spectrum: (DMSO-d₆) 1.68 (m, 4H), 1.97 (m, 2H), 2.22 (s, 3H), 2.43 (m, 4H), 2.55 (t, 2H), 3.96 (s, 3H), 4.22 (t, 2H), 6.93 (dd, 1H), 7.16 (s, 1H), 7.35 (m, 3H), 7.58 (s, 1H), 8.48 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 433 (MH)⁺

m.p. 75-77°C

Example 180

Using an analogous procedure to that described in Example 178, 4-chloro-6-methoxy-7-(2-piperidinoethoxy)quinazoline was used to give **6-methoxy-4-(3-methylindol-5-yloxy)-7-(2-piperidinoethoxy)quinazoline** (156mg, 80%).

¹H NMR Spectrum: (DMSO-d₆) 1.38 (m, 2H), 1.50 (m, 4H), 2.24 (s, 3H), 2.73 (t, 2H), 3.96 (s, 3H), 4.28 (t, 2H), 6.93 (dd, 1H), 7.16 (s, 1H), 7.32 (d, 1H), 7.37 (m, 2H), 7.58 (s, 1H), 8.47 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 433 (MH)⁺

Elemental analysis Found C 67.0 H 6.5 N 13.0

C₂₅H₂₈N₄O₃ 0.75 H₂O Requires C 67.3 H 6.6 N 12.6%

The starting material was prepared as follows:

1-(2-Chloroethyl)piperidine hydrochloride (0.83g, 4.5mmol) was added to 7-hydroxy-6-methoxy-4-phenoxyquinazoline (1.0g, 3.73mmol), (prepared as described for the starting
5 material in Example 1), and potassium carbonate (2.6g, 18.8mmol) in DMF (30ml), and the mixture heated at 110°C for 2.5 hours and allowed to cool. The insolubles were removed by filtration, and the volatiles were removed from the filtrate by evaporation. The residue was purified by column chromatography eluting with methylene chloride/methanol (9/1) to give 6-methoxy-4-phenoxy-7-(2-piperidinoethoxy)quinazoline (1.2g, 85%).

10 ¹H NMR Spectrum: (DMSO-d₆) 1.38(m, 2H); 1.50(m, 4H); 2.4-2.5(m, 4H); 2.75(t, 2H); 3.95(s, 3H); 4.27(t, 2H); 7.30(m, 3H); 7.40(s, 1H); 7.46(m, 2H); 7.54(s, 1H); 8.52(s, 1H)

MS - ESI: 380 [MH]⁺

A mixture of 6-methoxy-4-phenoxy-7-(2-piperidinoethoxy)quinazoline (1.15g, 3.0mmol) and 2M hydrochloric acid (20ml) was heated at 90°C for 2 hours and allowed to
15 cool. The mixture was neutralised with solid sodium hydrogen carbonate and extracted with methylene chloride. The organic phase was separated, passed through phase separating paper and the volatiles removed by evaporation to give a solid product (230mg). The aqueous phase was adjusted to pH10, the resulting precipitate was collected by filtration, washed with water and dried to give a second crop of product (220mg). The products were combined to give 6-methoxy-7-(2-piperidinoethoxy)-3,4-dihydroquinazolin-4-one (450mg, 50%).

20 MS - ESI: 304 [MH]⁺

A mixture of 6-methoxy-7-(2-piperidinoethoxy)-3,4-dihydroquinazolin-4-one (440mg, 1.45mmol), thionyl chloride (15ml) and DMF (3 drops) was heated at reflux for 3 hours then allowed to cool. The excess thionyl chloride was removed by evaporation and the residue was
25 azeotroped with toluene to give a crude 4-chloro-6-methoxy-7-(2-piperidinoethoxy)quinazoline hydrochloride (640mg).

4-Chloro-6-methoxy-7-(2-piperidinoethoxy)quinazoline hydrochloride was suspended in methylene chloride (10ml) and saturated aqueous sodium hydrogen carbonate solution (5ml) then stirred vigorously for 10 minutes at ambient temperature. The layers were
30 separated and the organic layer dried (MgSO₄) then evaporated to give a white solid. This solid was triturated with methanol (2.5ml), the resulting solid filtered off, washed with cold methanol and dried to give 4-chloro-6-methoxy-7-(2-piperidinoethoxy)quinazoline (0.36g).

Example 181

Using an analogous procedure to that described in Example 178, 4-chloro-6-methoxy-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline, (prepared as described for the starting material in Example 152), was used to give

6-methoxy-4-(3-methylindol-5-yloxy)-7-(3-(N-methyl-N-methylsulphonylamino)propoxy)quinazoline (104mg, 49%).

¹H NMR Spectrum: (DMSO-d₆) 2.08 (m, 2H), 2.22 (s, 3H), 2.80 (s, 3H), 2.88 (s, 3H), 3.27 (t, 2H), 3.97 (s, 3H), 4.22 (t, 2H), 6.95 (dd, 1H), 7.17 (s, 1H), 7.35 (m, 3H), 7.59 (s, 1H), 8.48 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 471 (MH)⁺

Elemental analysis	Found	C	H	N
		57.0	5.6	11.4

C ₂₃ H ₂₆ F ₄ N ₄ O ₅ S 0.5 H ₂ O,	Requires	C 57.5	H 5.7	N 11.7%
--	----------	--------	-------	---------

Example 182

A mixture of 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (218 mg, 0.68 mmol), (prepared as described for the starting material in Example 9), 5-hydroxy-1*H*-pyrrolo[2,3-*b*]pyridine (100 mg, 0.75 mmol) and potassium carbonate (280 mg, 2.0mmol) in DMF (4 ml) was stirred at 95°C for 6 hours and allowed to cool to ambient temperature. The reaction mixture was treated with 1.0 N aqueous sodium hydroxide solution and allowed to stir at ambient temperature for a few minutes. The resulting precipitate was filtered off, washed with water and air dried to give a crude product which was purified by column chromatography, eluting with dichloromethane/methanol/880 ammonia (100/8/1). The relevant fractions were combined and evaporated '*in vacuo*' to give a white solid. This was recolumned using dichloromethane/methanol (4/1) solvent to give a white solid which was triturated with acetone, filtered and dried to give **6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)-4-(1*H*-pyrrolo[2,3-*b*]pyridin-5-yloxy)quinazoline** (50 mg, 18%).

m.p. 184.0 - 185.5°C

¹H NMR Spectrum: (DMSO-d₆) 1.70 (m, 4H), 1.99 (m, 2H), 2.46 (m, 4H), 2.58 (t, 2H), 4.00 (s, 3H), 4.26 (t, 2H), 6.48 (t, 1H), 7.36 (s, 1H), 7.55 (t, 1H), 7.60 (s, 1H), 7.92 (d, 1H), 8.19 (d, 1H), 8.50 (s, 1H) and 11.78 (br s, 1H)

MS (ESI) : 420 (MH)⁺

- 210 -

Elemental analysis	Found	C 63.9	H 5.9	N 16.1
$C_{23}H_{25}N_5O_3 \cdot 0.7 H_2O$	Requires	C 63.9	H 6.2	N 16.2%

The starting material was prepared as follows:-

5 A suspension of 5-methoxy-1*H*-pyrrolo[2,3-*b*]pyridine (210 mg, 1.42 mmol), (Heterocycles 50, (2), 1065 - 1080, (1999)), in dichloromethane (10 ml) was stirred in an inert atmosphere, a 1.0M solution of boron tribromide in dichloromethane (4.3 ml, 4.3 mmol) added dropwise and the mixture stirred at ambient temperature overnight. The reaction mixture was taken to pH6 by the dropwise addition of 5N aqueous sodium hydroxide and
10 further diluted with water. The aqueous solution was extracted several times with ethyl acetate, the extracts combined, washed with water followed by brine and dried over magnesium sulphate. The ethyl acetate solvent was removed '*in vacuo*' and the residue purified by column chromatography, eluting with dichloromethane/methanol (95/5), to give a white solid. The solid was triturated with ether, filtered and dried to give 5-hydroxy-1*H*-
15 pyrrolo[2,3-*b*]pyridine (108 mg, 57%).

m.p. 206-209°C

¹H NMR Spectrum: (DMSO-d₆) 6.25 (s,1H), 7.27 (s,1H), 7.33 (s, 1H), 7.82 (s, 1H), 9.00 (s,1H) and 11.20 (s, 1H)

MS (ESI) : 135 (MH)⁺

Example 183

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (168 mg, 0.5 mmol), (prepared as described for the starting material in Example 67), potassium carbonate (345 mg, 5.0 mmol), 5-hydroxy-2-indolecarboxylic acid (106mg, 0.6 mmol) and DMA (2.0
25 ml) was purged with nitrogen for 5 minutes at 25°C. This mixture was then stirred at 100°C for 3 hours, allowed to cool to ambient temperature, filtered and the filtrate evaporated under vacuum. The residue was purified on octadecylsilane reverse phase silica eluting with acetonitrile/water/trifluoroacetic acid (as a gradient from 30/69.8/0.2 to 50/49.8/0.2) and the product further purified by silica column chromatography eluting with
30 dichloromethane/methanolic ammonia (7M) (90/10) to give 4-(2-carboxyindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline (85 mg 36%).

¹H NMR Spectrum: (DMSO-d₆) 1.43 (m, 2H), 1.56 (m, 4H), 2.04 (m, 2H), 2.59 (m, 6H), 3.97 (s, 3H), 4.24 (t, 2H), 7.01 (s, 1H), 7.11 (dd, 1H), 7.36 (s, 1H), 7.48 (m, 2H), 7.58 (s, 1H), 8.48 (s, 1H) and 11.53 (br s, 1H)

MS (ESI) : 477 (MH)⁺

5

Example 184

4-Chloro-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline (0.15 g, 0.45 mmol), (prepared as described for the starting material in Example 50), potassium carbonate (94 mg, 0.68 mmol) and 7-hydroxyquinoline (79 mg, 0.54 mmol) were suspended in anhydrous DMF (1.5 ml) and heated to 90°C overnight. The compound was precipitated upon addition of water. The precipitate was collected by filtration, washed with water and dried under vacuum over phosphorus pentoxide to give **6-methoxy-7-(3-methylsulphonylpropoxy)-4-(quinolin-7-yloxy)quinazoline** (161 mg, 81%).

10

¹H NMR Spectrum: (DMSO-d₆) 2.26 (m, 2H); 3.08 (s, 3H); 3.35 (m, 2H); 4.03 (s, 3H); 4.38 (m, 2H); 7.45 (s, 1H); 7.60 (m, 1H); 7.65 (m, 1H); 7.70 (s, 1H); 7.95 (d, 1H); 8.15 (d, 1H); 8.46 (d, 1H); 8.60 (s, 1H); 8.95 (d, 1H)

15


MS (ESI) : 440 [MH]⁺

Examples 185-188

Using an analogous procedure to that described in Example 184, 4-chloro-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline (0.15 g, 0.45 mmol), (prepared as described for the starting material in Example 50), was reacted with the appropriate phenols to give the compounds in Table X.





20

Table X



25

Example number	weight (mg)	yield %	MS-ESI [MH] ⁺	AR	note

185	199	93	474		a
186	171	85	422		b
187	183	88	460		c
188	83	40	455		d

a) Using 4-chloro-7-hydroxyquinoline (96 mg) gave **4-(4-chloroquinolin-7-yloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.24 (m, 2H); 3.04 (s, 3H); 3.35 (m, 2H); 3.99 (s, 3H); 4.32

5 (m, 2H); 7.42 (s, 1H); 7.64 (s, 1H); 7.80 (d, 2H); 8.04 (d, 1H); 8.29 (d, 1H); 8.55 (s, 1H); 8.87 (d, 1H)

b) Using 5-hydroxy-2-methylindole (80 mg) gave **6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-methylsulphonylpropoxy)quinazoline**.

10 ¹H NMR Spectrum: (DMSO-d₆) 2.24 (m, 2H); 2.40 (s, 3H); 3.05 (s, 3H); 3.35 (m, 2H); 4.0 (s, 3H); 4.32 (m, 2H); 6.13 (s, 1H); 6.88 (d, 1H); 7.25 (d, 1H); 7.32 (d, 1H); 7.39 (s, 1H); 7.60 (s, 1H); 8.50 (s, 1H)

c) Using 5-hydroxy-2-methylbenzothiazole (90 mg) gave **6-methoxy-4-(2-methyl-1,3-benzothiazol-5-yloxy)-7-(3-methylsulphonylpropoxy)quinazoline**.

15 ¹H NMR Spectrum: (DMSO-d₆) 2.24 (m, 2H); 2.28 (s, 3H); 3.05 (s, 3H); 3.35 (m, 2H); 4.0 (s, 3H); 4.32 (m, 2H); 7.36 (d, 1H); 7.41 (s, 1H); 7.65 (s, 1H); 7.87 (d, 1H); 8.11 (d, 1H); 8.53 (s, 1H)

20 d) Using 2,7-dihydroxynaphthalene (87 mg) gave **4-(7-hydroxy-2-naphthyloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.24 (m, 2H); 3.05 (s, 3H); 3.35 (m, 2H); 3.98 (s, 3H); 4.32 (m, 2H); 7.06 (d, 1H); 7.12 (s, 1H); 7.18 (d, 1H); 7.40 (d, 1H); 7.59 (m, 2H); 7.85 (m, 2H); 8.55 (d, 1H); 9.8 (br s, 1H)

5 Example 189

To a portion of 2-chloro-5-hydroxybenzimidazole (191 mg, 0.75 mmol) in DMF (3 ml) was added sodium hydride (60 mg, 1.5 mmol) under argon at ambient temperature. Ten minutes later 4-chloro-6-methoxy-7-(1-methylpiperidin-4-yl)methoxyquinazoline (200 mg, 0.62 mmol), (prepared as described for the starting material in Example 10), was added and the mixture heated at 100 °C for 2 hours. More 2-chloro-5-hydroxybenzimidazole (30 mg, 0.12 mmol) and sodium hydride (11 mg, 0.28 mmol) were then added as the reaction was found to be incomplete. The heating was continued for an additional 1 hour. Work-up using ethyl acetate and a saturated aqueous solution of ammonium chloride followed by drying of the organic phase (MgSO₄) and evaporation of the solvent gave a crude product which was adsorbed on alumina using dichloromethane/methanol and purified by flash chromatography using neutral alumina and dichloromethane/methanol (98:2) as the eluent.

Evaporation of the solvent and trituration in ether gave **4-(2-chloro-1H-benzimidazol-6-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline** (46 mg, 16%).

¹H NMR Spectrum: (DMSO-d₆ + TFA) 1.60 (m, 2H); 2.05 (d, 2H); 2.15 (m, 1H); 2.80 (s, 3H); 3.05 (m, 2H); 3.55 (m, 2H); 4.05 (s, 3H); 4.15 (d, 2H); 7.20 (dd, 1H); 7.50 (dd, 2H); 7.65 (d, 1H); 7.70 (s, 1H); 8.80 (s, 1H)
MS (ESI) : 454 [MH]⁺

The starting material was synthesised as follows:

25 2-Chloro-5-methoxybenzimidazole (0.3 g, 1.64 mmol) was suspended in dichloromethane (20 ml) under argon followed by the addition of boron tribromide (233 ul, 2.46 mmol). The reaction mixture was stirred for 2 hours at ambient temperature. The solvent was evaporated and the resulting powder was added in portions to methanol (30 ml). Silica was added and the solvent was evaporated. The resulting powder was placed on the top of a silica column and the product was eluted off using dichloromethane/methanol (95/5).
30 Evaporation of the solvent and trituration in ether gave 2-chloro-5-hydroxybenzimidazole (440 mg, 99%).

Example 190

Using an analogous procedure to that described in Example 189, 4-chloro-6-methoxy-7-(1-methylpiperidin-4-yl)methoxyquinazoline, (prepared as described for the starting material in Example 10), was reacted with 5-hydroxy-2-methylbenzimidazole (200 mg, 0.62 mmol) and after work-up and purification on a 10 g silica ISOLUTE column using successively dichloromethane, dichloromethane/methanol (95/5) and dichloromethane/methanol saturated with ammonia (95/5), gave **6-methoxy-4-(2-methyl-1H-benzimidazol-6-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline** (68 mg, 25%).

¹H NMR Spectrum: (DMSO-d₆ + TFA) 1.60 (m, 2H); 2.10 (m, 2H); 2.20 (m, 1H); 2.80 (s, 3H); 2.85 (s, 3H); 3.05 (m, 2H); 3.50 (m, 2H); 4.05 (s, 3H); 4.15 (d, 2H); 7.50 (s, 1H); 7.55 (d, 1H); 7.70 (s, 1H); 7.85 (d, 1H); 7.90 (d, 1H); 8.65 (s, 1H)

MS (ESI) : 434 [MH]⁺

The starting material was prepared as follows:

The free base of 4-methoxy-1,2-phenylenediamine dihydrochloride (10 g) was obtained by shaking it with a mixture of ethyl acetate and a saturated aqueous solution of sodium hydrogen carbonate. The organic phase was then washed with brine, dried (MgSO₄) and the solvent evaporated. The obtained dark oil (6.08 g, 50 mmol) was solubilised in toluene (60 ml) and p-toluene sulfonic acid (60 mg) and triethyl orthoacetate (9.15 ml, 50 mmol) were added in turn. The mixture was heated to 110 °C until no more ethanol distilled off. The remaining toluene was removed by rotary evaporation and the residue purified by flash chromatography using dichloromethane/methanol (95/5) as the eluent. The obtained dark oil was triturated in ether and the solid collected by filtration to give 5-methoxy-2-methylbenzimidazole (4.15 g, 51%).

¹H NMR Spectrum (DMSO-d₆ + TFA) 2.75 (s, 3H); 3.85 (s, 3H); 7.15 (dd, 1H); 7.25 (s, 1H); 7.70 (d, 1H)

Using an analogous procedure to that described for the synthesis of 2-chloro-5-hydroxybenzimidazole in Example 189, 5-methoxy-2-methylbenzimidazole (4.0 g, 25 mmol) was reacted with boron tribromide (7 ml, 74 mmol) in dichloromethane (150 ml) to give, after work-up and purification by flash chromatography using dichloromethane/methanol (90/10), 5-hydroxy-2-methylbenzimidazole (4.4 g, 76%).

¹H NMR Spectrum (DMSO-d₆) 2.70 (s, 3H); 6.95 (dd, 1H); 7.00 (d, 1H); 7.55 (d, 1H)

Example 191

4-Chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (200 mg, 0.62 mmol), (prepared as described for the starting material in Example 10), was suspended in DMF (3 ml) under argon. 3-Cyano-7-hydroxyquinoline (116 mg, 0.68 mmol) and potassium carbonate (129 mg, 0.93 mmol) were added and the reaction mixture was heated at 95 °C for 90 minutes. Upon cooling to ambient temperature the mixture was diluted with dichloromethane and poured on the top of an ISOLUTE silica column. Elution was done using successively dichloromethane, dichloromethane/methanol (95/5) and dichloromethane/methanol saturated with ammonia (95/5). Evaporation of the solvent and trituration of the solid in ether gave **4-(3-cyanoquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline** (244 mg, 86%).

¹H NMR Spectrum: (DMSO-d₆ + TFA) 1.60 (m, 2H); 2.10 (m, 3H); 2.85 (s, 3H); 3.05 (m, 2H); 3.55 (m, 2H); 4.05 (s, 3H); 4.20 (d, 2H); 7.55 (s, 1H); 7.80 (s, 1H); 7.85 (dd, 1H); 8.15 (s, 1H); 8.3 (d, 1H); 8.85 (s, 1H); 9.20 (s, 1H); 9.25 (s, 1H)

MS (ESI) : 456 [MH]⁺ 456

The starting material was prepared as follows:

m-Anisidine (50 g, 407 mmol) and diethyl ethoxymethylenemalonate (102 g, 407 mmol) were heated at 60 °C for 20 minutes. Diphenyl ether (270 ml) was then added and the temperature was raised to 240 °C over 30 minutes. The ethanol formed distilled off. Heating was maintained at this temperature for 1 hour then the reaction mixture was allowed to cool to 120 °C at which point the reaction mixture was diluted with heptane and allowed to stand overnight at ambient temperature. The brown solid was collected by filtration and washed with methanol and ether to give ethyl 7-methoxy-4-oxo-1,4-dihydroquinoline-3-carboxylate (45 g, 45%). This reaction was repeated twice.

¹H NMR Spectrum: (DMSO-d₆) 1.25 (t, 3H); 3.85 (s, 3H); 4.20 (q, 2H); 6.95 (d, 1H); 7.00 (s, 1H); 8.05 (d, 1H); 8.50 (s, 1H)

Phosphorus oxychloride (88 ml) was added to ethyl 7-methoxy-4-oxo-1,4-dihydroquinoline-3-carboxylate (58 g, 235 mmol) and the mixture was heated at reflux for 45 minutes under anhydrous conditions. Upon cooling to ambient temperature, phosphorus

oxychloride was evaporated and the solid residue was added in portions to a mixture of ammonia (150 ml) and ice (200g). External cooling as well as further addition of ammonia to maintain the pH around 8 was needed during this hydrolysis step. The aqueous phase was extracted with dichloromethane and the organic phase was washed with water and brine, dried (MgSO₄) and concentrated to about 300 ml. Pentane (400 ml) was added and the precipitate formed collected by filtration. Drying under vacuum gave 4-chloro-3-ethoxycarbonyl-7-methoxyquinoline (45.5 g, 73 %).

¹H NMR Spectrum: (DMSO-d₆) 1.40 (t, 3H); 4.00 (s, 3H); 4.45 (q, 2H); 7.45 (dd, 1H); 7.55 (d, 1H); 8.30 (d, 1H); 9.10 (s, 1H)

4-Chloro-3-ethoxycarbonyl-7-methoxyquinoline (43 g, 162 mmol) was dissolved in acetic acid (250 ml), with 10% palladium on charcoal (1.5 g) and hydrogenated at atmospheric pressure during 8 hours. The catalyst was removed by filtration over a pad of celite and the solvent evaporated. The residue was diluted with water and the pH adjusted to 7-8 with a saturated solution of sodium hydrogen carbonate. The solid was collected by filtration, washed with water and dried under vacuum over phosphorus pentoxide to give 3-ethoxycarbonyl-7-methoxyquinoline (33 g, 88%) as a beige powder.

¹H NMR Spectrum: (DMSO-d₆) 1.40 (t, 3H); 3.95 (s, 3H); 4.40 (q, 2H); 7.35 (dd, 1H); 7.50 (d, 1H); 8.15 (d, 1H); 8.90 (d, 1H); 9.25 (d, 1H)

3-Ethoxycarbonyl-7-methoxyquinoline (28 g, 120 mmol) was added to a methanol solution saturated with ammonia. The suspension was stirred at ambient temperature in a glass pressure vessel for 2 weeks. The white solid was collected by filtration, washed with methanol and dried under vacuum to give 3-carbamoyl-7-methoxyquinoline (21g, 86%).

¹H NMR Spectrum (DMSO-d₆) 3.95 (s, 3H); 7.35 (dd, 1H); 7.45 (d, 1H); 7.60 (br s, 1H); 8.00 (d, 1H); 8.20 (br s, 1H); 8.75 (s, 1H); 9.25 (s, 1H)

3-Carbamoyl-7-methoxyquinoline (4 g, 20 mmol) was suspended in anhydrous dichloromethane (60 ml) under argon. Anhydrous dimethyl sulphoxide (2.25 ml, 32 mmol) was added, the mixture was cooled to -78 °C and a solution of oxalyl chloride (2.08 ml, 24 mmol) in dichloromethane (20 ml) was added dropwise over the course of 1 hour. 15 Minutes after the end of the addition, triethylamine (8.3 ml, 60 mmol) was added dropwise and the heterogeneous reaction mixture stirred for an additional 1 hour at -78 °C then left to rise to ambient temperature. The unreacted starting material was removed by filtration and the filtrate was diluted with water and extracted with ethyl acetate. The organic phases were

combined, washed with brine, dried (MgSO_4) and the solvent evaporated. The residue was purified by flash chromatography using dichloromethane/methanol (97/3). The obtained solid was triturated with ether and gave, after drying under vacuum, 3-cyano-7-methoxyquinoline (1.47 g, 40%).

- 5 ^1H NMR Spectrum (DMSO-d_6) 4.00 (t, 3H); 7.40 (dd, 1H); 7.50 (d, 1H); 8.00 (d, 1H); 8.95 (s, 1H); 9.10 (d, 1H)

3-Cyano-7-methoxyquinoline (380 mg, 2.1 mmol) was suspended in benzene (10 ml), aluminium trichloride (826 mg, 6.2 mmol) was added and the mixture heated at reflux for 30 minutes. More aluminium trichloride (275 mg, 2.1 mmol) was added and the mixture
10 refluxed for a further 2 hours. The solvent was evaporated, the dark green solid was added to ice and extracted with ethyl acetate. The organic phase was washed with brine, dried (MgSO_4) and evaporated. The solid was found to contain some aluminium salts which were removed as follows. The solid was dissolved in dichloromethane (200 ml) was stirred vigorously with a saturated sodium hydrogen carbonate solution for 1 hour. The product was
15 collected by filtration of the aqueous phase and dried over phosphorus pentoxide under vacuum to give 3-cyano-7-hydroxyquinoline (238 mg, 68%).

^1H NMR Spectrum (DMSO-d_6) 7.25 (d, 1H); 7.30 (d, 1H); 7.95 (d, 1H); 8.85 (d, 1H); 9.00 (d, 1H)

20 Example 192

To 6-methoxy-7-(3-morpholinopropoxy)-4-((1-*tert*butoxycarbonyl-1,2,3,4-tetrahydroquinolin-6-yl)oxy)quinazoline (110 mg, 0.2 mmol) in solution in dichloromethane (3 ml) was added TFA (0.3 ml) and the mixture stirred for 1 hour at ambient temperature. The solvents were evaporated and the remaining oil was diluted with dichloromethane and the
25 pH adjusted to 9 with a saturated solution of sodium hydrogen carbonate. The organic phase was washed with, brine, dried (MgSO_4), filtered and the solvent evaporated to give 6-methoxy-7-(3-morpholinopropoxy)-4-(1,2,3,4-tetrahydroquinolin-6-yloxy)quinazoline (84 mg, 93%).

^1H NMR Spectrum: (CDCl_3) 1.95 (m, 2H); 2.15 (m, 2H); 2.45 (m, 4H); 2.60 (t, 2H); 2.80 (t, 2H); 3.35 (t, 2H); 3.75 (m, 4H); 3.90 (br s, 1H); 4.05 (s, 3H); 4.30 (t, 2H); 6.55 (d, 1H); 6.85 (m, 2H); 7.30 (s, 1H); 7.55 (s, 1H); 8.65 (s, 1H)

MS (ESI) : 451 $[\text{MH}]^+$

- 218 -

Elemental analysis:	Found	C 66.4	H 6.9	N 12.4
$C_2H_1N_1O_1$; 1 HCl, 2 H_2O	Requires	C 66.7	H 6.7	N 12.4%

The starting material was prepared as follows:

5 6-Hydroxyquinoline (1 g, 6.9 mmol) was dissolved in methanol and hydrogenated at 3 atmospheres pressure with platinum(IV) oxide (276 mg) over 24 hours. The catalyst was removed by filtration over a pad of celite and the solvent was evaporated. The solid was washed with ether to give 6-hydroxy-(1,2,3,4)-tetrahydroquinoline (698 mg, 68 %).

1H NMR Spectrum ($DMSO-d_6$) 1.75 (m, 2H); 2.60 (m, 2H); 3.05 (m, 2H); 4.90 (br s, 1H); 6.30 (m, 3H); 8.25 (br s, 1H)

10 6-Hydroxy-(1,2,3,4)-tetrahydroquinoline (250 mg, 1.7 mmol) was suspended in acetone (1 ml) and trichloromethane (1 ml) under argon. *Tert*-Butoxycarbonylanhydride (365 mg, 1.7 mmol) in solution in acetone was added dropwise followed by THF (2ml) to help the solubilisation. The reaction mixture was stirred overnight at ambient temperature, the solvent was evaporated, the residue was partitioned between ethyl acetate and water, the organic phase was washed with water, brine, dried ($MgSO_4$), filtered and the solvent evaporated. The resulting gum was purified by flash chromatography using dichloromethane/methanol (97/3) as solvent. Evaporation of the solvent gave 6-hydroxy-4-(1-*tert*butoxycarbonyl-1,2,3,4-tetrahydroquinoline (344 mg, 82%) as a brown foam.

20 1H NMR Spectrum: ($DMSO-d_6$) 1.50 (m, 9H); 1.90 (m, 2H); 2.70 (t, 2H); 3.65 (t, 2H); 4.75 (br s, 1H); 6.55 (d, 1H); 6.65 (dd, 1H); 7.45 (d, 1H)

25 6-Hydroxy-4-(1-*tert*butoxycarbonyl-1,2,3,4-tetrahydroquinoline (82 mg, 0.32 mmol) was dissolved in anhydrous dimethylformamide under argon, with potassium carbonate (61 mg, 0.44 mmol) and 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (100 mg, 0.3 mmol), (prepared as described for the starting material in Example 1). No reaction occurred after 2 hours at 60 °C. Sodium hydride (12 mg, 0.3 mmol) was added and the reaction mixture was heated at 120 °C for 90 minutes. The cooled mixture was poured into water and ethyl acetate. The organic phase was washed with water, brine, dried ($MgSO_4$), filtered and the solvent evaporated. The residue was purified by flash chromatography using first

30 dichloromethane/methanol (97/3) as solvent. Evaporation of the solvent gave 6-methoxy-7-(3-morpholinopropoxy)-4-((1-*tert*butoxycarbonyl-1,2,3,4-tetrahydroquinolin-6-yl)oxy)quinazoline (115 mg, 71%) as a white solid.

¹H NMR Spectrum: (DMSO-d₆) 1.55 (s, 9H); 1.95 (m, 2H); 2.15 (m, 2H); 2.50 (m, 4H); 2.60 (t, 2H); 2.85 (t, 2H); 3.75 (m, 6H); 4.05 (s, 3H); 4.30 (t, 2H); 7.00 (m, 2H); 7.35 (s, 1H); 7.55 (s, 1H); 7.80 (d, 1H); 8.65 (s, 1H)

5 Example 193

Using an analogous procedure to that described in Example 192, 4-(1-*tert*butoxycarbonyl-2,3-dihydro-indol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (169 mg, 0.32 mmol) was reacted with TFA (1 ml) to give, after work-up and purification, 4-(2,3-dihydro-1*H*-indol-5-yl)oxy-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (124 mg, 91%).

¹H NMR Spectrum: (CDCl₃) 1.90 (br, 4H); 2.30 (br, 2H); 2.70 (br d, 6H); 3.10 (t, 2H); 3.65 (t, 2H); 4.05 (s, 3H); 4.30 (t, 2H); 6.70 (d, 1H); 6.80 (dd, 1H); 7.00 (s, 1H); 7.30 (s, 1H); 7.55 (s, 1H); 8.65 (s, 1H)

MS (ESI) : 421 [MH]⁺

The starting material was prepared as follows:

5-Hydroxyindole (2 g, 15 mmol) was dissolved in methanol (60 ml) under argon. Sodium cyanoborohydride (1.89 g, 30 mmol) and trifluoroboron etherate (4.2 ml, 33 mmol) were added and the mixture was heated at reflux for 3 hours then left to cool to ambient temperature. The solvent was evaporated and the residue was partitioned between ethyl acetate and water. Ammonia was added to adjust the pH to 10 and the aqueous phase was extracted with more ethyl acetate. The combined organic phases were washed with water, brine, dried (MgSO₄), filtered and the solvent evaporated. The residue was purified by flash chromatography using dichloromethane/methanol (95/5) as solvent. Evaporation of the solvent gave 5-hydroxy-2,3-dihydro-1*H*-indole (1.45 g, 73%) as an off white solid.

¹H NMR Spectrum: (DMSO-d₆, TFA) 3.15 (t, 2H); 3.70 (t, 2H); 6.75 (dd, 1H); 6.85 (d, 1H); 7.30 (d, 1H)

5-Hydroxy-2,3-dihydro-1*H*-indole (1.5 g, 11.1 mmol) was suspended in a mixture of acetone (7 ml) trichloromethane (7 ml) and THF (6 ml). *tert*-Butoxycarbonylanhydride (2.42 g, 11 mmol) in solution in THF (7 ml) was added dropwise. The reaction mixture was stirred overnight at ambient temperature, the solvent was evaporated, the residue was partitioned between ethyl acetate and water, the organic phase was washed with water, brine, dried

(MgSO₄), filtered and the solvent evaporated. The solid was purified by flash chromatography using dichloromethane/methanol (95/5) as solvent. Evaporation of the solvent gave 5-hydroxy-(1-*tert*butoxycarbonyl)-2,3-dihydroindole (2.28 g, 87%) as an off white solid.

5 ¹H NMR Spectrum: (CDCl₃) 3.05 (t, 2H); 3.95 (br s, 2H); 4.70 (br s, 1H); 6.60 (d, 1H); 6.65 (s, 1H); 7.70 (br s, 1H)

Sodium hydride (22 mg, 0.56 mmol) was suspended in anhydrous dimethylformamide under argon. 5-Hydroxy-(1-*tert*butoxycarbonyl)-2,3-dihydroindole (131 mg, 0.56 mmol) was added followed 10 minutes later by 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (150 mg, 0.47 mmol), (prepared as described for the starting material in Example 9). The reaction mixture was heated at 110 °C for 3 hours, cooled to ambient temperature and partitioned between ethyl acetate and water. The organic phase was washed with water, brine, dried (MgSO₄), filtered and the solvent evaporated. The residue was purified by flash chromatography using increasingly polar solvent mixtures starting with dichloromethane/methanol (90/10) and ending with dichloromethane/methanol/methanol saturated with ammonia (80/15/5). Evaporation of the solvent gave 4-(1-*tert*butoxycarbonyl)-2,3-dihydro-indol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (178 mg, 73%) as a white solid.

15 ¹H NMR Spectrum: (DMSO-d₆) 1.60 (s, 9H); 1.80 (m, 4H); 2.20 (m, 2H); 2.55 (m, 4H); 2.70 (t, 2H); 3.15 (t, 2H); 4.05 (br s, 5H); 4.30 (t, 2H); 7.00 (d, 1H); 7.05 (s, 1H); 7.30 (s, 1H); 7.55 (s, 1H); 7.90 (br s, 1H); 8.60 (s, 1H)

Example 194

Using an analogous procedure to that described in Example 192, 4-(1-*tert*butoxycarbonyl)-2,3-dihydro-indol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (191 mg, 0.37 mmol) was reacted with TFA (1 ml) to give, after work-up and purification, 4-(2,3-dihydro-indol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (103 mg, 67%).

25 ¹H NMR Spectrum: (CDCl₃) 1.65 (m, 2H); 2.00 (m, 3H); 2.25 (m, 2H); 2.45 (s, 3H); 3.10 (m, 4H); 3.65 (t, 2H); 4.05 (s, 3H); 4.10 (d, 2H); 6.70 (d, 1H); 6.85 (dd, 1H); 7.0 (s, 1H); 7.25 (s, 1H); 7.55 (s, 1H); 8.60 (s, 1H)

30 MS (ESI) : 421 [MH]⁺

The starting material was prepared as follows:

Using an analogous procedure to that described in Example 193, 4-chloro-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (150 mg, 0.47 mmol), (prepared as described for the starting material in Example 10), was reacted with 5-hydroxy-(1-*tert*butoxycarbonyl)-2,3-dihydroindole (132 mg, 0.56 mmol), (prepared as described for the starting material in Example 193), to give, after work-up and purification, 4-(1-*tert*butoxycarbonyl-2,3-dihydroindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (197 mg, 81%) as a white solid.

¹H NMR Spectrum: (CDCl₃) 1.50 (br s, 11H); 2.00 (m, 5H); 2.30 (s, 3H); 2.90 (d, 2H); 3.15 (t, 2H); 4.05 (br s, 7H); 7.05 (br s, 2H); 7.30 (s, 1H); 7.55 (s, 1H); 7.95 (br s, 1H); 8.60 (s, 1H)

Example 195

To a suspension of 4-chloro-6-methoxy-7-(2-piperidinoethoxy)quinazoline (250mg, 0.78mmol), (prepared as described for the starting material in Example 180), in DMF (10ml) was added anhydrous potassium carbonate (320mg, 2.30mmol) and 7-hydroxyquinoline (135mg, 0.94mmol), and the reaction heated under reflux at 90C for 1 hour. The reaction was cooled to ambient temperature and 1N aqueous sodium hydroxide added. The resulting precipitate was filtered, washed with water and acetone, and dried under suction to give 6-methoxy-7-(2-piperidinoethoxy)-4-(quinolin-7-yloxy)quinazoline (248mg, 0.58mmol, 75%) as a white solid.

¹H NMR Spectrum: *d*_H (300MHz, CDCl₃): 1.5 (2H, m; NCH₂CH₂CH₂), 1.6 (4H, m; 2 x NCH₂CH₂), 2.6 (4H, t; 2 x NCH₂), 2.9 (2H, t; NCH₂), 4.1 (3H, s; OCH₃), 4.3 (2H, t; OCH₂), 7.3 (1H, s; ArH), 7.4 (1H, dd; ArH), 7.5 (1H, dd; ArH), 7.6 (1H, s; ArH), 7.9 (1H, d; ArH), 8.0 (1H, d; ArH), 8.2 (1H, d; ArH), 8.6 (1H, s; ArH) and 8.9 (1H, dd; ArH)
m/z (ESP+) 431 (MH⁺, 100%)

Example 196

To a suspension of 7-benzyloxy-4-chloro-6-methoxyquinazoline (1.82g, 6.1mmol), (prepared as described for the starting material in Example 1), in DMF (50ml) was added potassium carbonate (2.50g, 18.1mmol) and 7-hydroxyquinoline (1.06g, 7.3mmol), and the reaction heated under reflux at 90C for 4 hours. The reaction was poured into 1N aqueous

sodium hydroxide and the resulting precipitate filtered, washed with water, and dried under suction. Further drying in a vacuum oven gave **7-benzyloxy-6-methoxy-4-(quinolin-7-yloxy)quinazoline** (1.50g, 3.7mmol, 60%) as a cream solid.

¹H NMR Spectrum: d_H (300MHz, DMSO- d_6): 4.0 (3H, s; OCH₃), 5.4 (2H, s; OCH₂), 7.3-7.7 (9H, m; 9 x ArH), 7.9 (1H, br s; ArH), 8.1 (1H, d; ArH), 8.4 (1H, d; ArH), 8.5 (1H, s; ArH) and 8.9 (1H, d; ArH)

Example 197

A solution of **7-benzyloxy-6-methoxy-4-(quinolin-7-yloxy)quinazoline** (1.50g, 3.70mmol), (prepared as described in Example 196), in trifluoroacetic acid (50ml) was heated at reflux for 150 minutes. The reaction was concentrated *in vacuo* and the reaction neutralised with saturated aqueous ammonium hydroxide. The resulting precipitate was filtered, washed with acetone and dried under suction to give **7-hydroxy-6-methoxy-4-(quinolin-7-yloxy)quinazoline** (0.90g, 2.82mmol, 76%) as a white solid.

¹H NMR Spectrum: d_H (300MHz, DMSO- d_6): 4.0 (3H, s; OCH₃), 7.1 (1H, s; ArH), 7.3-7.4 (3H, m; 3 x ArH), 7.9 (1H, br s; ArH), 8.1 (1H, d; ArH), 8.4-8.5 (2H, d; 2 x ArH) and 8.9 (1H, d; ArH)

m/z (ESP+) 320 (MH⁺, 100%)

Example 198

To a suspension of **7-hydroxy-6-methoxy-4-(quinolin-7-yloxy)quinazoline** (450mg, 1.40mmol), (prepared as described in Example 197), in DMF (50ml) was added anhydrous potassium carbonate (773mg, 5.60mmol) and 4-(2-hydroxyethyl)morpholine (335mg, 1.80mmol), and the reaction heated under reflux for 2 hours. The DMF was evaporated *in vacuo*, and the residue partitioned between dichloromethane and 1N aqueous sodium hydroxide. The mixture was extracted with dichloromethane (3 x 200ml), dried (MgSO₄) and concentrated *in vacuo*. The crude product was triturated with hexane/ether to afford a solid which was filtered and dried under suction to give **6-methoxy-7-(2-morpholinoethoxy)-4-(quinolin-7-yloxy)quinazoline** (430mg, 1.00mmol, 71%) as a light brown solid.

¹H NMR Spectrum: d_H (300MHz, CDCl₃): 2.7 (4H, t; 2 x NCH₂), 3.0 (2H, t; NCH₂), 3.7 (4H, t; 2 x OCH₂), 4.1 (3H, s; OCH₃), 4.4 (2H, t; OCH₂), 7.2 (1H, s; ArH), 7.4 (1H, dd; ArH), 7.5

(1H, dd; ArH), 7.6 (1H, s; ArH), 7.9 (1H, d; ArH), 8.0 (1H, br s; ArH), 8.2 (1H, d; ArH), 8.6 (1H, s; ArH) and 8.9 (1H, dd; ArH)

m/z (ESP+) 433 (MH^+ , 100%)

Elemental analysis	Found	C 65.0	H 5.6	N 12.6
--------------------	-------	--------	-------	--------

5	$C_{24}H_{24}N_4O_4 \cdot 0.5H_2O$	Requires	C 65.3	H 5.7	N 12.7%
---	------------------------------------	----------	--------	-------	---------

Example 199

To a solution of 7-hydroxy-6-methoxy-4-(quinolin-7-yloxy)quinazoline (100mg, 0.31mmol), (prepared as described in Example 197), and (S)-(+)-5-(hydroxymethyl)-2-pyrrolidinone (101mg, 0.47mmol) in dichloromethane (10ml) was added triphenylphosphine (244mg, 0.93mmol) and DEAD (0.15ml, 162mg, 0.93mmol), and the reaction stirred at ambient temperature overnight. The reaction mixture was placed directly onto a 2g SCX ion-exchange column, and eluted with dichloromethane, then dichloromethane/methanol (4/1), then dichloromethane/methanol/ammonium hydroxide (20/5/1). The appropriate fractions were concentrated *in vacuo*, and the residue triturated with ether to give a solid which was filtered and dried under suction to give (5S)-6-methoxy-7-(5-oxo-pyrrolidin-2-ylmethoxy)-4-(quinolin-7-yloxy)quinazoline (55mg, 0.13mmol, 43%) as a yellow solid.

1H NMR Spectrum: d_H (300MHz, $CDCl_3$): 2.3-2.5 (4H, m; 2 x pyrrolidinone- CH_2), 4.0-4.1 (4H, m; pyrrolidinone- CH ; OCH_3), 4.2-4.3 (2H, m; OCH_2), 6.1 (1H, br s; NH), 7.3 (1H, s; ArH), 7.4 (1H, dd; ArH), 7.5 (1H, dd; ArH), 7.9 (1H, d; ArH), 8.0 (1H, br s; ArH), 8.2 (1H, d; ArH), 8.6 (1H, s; ArH) and 8.9 (1H, dd; ArH)

m/z (ESP+) 417 (MH^+ , 100%)

Example 200

To a solution of 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (100mg, 0.31mmol), (prepared as described for the starting material in Example 9), in DMF (10ml) was added potassium carbonate (124mg, 0.9mmol, 3eq.) followed by 2-hydroxycarbazole (66mg, 0.36mmol, 1.2eq.) and the reaction heated at 100°C for 4 hours. The DMF was removed *in vacuo*, the residue dissolved in dichloromethane and placed onto a 2g SCX ion-exchange column. Elution with dichloromethane, followed by 20% methanol/dichloromethane then 20% methanol/dichloromethane + 3% ammonium hydroxide, gave the crude product as a brown solid. Further purification by silica bond elut

chromatography eluting with dichloromethane to 15% methanol/dichloromethane + 1% ammonium hydroxide, followed by trituration with ether gave **4-(9H-carbazol-2-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (31mg, 22%) as a white solid.

¹H NMR Spectrum: *d*_H (300MHz, DMSO-*d*₆) 1.7 (4H, m; 2 x pyrrolidine-CH₂), 2.0 (2H, t; OCH₂CH₂), 2.5 (4H, m; 2 x pyrrolidine-NCH₂), 2.6 (2H, t; NCH₂), 4.0 (3H, s; OCH₃), 4.2 (2H, t; OCH₂), 7.1 (1H, br d; ArH), 7.2 (1H, t; ArH), 7.3-7.4 (3H, m; 3 x ArH), 7.5 (1H, br d; ArH), 7.6 (1H, s; ArH), 8.1-8.2 (2H, m; 2 x ArH), 8.5 (1H, s; ArH), 11.3 (1H, s; carbazole NH)

m/z (ESP+) 469 (MH⁺, 100%)

Example 201

To a solution of 7-hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg, 0.32 mmol), 2-((N-(3,6-dichloropyridazin-4-yl)-N-methyl)amino)ethanol (107 mg, 0.48 mmol), (prepared as described for the starting material in Example 142), triphenylphosphine (168 mg, 0.64 mmol) in methylene chloride (1 ml) and DMF (0.5 ml) cooled at 4°C was added a solution of diethyl azodicarboxylate (101 µl, 0.64 mmol) in methylene chloride (0.4 ml). The mixture was stirred for 12 hours at 4°C and overnight at ambient temperature. The precipitate was filtered, washed with ether and dried under vacuum to give **7-(2-((N-(3,6-dichloropyridazin-4-yl)-N-methyl)amino)ethoxy)-4-(indol-5-ylamino)-6-methoxyquinazoline** (72 mg, 44 %).

MS-ESI : 510-512 [MH]⁺

¹H NMR Spectrum: (DMSO-*d*₆) 3.12 (s, 3H) ; 3.85 (s, 3H) ; 4.1 (t, 2H) ; 4.45 (t, 2H) ; 6.45 (s, 1H) ; 7.2 (s, 1H) ; 7.3 (s, 1H) ; 7.35 (m, 2H) ; 7.42 (d, 1H) ; 7.8 (s, 1H) ; 7.85 (s, 1H) ; 8.35 (s, 1H) ; 9.45 (s, 1H)

The starting material was prepared as follows :

A solution of 7-benzyloxy-4-chloro-6-methoxyquinazoline (5g, 16.6 mmol), (prepared as described for the starting material in Example 1), 5-aminoindole (2.4 g, 18.2 mmol) in isopropanol (60 ml) containing 5N hydrogen chloride in isopropanol (260 µl, 1.6 mmol) was refluxed for 90 minutes. After cooling the volatiles were removed under vacuum. The solid was triturated with isopropanol, filtered, washed with isopropanol followed by ether and dried

under vacuum to give 7-benzyloxy-4-(indol-5-ylamino)-6-methoxyquinazoline hydrochloride (6.9g, 96%).

¹H NMR Spectrum: (DMSO-d₆) 4.05 (s, 3H) ; 5.35 (s, 2H) ; 6.5 (s, 1H) ; 7.3 (d, 1H) ; 7.4-7.65 (m, 9H) ; 7.8 (s, 1H) ; 8.3 (s, 1H) ; 8.7 (s, 1H)

- 5 A solution of give 7-benzyloxy-4-(indol-5-ylamino)-6-methoxyquinazoline hydrochloride (10g, 23.1 mmol) in methanol (300ml) and DMF (100ml) containing ammonium formate (22gr, 347 mmol) and 10% palladium on charcoal (1g) was stirred overnight at ambient temperature. The solution was filtered over celite and washed with DMF followed by methanol. The filtrate was evaporated. The residue was dissolved in aqueous
- 10 ammonia 2mM (300ml) and stirred for 15 minutes. The solid was filtered, washed with water followed by ethyl acetate and ether and dried under vacuum at 50°C for 2 days. The solid was purified by column chromatography eluting with methanol/methylene chloride (1/9). The volatiles were removed under vacuum and the solid was left under vacuum at 70°C for 2 days to give 7-hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (6.8 g, 97%)


15 MS-ESI : 307 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 3.98 (s, 3H) ; 6.42 (s, 1H) ; 7.0 (s, 1H) ; 7.3-7.45 (m, 3H) ; 7.85 (s, 2H) ; 8.28 (s, 1H) ; 9.35 (s, 1H) ; 10.25 (br s, 1H) ; 11.05 (s, 1H)

Examples 202-204




- 20 Using an analogous procedure to that described in Example 201, 7-hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline, (prepared as described for the starting material in Example 201), was used in the synthesis of the compounds described in Table XI.

Table XI



25

Example number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	Note	R
----------------	-------------	---------	--------------------------	------	---

202	83	59	441	a	
203	91	72	398	b	
204	76	55	432	c	

a) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline was reacted with 2-(N-methyl-N-(4-pyridyl)amino)ethanol (73mg), (EP 0359389), to give **4-(indol-5-ylamino)-6-methoxy-7-(2-(N-methyl-N-(4-pyridyl)amino)ethoxy)quinazoline**.

- 5 ¹H NMR Spectrum: (DMSO-d₆) 3.08 (s, 3H) ; 3.9 (t, 2H) ; 3.95 (s, 3H) ; 4.35 (t, 2H) ; 6.45 (s, 1H) ; 6.75 (d, 2H) ; 7.15 (s, 1H) ; 7.35 (m, 2H) ; 7.4 (d, 1H) ; 7.85 (s, 1H) ; 7.9 (s, 1H) ; 8.15 (d, 2H) ; 8.38 (s, 1H) ; 9.45 (s, 1H)

- 10 b) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline was reacted with 3-hydroxymethyl pyridine (53 mg) to give **4-(indol-5-ylamino)-6-methoxy-7-((3-pyridyl)methoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 4.0 (s, 3H) ; 5.35 (s, 2H) ; 6.42 (s, 1H) ; 7.3-7.55 (m, 5H) ; 7.8-8.0 (m, 3H) ; 8.4 (s, 1H) ; 8.6 (d, 1H) ; 8.75 (s, 1H) ; 9.5 (s, 1H)

- 15 c) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline was reacted with 5-(2-hydroxyethyl)-4-methylthiazole (69 mg) to give **4-(indol-5-ylamino)-6-methoxy-7-(2-(4-methyl-1,3-thiazol-5-yl)ethoxy)quinazoline**.

- 20 ¹H NMR Spectrum: (DMSO-d₆) 2.45 (s, 3H) ; 3.32 (t, 2H) ; 3.95 (s, 3H) ; 4.32 (t, 2H) ; 6.45 (s, 1H) ; 7.15 (s, 1H) ; 7.3-7.45 (m, 3H) ; 7.85 (s, 1H) ; 7.9 (s, 1H) ; 8.35 (s, 1H) ; 8.85 (s, 1H) ; 9.45 (s, 1H)

Example 205

To a solution of 7-hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (102 mg, 0.32 mmol), 4-(3-hydroxypropyl)morpholine (70 mg, 0.48 mmol), (prepared as described for the starting material in Example 60), triphenylphosphine (168 mg, 0.64 mmol) in methylene chloride (1 ml) and DMF (0.5 ml) cooled at 4°C was added a solution of diethyl azodicarboxylate (101 µl, 0.64 mmol) in methylene chloride (0.4 ml). The mixture was stirred for 12 hours at 4°C and overnight at ambient temperature. The mixture was poured onto a column of silica (IST isolate ® 10 g of silica) and was eluted with methylene chloride (15 ml) followed by 5 % methanol in methylene chloride (45 ml) followed by 5 % methanol (saturated with ammonia) in methylene chloride (30 ml) followed by 10 % methanol (saturated with ammonia) in methylene chloride (45 ml) followed by 15 % methanol (saturated with ammonia) in methylene chloride (30 ml). The fractions containing the expected product were evaporated to give **6-methoxy-4-(2-methylindol-5-ylamino)-7-(3-morpholinopropoxy)quinazoline** (63 mg, 44 %).

MS-ESI : 448 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.0 (m, 2H) ; 2.4 (s, 3H) ; 2.3-2.6 (m, 6H) ; 3.6 (t, 4H) ; 3.95 (s, 3H) ; 4.2 (t, 2H) ; 6.12 (s, 1H) ; 7.12 (s, 1H) ; 7.3 (br s, 2H) ; 7.7 (s, 1H) ; 7.85 (s, 1H) ; 8.35 (s, 1H) ; 9.4 (s, 1H)

The starting material was prepared as follows:

A solution of 2-methyl-5-nitroindole (1 g, 5.7 mmol) in ethanol (25ml) and THF (25 ml) containing 10% palladium on charcoal (128mg) was hydrogenated until uptake of hydrogen ceased. The mixture was filtered and the filtrate was evaporated to give 5-amino-2-methylindole (830 mg, quant.).

¹H NMR Spectrum: (DMSO-d₆) 2.3 (s, 3H) ; 4.3 (br s, 2H) ; 5.8 (s, 1H) ; 6.35 (d, 1H) ; 6.55 (s, 1H) ; 6.95 (d, 1H) ; 10.35 (br s, 1H)

Using an analogous procedure to that described for the synthesis of the starting material in Example 201, 7-benzyloxy-4-chloro-6-methoxyquinazoline (2 g, 6.6 mmol), (prepared as described for the starting material in Example 1), was reacted with 5-amino-2-methylindole (1.07g, 7.3 mmol) to give 7-benzyloxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline hydrochloride (2.9 g, quanti.).

MS-ESI : 411 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.41 (s, 3H) ; 4.01 (s, 3H) ; 5.33 (s, 2H) ; 6.18 (s, 1H) ; 7.25 (d, 1H) ; 7.3-7.7 (m, 8H) ; 8.3 (s, 1H) ; 8.7 (s, 1H) ; 11.1 (s, 1H) ; 11.4 (s, 1H)

Using an analogous procedure to that described for the synthesis of the starting material in Example 201, 7-benzyloxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline hydrochloride (2.87g, 6.4 mmol) was reacted with ammonium formate (6g, 9.6 mmol) to give 7-hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (1.91g, 93%).


MS-ESI : 321 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H) ; 3.95 (s, 3H) ; 6.12 (s, 1H) ; 7.0 (s, 1H) ; 7.25 (s, 1H) ; 7.7 (s, 1H) ; 7.85 (s, 1H) ; 8.3 (s, 1H) ; 9.35 (s, 1H) ; 10.2 (br s, 1H) ; 10.9 (s, 1H)

Examples 206-207

Using an analogous procedure to that described for Example 205, 7-hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline, (prepared as described for the starting material in Example 205), was used in the synthesis of the compounds described in Table XII.

Table XII



Example Number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	Note	R
206	65	41	496	a	
207	62	45		b	

a) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (98 mg) was reacted with

3-(1,1-dioxothiomorpholino)-1-propanol (93 mg), (prepared as described for the starting

material in Example 5), to give **7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.0 (m, 2H) ; 2.4 (s, 3H) ; 2.7 (t, 2H) ; 2.95 (m, 4H) ; 3.15 (m, 4H) ; 3.95 (s, 3H) ; 4.2 (t, 2H) ; 6.15 (s, 1H) ; 7.18 (s, 1H) ; 7.28 (m, 2H) ; 7.7 (s, 1H) ; 7.85 (s, 1H) ; 8.35 (s, 1H) ; 9.4 (s, 1H)

b) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (98 mg) was reacted with 1-(2-hydroxyethyl)piperidine (62 mg) to give **6-methoxy-4-(2-methylindol-5-ylamino)-7-(2-piperidinoethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.4 (m, 2H) ; 1.45-1.6 (m, 4H) ; 2.42 (s, 3H) ; 2.45 (br s, 4H) ; 2.75 (t, 2H) ; 3.95 (s, 3H) ; 4.25 (t, 2H) ; 6.15 (s, 1H) ; 7.15 (s, 1H) ; 7.25 (br s, 2H) ; 7.7 (s, 1H) ; 7.88 (s, 1H) ; 8.35 (s, 1H) ; 9.4 (s, 1H)

Example 208

Using an analogous procedure to that described in Example 205, 7-hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg, 0.32 mmol), (prepared as described for the starting material in Example 201), was reacted with 3-(1,2,3-triazol-1-yl)propan-1-ol (61 mg, 0.48 mmol) to give **4-(indol-5-ylamino)-6-methoxy-7-(3-(1,2,3-triazol-1-yl)propoxy)quinazoline** (56 mg, 42 %).

MS-ESI : 416 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.4 (m, 2H) ; 4.0 (s, 3H) ; 4.2 (t, 2H) ; 4.65 (t, 2H) ; 6.45 (s, 1H) ; 7.15 (s, 1H) ; 7.35 (m, 2H) ; 7.42 (d, 1H) ; 7.75 (s, 1H) ; 7.88 (s, 1H) ; 7.9 (s, 1H) ; 8.2 (s, 1H) ; 8.38 (s, 1H) ; 9.42 (s, 1H)

The starting material was prepared as follows:

A mixture of 1,2,3-triazole (5g, 72.4 mmol) and ethyl acrylate (7.8 ml, 72.4 mmol) containing pyridine (50 drops) was heated at 90°C for 4 hours. After cooling, the volatiles were removed under vacuum and the residue was purified by column chromatography eluting with methylene chloride/ether to give ethyl (1H-1,2,3-triazol-1-yl)propanoate (8.96g, 73%).

¹H NMR Spectrum: (CDCl₃) 1.25 (t, 3H) ; 2.95 (t, 2H) ; 4.15 (q, 2H) ; 4.7 (t, 2H) ; 7.65 (s, 1H) ; 7.7 (s, 1H)


A solution of ethyl (1*H*-1,2,3-triazol-1-yl)propanoate (8.96g, 53 mmol) in THF (50ml) was added dropwise to a suspension of lithium aluminium hydride (3 g, 79 mmol) in THF (250ml) cooled at 0°C. After stirring for 1 hour at 5°C, the mixture was stirred for 1 hour at ambient temperature. The mixture was cooled at 0°C and 4N sodium hydroxide (30ml) was added dropwise. The mixture was filtered and the solid was washed with THF followed by ethyl acetate. The filtrate was dried (MgSO₄) and evaporated. The residue was purified by column chromatography, eluting with methylene chloride/methanol (94/6) to give 3-(1,2,3-triazol-1-yl)propan-1-ol (6.2 g, 92%).

¹H NMR Spectrum: (CDCl₃) : 2.1-2.2 (m, 3H) ; 3.65 (m, 2H) ; 4.6 (t, 2H) ; 7.6 (s, 1H) ; 7.72 (s, 1H)







Examples 209–216

Using an analogous procedure to that described in Example 208, 7-hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline, (prepared as described for the starting material in Example 201), was used in the synthesis of the compounds described in Table XIII.

Table XIII



Example Number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	Note	R
209	77	57	422	a	
210	64	45	446	b	

211	76	49	482	c	
212	70	48	462	d	
213	85	59	447	e	
214	62	54	365	f	
215	71	54	409	g	
216	73	55	418	h	

a) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 2-(N-(2-methoxyethyl)-N-methylamino)ethanol (64 mg), (prepared as described for the starting material in Example 59), to give **4-(indol-5-ylamino)-6-methoxy-7-(2-(N-(2-methoxyethyl)-N-methylamino)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.35 (s, 3H) ; 2.68 (t, 2H) ; 2.82 (t, 2H) ; 3.25 (s, 3H) ; 3.5 (t, 2H) ; 3.97 (s, 3H) ; 4.22 (t, 2H), 6.45 (s, 1H) ; 7.18 (s, 1H) ; 7.3-7.45 (m, 3H) ; 7.88 (m, 2H) ; 8.35 (s, 1H) ; 9.42 (s, 1H)

10 b) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 1-(3-hydroxypropyl)pyrrolidin-2,5-dione (76 mg) to give **7-(3-(2,5-dioxopyrrolidin-1-yl)propoxy)-4-(indol-5-ylamino)-6-methoxyquinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.05 (m, 2H) ; 2.65 (s, 3H) ; 3.6 (t, 2H) ; 3.98 (s, 2H) ; 4.15 (t, 2H) ; 6.45 (s, 1H) ; 7.1 (s, 1H) ; 7.3-7.45 (m, 3H) ; 8.7 (s, 1H) ; 8.8 (s, 1H) ; 8.35 (s, 1H) ;

15 9.45 (s, 1H)

The starting material was prepared as follows:

A solution of pyrrolidine-2,5-dione (5g, 50.5 mmol) and 3-bromopropan-1-ol (6.85 ml, 76 mmol) in acetonitrile (80ml) containing potassium carbonate (14g, 100 mmol) was refluxed overnight. After cooling, the mixture was filtered and the filtrate was evaporated.

- 5 The residue was dissolved in methylene chloride and purified by column chromatography, eluting with ethylacetate/petroleum ether (4/1). After evaporation of the volatiles, the residue was distilled at 100-125°C under about 0.1 mm Hg to give 1-(3-hydroxypropyl)pyrrolidin-2,5-dione (2.6 g, 34%).

¹H NMR Spectrum: (CDCl₃) 1.8 (m, 2H) ; 2.52 (t, 1H) ; 2.78 (s, 4H) ; 3.58 (q, 2H) ; 3.7 (t, 10 2H)

- c) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 3-(1,1-dioxothiomorpholino)-1-propanol (93mg), (prepared as described for the starting material in Example 5), to give **7-(3-(1,1-dioxothiomorpholino)propoxy)-4-(indol-5-ylamino)-6-methoxyquinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.0 (m, 2H) ; 2.7 (t, 2H) ; 2.95 (br s, 4H) ; 3.15 (br s, 4H) ; 3.97 (s, 3H) ; 4.2 (t, 2H) ; 6.45 (s, 1H) ; 7.2 (s, 1H) ; 7.3-7.5 (m, 3H) ; 7.9 (2s, 2H) ; 8.35 (s, 1H) ; 9.42 (s, 1H)

- 20 d) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 3-((4-methyl-4*H*-1,2,4-triazol-3-yl)sulphanyl)propan-1-ol (83 mg) to give **4-(indol-5-ylamino)-6-methoxy-7-(3-((4-methyl-4*H*-1,2,4-triazol-3-yl)sulphanyl)propoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.22 (m, 2H) ; 3.3 (m, 2H) ; 3.65 (s, 3H) ; 3.95 (s, 3H) ; 4.25 (t, 2H) ; 6.45 (s, 1H) ; 7.15 (s, 1H) ; 7.3-7.45 (m, 3H) ; 7.88 (s, 1H) ; 8.0 (s, 1H) ; 8.35 (s, 1H) ; 8.58 (s, 1H) ; 9.45 (s, 1H)

The starting material was prepared as follows:

- A solution of 4-methyl-4-*H*-1,2,4-triazole-3-thiol (1.72g, 15 mmol) and 3-bromopropan-1-ol (1.39g, 10 mmol) in DMF (10ml) containing potassium carbonate (1.57g, 14 mmol) was heated at 40°C for 30 minutes. The mixture was then partitioned between saturated ammonium chloride and ethyl acetate. The aqueous layer was evaporated to dryness and the residue was triturated with ethyl acetate and methylene chloride. The suspension was

filtered and the filtrate was dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with methylene chloride/methanol (9/1) to give 3-((4-methyl-4*H*-1,2,4-triazol-3-yl)sulphanyl)propan-1-ol (510mg, 30%).

¹H NMR Spectrum: (CDCl₃) 2.02 (m, 2H); 3.45 (t, 2H); 3.55 (s, 3H); 3.75 (t, 2H); 8.15 (s, 1H)

e) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 1-(3-hydroxypropyl)-4-methylpiperazine (76 mg), (prepared as described for the starting material in Example 133), to give **4-(indol-5-ylamino)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.0 (m, 2H); 2.2 (s, 3H); 2.25-2.55 (m, 10H); 4.0 (s, 3H); 4.2 (t, 2H); 6.45 (s, 1H); 7.15 (s, 1H); 7.35 (m, 2H); 7.42 (d, 1H); 7.88 (br s, 2H); 8.38 (s, 1H); 9.42 (s, 1H)

f) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 2-methoxyethanol (37 mg) to give **4-(indol-5-ylamino)-6-methoxy-7-(2-methoxyethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 3.4 (s, 3H); 3.75 (t, 2H); 3.98 (s, 3H); 4.38 (t, 2H); 6.45 (s, 1H); 7.18 (s, 1H); 7.35 (m, 2H); 7.42 (d, 1H); 7.85 (s, 1H); 7.9 (s, 1H); 8.38 (s, 1H); 9.5 (s, 1H)

g) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 2-(2-methoxyethoxy)ethanol (58 mg) to give **4-(indol-5-ylamino)-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 3.3 (s, 3H); 3.5 (t, 2H); 3.65 (t, 2H); 3.85 (t, 2H); 4.0 (s, 3H); 4.28 (t, 2H); 6.45 (s, 1H); 7.18 (s, 1H); 7.35 (m, 2H); 7.45 (d, 1H); 7.88 (s, 1H); 7.9 (s, 1H); 8.35 (s, 1H); 9.45 (s, 1H)

h) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 1-(2-hydroxyethyl)piperidine (62 mg) to give **4-(indol-5-ylamino)-6-methoxy-7-(2-piperidinoethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.6 (m, 6H) ; 2.5 (br s, 4H) ; 2.7 (t, 2H) ; 3.98 (s, 3H) ; 4.25 (t, 2H) ; 6.45 (s, 1H) ; 7.18 (s, 1H) ; 7.35 (m, 2H) ; 7.42 (d, 1H) ; 7.9 (br s, 2H) ; 8.38 (s, 1H) ; 9.42 (s, 1H)

5 Example 217-223

Using an analogous procedure to that described in Example 205, 7-hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline was used in the synthesis of the compounds described in Table XIV.

10 The starting material was prepared as follows:

Using an analogous procedure to that described for the preparation of the starting material in Example 201, 6-nitroindole (500mg, 3 mmol) was hydrogenated to give 6-aminoindole (395mg, quant.).

¹H NMR Spectrum: (DMSO-d₆) 6.41 (s, 1H) ; 6.6 (dd, 1H) ; 6.63 (s, 1H) ; 7.0 (t, 1H) ; 7.4 (d, 15 1H) ; 7.87 (br s, 1H)

Using an analogous procedure to that described for the preparation of the starting material in Example 201, 7-benzyloxy-4-chloro-6-methoxyquinazoline (2.5 g, 8.3 mmol), (prepared as described for the starting material in Example 1), was reacted with 6-aminoindole (1.5g, 11.4 mmol) to give 7-benzyloxy-4-(indol-6-ylamino)-6-methoxyquinazoline 20 hydrochloride (3.18g, 89%).

MS-ESI : 397 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 4.02 (s, 3H) ; 5.35 (s, 2H) ; 6.5 (s, 1H) ; 7.25 (dd, 1H) ; 7.35-7.6 (m, 5H) ; 7.63 (d, 1H) ; 7.72 (s, 1H) ; 8.3 (s, 1H) ; 8.75 (s, 1H) ; 11.3 (br s, 1H)


Using an analogous procedure to that described for the preparation of the starting material in Example 201, 7-benzyloxy-4-(indol-6-ylamino)-6-methoxyquinazoline 25 hydrochloride was treated with ammonium formate (655mg, 10.4 mmol) to give 7-hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (162 mg, 76%).

MS-ESI : 307 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 4.0 (s, 3H) ; 6.4 (s, 1H) ; 7.0 (s, 1H) ; 7.3 (m, 2H) ; 7.5 (d, 1H) 30 ; 7.85 (s, 1H) ; 8.0 (s, 1H) ; 8.35 (s, 1H) ; 9.35 (s, 1H) ; 11.05 (s, 1H)

Table XIV

- 235 -



Example number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	Note	R
217	46	35	416	a	
218	57	37	482	b	
219	37	25	462	c	
220	38	29	418	d	
221	10	7	418	e	
222	94	61	483	f	
223	56	44	398	g	

a) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 3-(1,2,3-triazol-1-yl)propan-1-ol (61 mg), (prepared as described for the starting material in Example 208), to give **4-(indol-6-ylamino)-6-methoxy-7-(3-(1,2,3-triazol-1-yl)propoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.42 (t, 2H) ; 4.02 (s, 3H) ; 4.2 (t, 2H) ; 4.62 (t, 2H) ; 6.42 (s, 1H) ; 7.15 (s, 1H) ; 7.3 (m, 2H) ; 7.55 (d, 1H) ; 7.75 (s, 1H) ; 7.92 (s, 1H) ; 8.02 (s, 1H) ; 8.2 (s, 1H) ; 8.42 (s, 1H) ; 9.45 (s, 1H)

b) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 3-(1,1-dioxothiomorpholino)-1-propanol (93mg), (prepared as described for the starting material in Example 5), to give **7-(3-(1,1-dioxothiomorpholino)propoxy)-4-(indol-6-ylamino)-6-methoxyquinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.0 (m, 2H) ; 2.7 (t, 2H) ; 2.95 (br s, 4H) ; 3.12 (br s, 4H) ; 4.0 (s, 3H) ; 4.2 (t, 2H) ; 6.42 (s, 1H) ; 7.2 (s, 1H) ; 7.3 (m, 2H) ; 7.55 (d, 1H) ; 7.9 (s, 1H) ; 8.02 (s, 1H) ; 8.42 (s, 1H) ; 9.48 (s, 1H)

c) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 3-((4-methyl-4*H*-1,2,4-triazol-3-yl)sulphanyl)propan-1-ol (83 mg), (prepared as described for the starting material in Example 212), to give **4-(indol-6-ylamino)-6-methoxy-7-(3-((4-methyl-4*H*-1,2,4-triazol-3-yl)sulphanyl)propoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.22 (t, 2H) ; 3.3 (t, 2H) ; 3.6 (s, 3H) ; 4.0 (s, 3H) ; 4.28 (t, 2H) ; 6.4 (s, 1H) ; 7.18 (s, 1H) ; 7.3 (m, 2H) ; 7.53 (d, 1H) ; 7.9 (s, 1H) ; 8.02 (s, 1H) ; 8.42 (s, 1H) ; 8.58 (s, 1H) ; 9.45 (s, 1H)

d) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 1-(2-hydroxyethyl)piperidine (62 mg) to give **4-(indol-6-ylamino)-6-methoxy-7-(2-piperidinoethoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.6 (m, 6H) ; 2.5 (br s, 4H) ; 2.75 (t, 2H) ; 4.0 (s, 3H) ; 4.25 (t, 2H) ; 6.42 (s, 1H) ; 7.2 (s, 1H) ; 7.3 (m, 2H) ; 7.55 (d, 1H) ; 7.9 (s, 1H) ; 8.02 (s, 1H) ; 8.42 (s, 1H) ; 9.45 (s, 1H)

e) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 1-(3-hydroxypropyl)pyrrolidine (62 mg) to give **4-(indol-6-ylamino)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline**.

5 The starting material was prepared as follows:

A solution of pyrrolidine (50 g, 0.7 mol) and 3-chloropropan-1-ol (66.15 g, 0.7 mol) in acetonitrile (1 l) containing potassium carbonate (145 g, 1.05 mol) was refluxed for 20 hours. After cooling, the mixture was filtered, the solid was washed with acetonitrile and the filtrate was evaporated. The residue was distilled at about 130°C under about 70 mmHg to give 1-(3-hydroxypropyl)pyrrolidine (62.1 g, 69%).

MS-ESI : 130 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 1.6-1.8 (m, 6H) ; 2.55 (br s, 4H) ; 2.75 (t, 2H) ; 3.85 (t, 2H) ; 5.2-5.8 (br s, 1H)

15 f) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 3-((N-(2,6-dimethyl-4-pyridyl)-N-methyl)amino)propan-1-ol (93mg) to give **7-(3-((N-(2,6dimethyl-4-pyridyl)-N-methyl)amino)propoxy)-4-(indol-6-ylamino)-6-methoxyquinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.08 (m, 2H) ; 2.22 (s, 6H) ; 2.95 (s, 3H) ; 3.6 (t, 2H) ; 4.05 (s, 3H) ; 4.15 (t, 2H) ; 6.35 (s, 2H) ; 6.42 (s, 1H) ; 7.15 (s, 1H) ; 7.3 (m, 2H) ; 7.55 (d, 1H) ; 7.92
20 (s, 1H) ; 8.02 (s, 1H) ; 8.4 (s, 1H) ; 9.45 (s, 1H)

The starting material was prepared as follows:

A solution of 4-chloro-2,6-dimethylpyridine (2.12 g, 15 mmol) and 3-(N-methylamino)-propan-1-ol (4g, 45 mmol) containing 2N hydrogen chloride in ether (10 drops)
25 was heated at 140°C for 1 hour. The mixture was diluted with water (10ml) and poured onto a suspension of MgSO₄ (125g) in ethyl acetate (200ml). The mixture was filtered. The filtrate was evaporated and the residue was triturated with ether. The solid was filtered and dried under vacuum to give 3-((N-(2,6-dimethyl-4-pyridyl)-N-methyl)amino)propan-1-ol (1.76 g, 61%).

30 MS-EI : 194 [M]⁺

¹H NMR Spectrum: (CDCl₃) 1.75-1.95 (m, 2H) ; 2.4 (s, 6H) ; 3.0 (s, 3H) ; 3.48 (t, 2H) ; 3.7 (t, 2H) ; 6.25 (s, 2H)

g) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 3-hydroxymethyl pyridine (53 mg) to give **4-(indol-6-ylamino)-6-methoxy-7-((3-pyridyl)methoxy)quinazoline**.

- 5 ¹H NMR Spectrum: (DMSO-d₆) 4.02 (s, 3H) ; 5.35 (s, 2H) ; 6.42 (s, 1H) ; 7.22-7.4 (m, 3H) ; 7.5 (m, 1H) ; 7.55 (d, 1H) ; 7.95 (s, 1H) ; 7.97 (d, 1H) ; 8.0 (s, 1H) ; 8.42 (s, 1H) ; 8.6 (d, 1H) ; 8.78 (s, 1H) ; 9.5 (s, 1H)

Example 224

- 10 Using an analogous procedure to that described in Example 208, 7-hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline (98 mg, 0.32 mmol), (prepared as described for the starting material in Example 201), was reacted with (E)-4-(pyrrolidin-1-yl)but-2-en-1-ol (68 mg, 0.48 mmol), (prepared as described for the starting material in Example 129). After evaporation of the fractions containing the expected product, the residue was triturated with isopropanol (1
15 ml) containing 6.2 N hydrogen chloride in isopropanol (100 µl). After stirring at ambient temperature for 10 minutes, ether (500 µl) was added. The precipitate was filtered and washed several times with ether to give **4-(indol-5-ylamino)-6-methoxy-7-((E)4-(pyrrolidin-1-yl)but-2-en-1-yloxy)quinazoline hydrochloride** (14 mg, 10 %).

MS-ESI : 430 [MH]⁺


- 20 ¹H NMR Spectrum: (DMSO-d₆) 1.85-2.7 (br s, 4H) ; 2.95-3.1 (br s, 2H) ; 3.0 (m, 2H) ; 3.4-3.5 (m, 2H) ; 3.8 (d, 2H) ; 4.0 (s, 3H) ; 4.8 (d, 2H) ; 6.0-6.3 (m, 2H) ; 6.5 (s, 1H) ; 7.2-7.53 (m, 4H) ; 7.75 (s, 1H) ; 8.25 (s, 1H) ; 8.8 (br s, 1H)

Example 225

- 25 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline, (prepared as described for the starting material in Example 201), was treated as follows. After purification by chromatography and evaporation of the solvent, the residue was triturated in a solution of isopropanol (1 ml) containing 6.2 N hydrogen chloride in isopropanol (100 µl). After stirring for 10 minutes at ambient temperature, ether (500 µl) was added. The solid was filtered and
30 dried under vacuum to give 7-hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline hydrochloride.

Using an analogous procedure to that described in Example 224, 7-hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline hydrochloride was used in the synthesis of the compounds described in Table XV.

5 **Table XV**



Example Number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	Note	R
225	77	50	483	a	

a) 7-Hydroxy-4-(indol-5-ylamino)-6-methoxyquinazoline hydrochloride (98 mg) was reacted with 3-((N-(2,6-dimethyl-4-pyridyl)-N-methyl)amino)propan-1-ol (93mg), (prepared as described for the starting material in Example 222), to give 7-(3-((N-(2,6-dimethyl-4-pyridyl)-N-methyl)amino)propoxy)-4-(indol-5-ylamino)-6-methoxyquinazoline.

¹H NMR Spectrum: (DMSO-d₆) 2.2 (m, 2H) ; 2.5 (2br s, 6H) ; 3.2 (s, 3H) ; 3.8 (t, 2H) ; 4.1 (s, 3H) ; 4.25 (t, 2H) ; 6.52 (s, 1H) ; 6.75 (br s, 1H) ; 6.9 (br s, 1H) ; 7.35 (dd, 1H) ; 7.45 (br s, 2H) ; 7.5 (d, 1H) ; 7.8 (s, 1H) ; 8.4 (s, 1H) ; 8.75 (s, 1H)

Example 226

Using an analogous procedure to that described in Example 224, 7-hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline, (prepared as described for the starting material in Example 217), (98 mg, 0.32 mmol) was reacted with 4-(3-hydroxypropyl)morpholine (70 mg, 0.48 mmol), (prepared as described for the starting material in Example 60), to give 4-(indol-6-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride (26 mg, 19 %).


MS-ESI : 434 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.35 (m, 2H) ; 3.15 (m, 2H) ; 3.3 (t, 2H) ; 3.52 (d, 2H) ; 3.8 (t, 2H) ; 4.0 (d, 2H) ; 4.1 (s, 3H) ; 4.3 (t, 2H) ; 6.5 (s, 0.5 H, partly exchanged) ; 7.3 (d, 1H) ; 7.4 (s, 1H) ; 7.45 (s, 1H) ; 7.65 (d, 1H) ; 7.75 (s, 1H) ; 8.3 (s, 1H) ; 8.75 (s, 1H)

5 Examples 227-229

Using an analogous procedure to that described in Example 226, 7-hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline, (prepared as described for the starting material in Example 217), was used in the synthesis of the compounds described in Table XVI.

10 Table XVI



Example number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	Note	R
227	24	17	441	a	
228	14	10	430	b	
229	15	10	447	c	

a) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 2-((N-methyl-N-(4-pyridyl))amino)ethanol (73 mg), (EP 0359389A1), to give 4-(indol-6-ylamino)-6-methoxy-7-(2-((N-methyl-N-(4-pyridyl))amino)ethoxy)quinazoline hydrochloride.

¹H NMR Spectrum: (DMSO-d₆) 3.3 (s, 3H) ; 4.0 (s, 3H) ; 4.18 (t, 2H) ; 4.45 (t, 2H) ; 6.5 (s, 1H) ; 7.35 (d, 1H) ; 7.35-7.5 (m, 4H) ; 7.62 (d, 1H) ; 7.75 (s, 1H) ; 8.3 (d, 2H) ; 8.4 (s, 1H) ; 8.75 (s, 1H)

- 5 b) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with (E)-4-(pyrrolidin-1-yl)but-2-en-1-ol (68 mg, 0.48 mmol), (prepared as described for the starting material in Example 129) to give **4-(indol-6-ylamino)-6-methoxy-7-((E)-4-(pyrrolidin-1-yl)but-2-en-1-yloxy)quinazoline** hydrochloride.

¹H NMR Spectrum: (DMSO-d₆) 1.8-2.1 (m, 4H) ; 2.9-3.1 (m, 2H) ; 3.4-3.5 (br s, 2H) ; 3.87 (d, 10 2H) ; 4.05 (s, 3H) ; 4.9 (d, 2H) ; 6.1 (m, 1H) ; 6.3 (m, 1H) ; 6.5 (s, 1H) ; 7.25 (d, 1H) ; 7.45 (m, 2H) ; 7.65 (d, 1H) ; 7.75 (s, 1H) ; 8.3 (s, 1H) ; 8.8 (s, 1H)

- c) 7-Hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg) was reacted with 1-(3-hydroxypropyl)-4-methylpiperazine (76 mg), (prepared as described for the starting material 15 in Example 133), to give **4-(indol-6-ylamino)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline** hydrochloride.

Example 230

Using an analogous procedure to that described in Example 224, 7-hydroxy-6- 20 methoxy-4-(2-methylindol-5-ylamino)quinazoline (102 mg, 0.32 mmol), (prepared as described for the starting material in Example 205), was reacted with 1-(3-hydroxypropyl)-2-methylimidazole (67 mg, 0.48 mmol), (EP 0060696 A1), to give **6-methoxy-7-(3-(2-methylimidazol-1-yl)propoxy)-4-(2-methylindol-5-ylamino)quinazoline** (53 mg, 37 %).

MS-ESI : 443 [MH]⁺


¹H NMR Spectrum: (DMSO-d₆) 2.42 (s, 3H) ; 2.62 (s, 3H) ; 4.03 (s, 3H) ; 4.3 (t, 2H) ; 4.35 (t, 25 2H) ; 6.2 (s, 1H) ; 7.22 (d, 1H) ; 7.35 (d, 1H) ; 7.45 (s, 1H) ; 7.6 (dd, 1H) ; 7.65 (dd, 1H) ; 7.7 (s, 1H) ; 8.35 (s, 1H) ; 8.75 (s, 1H)

Examples 231-235

30 Using an analogous procedure to that described in Example 224, 7-hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (102 mg, 0.32 mmol), (prepared as

described for the starting material in Example 205), was used in the synthesis of the compounds described in Table XVII.

Table XVII



5

Example number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	Note	R
231	49	31	497	a	
232	25	18	444	b	
233	23	15	476	c	
234	33	22	461	d	
235	26	19	423	e	

a) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (102 mg) was reacted with 3-((N-(2,6-dimethyl-4-pyridyl)-N-methyl)amino)propan-1-ol (93mg), (prepared as described for the starting material in Example 222), to give 7-(3-((N-(2,6-dimethyl-4-pyridyl)-N-methyl)amino)propoxy)-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline.

10

¹H NMR Spectrum: (DMSO-d₆) 2.2 (m, 2H) ; 2.4 (s, 6H) ; 2.45 (s, 3H) ; 3.15 (s, 3H) ; 3.75 (t, 2H) ; 4.02 (s, 3H) ; 4.25 (t, 2H) ; 6.2 (s, 1H) ; 6.72 (br s, 1H) ; 6.85 (br s, 1H) ; 7.2 (dd, 1H) ; 7.3-7.4 (m, 2H) ; 7.62 (s, 1H) ; 8.3 (s, 1H) ; 8.7 (s, 1H)

- 5 b) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (102 mg) was reacted with (E)-4-(pyrrolidin-1-yl)but-2-en-1-ol (68 mg, 0.48 mmol), (prepared as described for the starting material in Example 129) to give **6-methoxy-4-(2-methylindol-5-ylamino)-7-((E)-4-(pyrrolidin-1-yl)but-2-en-1-yloxy)quinazoline**.

10 ¹H NMR Spectrum: (DMSO-d₆) 1.8-2.1 (m, 4H) ; 2.4 (s, 3H) ; 2.9-3.1 (m, 2H) ; 3.4-3.6 (m, 2H) ; 3.9 (d, 2H) ; 4.05 (s, 3H) ; 4.9 (d, 2H) ; 6.1 (m, 1H) ; 6.2 (s, 1H) ; 6.3 (d,t, 1H) ; 7.2 (m, 1H) ; 7.37 (d, 1H) ; 7.4 (s, 1H) ; 7.32 (s, 1H) ; 8.3 (s, 1H) ; 8.75 (s, 1H)

- 15 c) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (102 mg) was reacted with 3-((4-methyl-4*H*-1,2,4-triazol-3-yl)sulphanyl)propan-1-ol (83 mg), (prepared as described for the starting material in Example 212), to give **6-methoxy-4-(2-methylindol-5-ylamino)-7-(3-((4-methyl-4*H*-1,2,4-triazol-3-yl)sulphanyl)propoxy)quinazoline**.

20 ¹H NMR Spectrum: (DMSO-d₆) 2.25 (m, 2H) ; 2.45 (s, 3H) ; 3.35 (t, 2H) ; 3.65 (s, 3H) ; 4.05 (s, 3H) ; 4.35 (t, 2H) ; 6.2 (s, 1H) ; 7.2 (d, 1H) ; 7.35 (s, 1H) ; 7.37 (d, 1H) ; 7.62 (s, 1H) ; 8.25 (s, 1H) ; 8.75 (s, 1H) ; 8.9 (s, 1H)

- 25 d) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (102 mg) was reacted with 1-(3-hydroxypropyl)-4-methylpiperazine (76 mg), (prepared as described for the starting material in Example 133), to give **6-methoxy-4-(2-methylindol-5-ylamino)-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline**.

- 30 e) 7-Hydroxy-6-methoxy-4-(2-methylindol-5-ylamino)quinazoline (102 mg) was reacted with 2-(2-methoxyethoxy)ethanol to give **6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(2-methylindol-5-ylamino)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.45 (s, 3H) ; 3.28 (s, 3H) ; 3.5 (t, 2H) , 3.65 (t, 2H) ; 3.9 (t, 2H) ; 4.02 (s, 3H) ; 4.33 (t, 2H) ; 6.2 (s, 1H) ; 7.2 (d, 1H) ; 7.4 (m, 2H) ; 7.63 (s, 1H) ; 8.28 (s, 1H) ; 8.73 (s, 1H)

Example 236

A solution of 4-chloro-6-methoxy-7-((1-cyanomethylpiperidin-4-yl)methoxy)quinazoline (200 mg, 0.58 mmol) and 5-hydroxyindole (85 mg, 0.63 mmol) in DMF (3 ml) containing cesium carbonate (282 mg, 0.86 mmol) was stirred at 90°C for 90 minutes. After cooling, the mixture was poured onto water (25 ml). The precipitate was filtered, dried under vacuum and purified by reverse phase column chromatography on silica (kromasil ® C18) eluting with methanol/water (1 % acetic acid) (1/1). The fractions containing the expected product were combined and evaporated to give 7-((1-cyanomethyl)piperidin-4-ylmethoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (44 mg, 17 %).

MS-ESI : 444 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.7 (m, 2H) ; 2.15 (d, 2H) ; 2.2-2.35 (m, 1H) ; 3.20 (t, 2H) ; 3.65 (d, 2H) ; 4.1 (s, 3H) ; 4.25 (d, 2H) ; 4.62 (s, 2H) ; 6.5 (s, 0.5 H, partly exchanged) ; 7.1 (dd, 1H) ; 7.5 (s, 1H) ; 7.5-7.6 (m, 3H) ; 7.85 (s, 1H) ; 9.1 (s, 1H)

The starting material was prepared as follows:

To a suspension of 6-methoxy-7-(piperidin-4-ylmethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one hydrochloride (34 g, 84 mmol), (prepared as described for the starting material in Example 12), in water cooled at 0°C was added 1N sodium hydroxide until the mixture was at pH8. The solution was extracted with trichloromethane and the organic layer was dried (MgSO₄), filtered and evaporated to give 6-methoxy-7-(piperidin-4-ylmethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (29g).


To a solution of 6-methoxy-7-(piperidin-4-ylmethoxy)-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (28.9 g, 72 mmol) and aqueous formaldehyde 12 M (11.95 ml, 141 mmol) in methanol/THF (1/1) (580 ml) was added sodium cyanoborohydride (5.7 g, 86 mmol) in portions. After stirring for 90 minutes at ambient temperature, the volatiles were removed under vacuum and the residue was partitioned between methylene chloride and water. The organic layer was separated, dried (MgSO₄) and evaporated. The residue was dissolved in methanol saturated with ammonia (500 ml). The mixture was stirred for 36 hours at ambient temperature. The volatiles were removed under vacuum. The residue was triturated with a mixture ether/methylene chloride, filtered, washed with ether and dried under vacuum. The solid was dissolved in thionyl chloride (180 ml) and DMF (1.8 ml) was added.

After stirring at 80°C for 75 minutes the volatiles were removed under vacuum. The residue was azeotroped with toluene twice and the solid was partitioned between methylene chloride and water and the pH of the aqueous layer was adjusted to 9 with 2N sodium hydroxide. The organic layer was dried (MgSO₄) and evaporated. The residue was purified by column chromatography on aluminium oxide eluting with methylene chloride, followed by methylene chloride/ethyl acetate (70/30 followed by 50/50) followed by ethyl acetate and ethyl acetate/methanol (80/20) to give 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (11.2 g) (identical to the starting material prepared in Example 10) and 4-chloro-6-methoxy-7-((1-(cyanomethyl)piperidin-4-yl)methoxy)quinazoline (2.55 g).

MS-ESI : 347 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.42 (m, 2H) ; 1.85 (d, 2H) ; 1.8-1.9 (m, 1H) ; 2.2 (t, 2H) ; 2.85 (d, 2H) ; 3.75 (s, 2H) ; 4.05 (s, 3H) ; 4.15 (d, 2H) ; 7.42 (s, 1H) ; 7.5 (s, 1H) ; 8.9 (s, 1H)

Example 237



A solution of 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (2 gr, 6.22 mmol), (prepared as described for the starting material in Example 10), and 4-fluoro-5-hydroxy-2-methylindole (1.23 g, 7.46 mmol) in DMF (30 ml) containing potassium carbonate (1.28 g, 9.33 mmol) was stirred at 95°C for 2 hours. After cooling, the volatiles were removed under vacuum and the residue was triturated with ether, filtered and dried under vacuum. The residue was purified by column chromatography eluting with methanol/methylene chloride (1/9) followed by methanol/methanol saturated with ammonia/methylene chloride (20/1/79 followed by 20/5/75). The fractions containing the expected product were combined and evaporated. The solid was triturated with methanol, filtered and dried under vacuum to give 4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (1.95 g, 69 %).

MS-ESI : 451 [MH]⁺

¹H NMR Spectrum (DMSO-d₆) 1.4 (m, 2H) ; 1.8 (d, 2H) ; 1.7-1.9 (m, 1H) ; 1.9 (t, 2H) ; 2.2 (s, 3H) ; 2.45 (s, 3H) ; 2.8 (d, 2H) ; 4.02 (s, 3H) ; 4.1 (d, 2H) ; 6.25 (s, 1H) ; 7.0 (dd, 1H) ; 7.2 (d, 1H) ; 7.4 (s, 1H) ; 7.62 (s, 1H) ; 8.5 (s, 1H)

Elemental analysis:

Found C 64.2 H 6.5 N 11.7

5 C₂₅H₂₇FN₄O₃ 0.91methanol 0.08CH₂Cl₂ 0.1H₂O Requires C 63.9 H 6.4 N 11.5%

The starting material was prepared as follows:

To a solution of 2-fluoro-4-nitroanisole (9.9 g, 58 mmol) and 4-chlorophenoxyacetonitrile (10.7 g, 64 mmol) in DMF (50 ml) cooled at -15°C was added
10 potassium *tert*-butoxide (14.3 g, 127 mmol) in DMF (124 ml). After stirring for 30 minutes at -15°C, the mixture was poured onto cooled 1N hydrochloric acid. The mixture was extracted with ethyl acetate. The organic layer was washed with 1N sodium hydroxide, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with methylene chloride. The fractions containing the expected product were combined and
15 evaporated. The residue was dissolved in ethanol (180 ml) and acetic acid (24 ml) containing 10 % palladium on charcoal (600 mg) and the mixture was hydrogenated under 3 atmospheres pressure for 2 hours. The mixture was filtered, and the volatiles were removed under vacuum. The residue was partitioned between ethyl acetate and water. The organic layer was separated, and washed with saturated sodium hydrogen carbonate followed by brine, dried
20 (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with methylene chloride to give a mixture of 4-fluoro-5-methoxyindole and 6-fluoro-5-methoxyindole (5.64 g, 59 %) in a ratio 1/2.

¹H NMR Spectrum: (DMSO-d₆) 3.85 (s, 3H) ; 6.38 (s, 1H, 6-Fluoro) ; 6.45 (s, 1H ; 4-Fluoro) ; 6.9-7.4 (m, 3H)

25 A solution of 4-fluoro-5-methoxyindole and 6-fluoro-5-methoxyindole in a ratio 1/2 (496 mg, 3 mmol), di-*tert*butyl dicarbonate (720 mg, 3.3 mmol) in acetonitrile (12 ml) containing DMAP (18 mg, 0.15 mmol) was stirred at ambient temperature for 24 hours. The volatiles were removed under vacuum. The residue was dissolved in ethyl acetate, washed with 1N hydrochloric acid, followed by water, brine, dried (MgSO₄) and evaporated to give a
30 mixture of 4-fluoro-5-methoxy-1-*tert*-butoxycarbonylindole and 6-fluoro-5-methoxy-1-*tert*-butoxycarbonylindole in a ratio 1/2 (702 mg, 88 %).

¹H NMR Spectrum: (DMSO-d₆) 1.65 (s, 9H) ; 3.9 (s, 3H) ; 6.6 (d, 1H, 6-fluoro) ; 6.72 (d, 1H, 4-fluoro) ; 7.2 (t, 1H, 6-fluoro) ; 7.4 (d, 1H, 4-fluoro) ; 7.62 (d, 1H, 6-fluoro) ; 7.68 (d, 1H, 4-fluoro) ; 7.78 (s, 1H, 4-fluoro) ; 7.85 (s, 1H, 6-fluoro)

To a solution of 4-fluoro-5-methoxy-1-*tert*-butoxycarbonylindole and 6-fluoro-5-methoxy-1-*tert*-butoxycarbonylindole in a ratio 1/2 (8.1 g, 30.5 mmol) in THF (100 ml) cooled at -65°C was added *tert*-butyllithium (1.7 M) (23 ml, 35.7 mmol). After stirring for 4 hours at -70°C, methyl iodide (8.66 g, 61 mmol) was added and the mixture was left to warm-up to ambient temperature. Water was added and the mixture was extracted with ether. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated and was used directly in the next step.

The crude product was dissolved in methylene chloride (100 ml) and TFA (25 ml) was added. After stirring for 1 hour at ambient temperature, the volatiles were removed under vacuum. The residue was dissolved in ethyl acetate and the organic layer was washed with 1N sodium hydroxide, followed by water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography, eluting with ethyl acetate/petroleum ether (3/7) to give 6-fluoro-5-methoxy-2-methylindole (1.6 g) and 4-fluoro-5-methoxy-2-methylindole (0.8 g, 48 %).

6-fluoro-5-methoxy-2-methylindole:

MS-ESI : 180 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.35 (s, 3H) ; 3.8 (s, 3H) ; 6.05 (s, 1H) ; 7.1 (s, 1H) ; 7.12 (s, 1H) ; 10.8 (s, 1H)

4-fluoro-5-methoxy-2-methylindole:

MS-ESI : 180 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.35 (s, 3H) ; 3.8 (s, 3H) ; 6.15 (s, 1H) ; 6.9 (t, 1H) ; 7.05 (d, 1H) ; 11.0 (s, 1H)

To a solution of 4-fluoro-5-methoxy-2-methylindole (709 mg, 3.95 mmol) in methylene chloride (9 ml) cooled at -30°C was added a solution of boron tribromide (2.18 g, 8.7 mmol) in methylene chloride (1 ml). After stirring for 1 hour at ambient temperature, the mixture was poured onto water and was diluted with methylene chloride. The pH of the aqueous layer was adjusted to 6. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography, eluting

with ethyl acetate/petroleum ether (3/7) to give 4-fluoro-5-hydroxy-2-methylindole (461 mg, 70 %).

MS-ESI : 166 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.35 (s, 3H) ; 6.05 (s, 1H) ; 6.65 (dd, 1H) ; 6.9 (d, 1H) ; 8.75 (s, 1H) ; 10.9 (s, 1H)

¹³C NMR Spectrum: (DMSO-d₆) 13.5 ; 94,0 ; 106,0 ; 112 ; 118.5 (d) ; 132 (d) ; 136 (d) ; 136.5 ; 142.5 (d)

Alternatively the 4-fluoro-5-hydroxy-2-methylindole may be prepared as follows:

To a suspension of sodium hydride (5.42 g, 226 mmol) (prewashed with pentane) in THF (100 ml) cooled at 10°C was added ethyl acetoacetate (29.4 g, 226 mmol) while keeping the temperature below 15°C. After completion of addition, the mixture was further stirred for 15 minutes and cooled to 5°C. A solution of 1,2,3-trifluoro-4-nitrobenzene (20 g, 113 mmol) in THF (150 ml) was added while keeping the temperature below 5°C. The mixture was then left to warm up to ambient temperature and stirred for 24 hours. The volatiles were removed under vacuum and the residue was partitioned between ethyl acetate and 2N aqueous hydrochloric acid. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was dissolved in concentrated hydrochloric acid (650 ml) and acetic acid (600 ml) and the mixture was refluxed for 15 hours. After cooling, the volatiles were removed under vacuum and the residue was partitioned between aqueous sodium hydrogen carbonate (5 %) and ethyl acetate. The organic layer was washed with sodium hydrogen carbonate, water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with ethylacetate/petroleum ether (75/25) to give 3-acetylmethyl-1,2-difluoro-4-nitrobenzene (17.5 g, 72 %).

¹H NMR Spectrum: (CDCl₃) 2.4 (s, 3H) ; 4.25 (s, 2H) ; 7.25 (dd, 1H) ; 8.0 (dd, 1H)

A solution of 3-acetylmethyl-1,2-difluoro-4-nitrobenzene (500 mg, 2.3 mmol) in methylene chloride (5 ml) containing montmorillonite K10 (1 g) and trimethyl orthoformate (5 ml) was stirred for 24 hours at ambient temperature. The solid was filtered, washed with methylene chloride and the filtrate was evaporated to give 1,2-difluoro-3-(2,2-dimethoxypropyl)-4-nitrobenzene (534 mg, 88 %).

¹H NMR Spectrum: (CDCl₃) 1.2 (s, 3H) ; 3.2 (s, 6H) ; 3.52 (s, 2H) ; 7.18 (dd, 1H) ; 7.6 (m, 1H)

To a solution of benzyl alcohol (221 mg, 2.05 mmol) in DMA (1.5 ml) was added 60% sodium hydride (82 mg, 2.05 mmol). The mixture was stirred for 1 hour at ambient temperature. A solution of 1,2-difluoro-3-(2,2-dimethoxypropyl)-4-nitrobenzene (534 mg, 2.05 mmol) in DMA (1.5 ml) was added and the mixture was stirred for 3 hours at ambient temperature. The mixture was diluted with 1N hydrochloric acid (10 ml) and extracted with ethyl acetate. The organic layer was evaporated and the residue was dissolved in THF (2 ml) and 6N hydrochloric acid (0.3 ml) was added. The mixture was stirred for 1 hour at ambient temperature and the solvents were removed under vacuum. The residue was partitioned between ethyl acetate and water. The organic layer was separated, washed with brine, dried (MgSO₄) and evaporated. The solid was triturated with ether, filtered, washed with ether and dried under vacuum to give 3-acetylmethyl-1-benzyloxy-2-fluoro-4-nitrobenzene (350 mg, 56 %).

¹H NMR Spectrum: (CDCl₃) 2.35 (s, 3H) ; 4.25 (s, 2H) ; 5.25 (s, 2H) ; 7.0 (dd, 1H) ; 7.32-7.5 (m, 5H) ; 8.0 (dd, 1H)

A solution of 3-acetylmethyl-1-benzyloxy-2-fluoro-4-nitrobenzene (300 mg, 0.99 mmol) in ethanol (10 ml) and acetic acid (1 ml) containing 10 % palladium on charcoal (30 mg) was hydrogenated at 2 atmospheres pressure for 2 hours. The mixture was filtered and the filtrate was evaporated. The residue was dissolved in ethyl acetate and the organic layer was washed with aqueous sodium hydrogen carbonate, brine and evaporated to give 4-fluoro-5-hydroxy-2-methylindole. The residue was purified by column chromatography eluting with ethyl acetate/petroleum ether (3/7) to give 4-fluoro-5-hydroxy-2-methylindole (63 mg, 30%). Analytical data as above.

Alternatively the 4-fluoro-5-methoxy-2-methylindole can be prepared as follows:

A solution of sodium methoxide (freshly prepared from sodium (1.71g) and methanol (35ml)) was added to a solution of 1,2-difluoro-3-(2,2-dimethoxypropyl)-4-nitrobenzene (16.2 g, 62 mmol), (prepared as described above), in methanol (200ml) cooled at 5°C. The mixture was left to warm to ambient temperature and was stirred for 3 days. The volatiles were removed under vacuum and the residue was partitioned between ethyl acetate and 2N hydrochloric acid (1ml). The organic layer was concentrated to a total volume of 100ml and THF (100ml) and 6N hydrochloric acid (25ml) were added. The mixture was stirred for 1 hour at ambient temperature. The volatiles were removed under vacuum and the residue was

partitioned between ethyl acetate and water. The organic layer was separated, washed with water, brine, dried (MgSO_4) and evaporated. The residue was purified by column chromatography eluting with ethyl acetate/petroleum ether (3/7) to give 3-acetylmethyl-2-fluoro-1-methoxy-4-nitrobenzene (12.7 g, 90%).

5 MS-ESI : 250 $[\text{MNa}]^+$

^1H NMR Spectrum: (CDCl_3) 2.38 (s, 3H) ; 4.0 (s, 3H) ; 4.25 (s, 2H) ; 7.0 (dd, 1H) ; 8.05 (d, 1H)


To a solution of 3-acetylmethyl-2-fluoro-1-methoxy-4-nitrobenzene (11.36g, 50 mmol) in acetone (200ml) was added 4M aqueous ammonium acetate (700ml) followed by a
10 solution of titanium trichloride (15% in water, 340ml) dropwise. The mixture was stirred for 10 minutes at ambient temperature and the mixture was extracted with ether. The organic layer was washed with 0.5N aqueous sodium hydroxide followed by water, brine, dried (MgSO_4) and the volatiles were removed under vacuum. The residue was purified by column chromatography eluting with methylene chloride to give 4-fluoro-5-methoxy-2-methylindole
15 (8.15g, 90%).

^1H NMR Spectrum: (DMSO) 2.35 (s, 3H) ; 3.8 (s, 3H) ; 6.1 (s, 1H) ; 6.85 (dd, 1H) ; 7.02 (d, 1H)

Cleavage of 4-fluoro-5-methoxy-2-methylindole with boron tribromide to give 4-fluoro-5-hydroxy-2-methylindole is described above.

20

Example 238




Using an analogous procedure to that described in Example 237, 4-chloro-6-methoxy-
25 7-(3-piperidinopropoxy)quinazoline (1.65 g, 4.89 mmol), (prepared as described for the starting material in Example 67), was reacted with 4-fluoro-5-hydroxy-2-methylindole (970 mg, 5.88 mmol), (prepared as described for the starting material in Example 237), to give 4-

(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline (1.9 g, 83 %).

MS-ESI : 465 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.4 (br s, 2H) ; 1.5 (m, 4H) ; 1.95 (m, 2H) ; 2.25-2.5 (m, 6H) ;
2.45 (s, 3H) ; 4.0 (s, 3H) ; 4.25 (t, 2H) ; 6.25 (s, 1H) ; 7.0 (dd, 1H) ; 7.15 (d, 1H) ; 7.4 (s, 1H) ;
7.6 (s, 1H) ; 8.5 (s, 1H)

Example 239




Using an analogous procedure to that described in Example 237, 4-chloro-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (106 mg, 0.30 mmol), (prepared as described for the starting material in Example 176), was reacted with 4-fluoro-5-hydroxy-2-methylindole (60 mg, 0.36 mmol), (prepared as described for the starting material in Example 237), to give 4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (100 mg, 70 %).

MS-ESI : 480 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.0 (t, 2H) ; 2.15 (s, 3H) ; 2.45 (s, 3H) ; 2.2-2.6 (m, 10H) ;
4.02 (s, 3H) ; 4.25 (t, 2H) ; 6.25 (s, 1H) ; 7.0 (dd, 1H) ; 7.18 (d, 1H) ; 7.4 (s, 1H) ; 7.62 (s, 1H)
; 8.5 (s, 1H)

Example 240



Using a procedure identical to that described in Example 237, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (2 g, 6.22 mmol), (prepared as described for the

starting material in Example 9), was reacted with 4-fluoro-5-hydroxy-2-methylindole (1.23 g, 7.46 mmol), (prepared as described for the starting material in Example 237), to give **4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (1.41 g, 50 %).

5 MS-ESI : 451 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.7 (br s, 4H) ; 2.0 (m, 2H) ; 2.41 (s, 3H) ; 2.5 (br s, 4H) ; 2.6 (t, 2H) ; 4.0 (s, 3H) ; 4.25 (t, 2H) ; 6.25 (s, 1H) ; 7.0 (dd, 1H) ; 7.2 (d, 1H) ; 7.4 (s, 1H) ; 7.6 (s, 1H) ; 8.5 (s, 1H)

Elemental analysis :

Found C 63.3 H 6.4 N 11.9

10 C₂₅H₂₇N₄O₃ · 1.08 H₂O ; 0.16 methanol

Requires C 63.6 H 6.3 N 11.8%

Example 241

A solution of 4-chloro-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline (300 mg, 0.9 mmol) and 4-fluoro-5-hydroxyindole (162 mg, 1 mmol), (prepared as described for the starting material in Example 242), in DMF (4.5 ml) containing potassium carbonate (185 mg, 1.3 mmol) was stirred at 90°C for 1 hour. After cooling, the mixture was filtered and the solid was washed with DMF. The filtrate was evaporated and the residue was purified by column chromatography, eluting with methylene chloride followed by methanol/methylene chloride (1/99) followed by methanol saturated with ammonia/methylene chloride (2/98). The fractions containing the expected product were combined and evaporated. The solid was triturated with ether, filtered, washed with ether and dried under vacuum to give **4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline** (282 mg, 69%).

MS-ESI : 451 [MH]⁺

25 ¹H NMR Spectrum: (DMSO-d₆) 1.2-1.3 (m, 2H) ; 1.4-1.55 (m, 1H) ; 1.7-1.9 (m, 6H) ; 2.15 (s, 3H) ; 2.75 (d, 2H) ; 4.0 (s, 3H) ; 4.3 (t, 2H) ; 6.55 (s, 1H) ; 7.1 (dd, 1H) ; 7.3 (d, 1H) ; 7.4 (s, 1H) ; 7.5 (s, 1H) ; 7.6 (s, 1H) ; 8.5 (s, 1H) ; 11.5 (s, 1H)

The starting material was prepared as follows:

30 To a solution of 4-(2-hydroxyethyl)-(1-*tert*-butoxycarbonyl)piperidine (12.9 g, 56 mmol), (prepared as described for the starting material in Example 126), in *tert*-butyl methyl ether (120 ml) containing 1,4-diazabicyclo[2.2.2]octane (9.8 g, 87 mmol) cooled at -5°C was

added a solution of tosyl chloride (14.5 gr, 76 mmol) in *tert*-butyl methyl ether (120ml) dropwise whilst keeping the temperature below 0°C. After completion of addition, the mixture was left to warm up to ambient temperature and stirred for 1 hour. The mixture was poured onto petroleum ether (240 ml). The precipitate was filtered and washed with petroleum ether. The filtrate was evaporated and the residue was dissolved in ether. The ether layer was washed with 0.5 N hydrochloric acid, followed by saturated sodium hydrogen carbonate, dried (MgSO₄) and evaporated to give 4-(2-(4-methylphenylsulphonyloxy)ethyl)-1-*tert*-butoxycarbonylpiperidine (20.9 g, 97%).

¹H NMR Spectrum: (CDCl₃) 0.95-1.05 (m, 4H) ; 1.45 (s, 9H) ; 1.4-1.6 (m, 3H) 2.45 (s, 3H) ; 2.62 (t, 2H) ; 3.9-4.1 (m, 2H) ; 4.1 (t, 2H) ; 7.35 (d, 2H) ; 7.8 (d, 2H)

A suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (7 g, 23 mmol), (prepared as described for the starting material in Example 12), 4-(2-(4-methylphenylsulphonyloxy)ethyl)-1-*tert*-butoxycarbonylpiperidine (11.4 g, 30 mmol) in DMF (70 ml) containing potassium carbonate (6.32 g, 46 mmol) was stirred at 100°C for 3 hours. After cooling, the volatiles were removed under vacuum and the residue was partitioned between ether and water. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The solid was triturated with pentane, filtered and dried under vacuum to give 7-(2-(1-*tert*butoxycarbonylpiperidin-4-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (10.5 g, 88 %).

MS-ESI : 540 [MNa]⁺

¹H NMR Spectrum: (CDCl₃) 1.2 (s, 9H) ; 1.15-1.25 (m, 2H) ; 1.48 (s, 9H) ; 1.65-1.75 (m, 1H) ; 1.7 (d, 2H) ; 1.9 (dd, 2H) ; 2.72 (t, 2H) ; 4.0 (s, 3H) ; 4.0-4.2 (m, 2H) ; 4.2 (t, 2H) ; 5.95 (s, 2H) ; 7.1 (s, 1H) ; 7.65 (s, 1H) ; 8.2 (s, 1H)

A solution of 7-(2-(1-*tert*butoxycarbonylpiperidin-4-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (10.5 g, 20 mmol) in methylene chloride (100 ml) containing TFA (25 ml) was stirred for 1 hour at ambient temperature. Water (50 ml) and methylene chloride (100 ml) were added and the pH of the aqueous layer was adjusted to 8 with solid sodium hydrogen carbonate. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was triturated with ether and the solid was filtered and dried under vacuum to give 7-(2-(piperidin-4-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (8.3 g, 100 %).

¹H NMR Spectrum: (CDCl₃) 1.2 (s, 9H) ; 1.65 (m, 2H) ; 1.9 (br s, 2H) ; 1.8-1.9 (m, 1H) ; 2.0 (d, 2H) ; 2.9 (t, 2H) ; 3.45 (d, 2H) ; 4.0 (s, 3H) ; 4.2 (t, 2H) ; 5.95 (s, 2H) ; 7.1 (s, 1H) ; 7.65 (s, 1H) ; 8.2 (s, 1H)

To a solution of 7-(2-(piperidin-4-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (6 g, 14.4 mmol) in methanol (30 ml) and methylene chloride (60 ml) was added 37% aqueous formaldehyde (2.2 ml ; 28.9 mmol) followed by acetic acid (990 µl ; 17.3 mmol). Sodium borohydride triacetate (4.6 g, 21.6 mmol) was added in portions. After stirring for 1 hour at ambient temperature, the volatiles were removed under vacuum and the residue was partitioned between water (50 ml) and methylene chloride (50 ml). The pH of the aqueous layer was adjusted to 7, washed with water, brine, dried (MgSO₄) and evaporated. The solid was triturated with ether, filtered, washed with ether and dried under vacuum to give 7-(2-(1-methylpiperidin-4-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (4.2 g, 68 %).

MS-ESI : 432 [MH]⁺

¹H NMR Spectrum: (CDCl₃) 1.22 (s, 9H) ; 1.68 (br s, 3H) ; 1.9 (m, 4H) ; 2.32 (br s, 2H) ; 2.52 (s, 3H) ; 3.18 (d, 2H) ; 4.0 (s, 3H) ; 4.2 (t, 2H) ; 5.95 (s, 2H) ; 7.1 (s, 1H) ; 7.65 (s, 1H) ; 8.2 (s, 2H)

A solution of 7-(2-(1-methylpiperidin-4-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (4.2 g, 9.7 mmol) in methanol saturated with ammonia (150 ml) was stirred overnight at ambient temperature. The volatiles were removed under vacuum and the residue was triturated with ether. The solid was filtered, washed with ether and dried under vacuum to give 7-(2-(1-methylpiperidin-4-yl)ethoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (3.12 g, 100 %).

MS-ESI : 318 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.3 (m, 2H) ; 1.58 (br s, 1H) ; 1.72 (dd, 2H) ; 1.8 (d, 2H) ; 2.4 (s, 3H) ; 2.2-2.45 (m, 2H) ; 3.0 (br s, 2H) ; 3.85 (s, 3H) ; 4.15 (t, 2H) ; 7.15 (s, 1H) ; 7.45 (s, 1H) ; 8.0 (s, 1H)

A solution of 7-(2-(1-methylpiperidin-4-yl)ethoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (3.1 g, 9.8 mmol) in thionyl chloride (40 ml) containing DMF (400 µl) was refluxed for 4 hours. After cooling, the volatiles were removed under vacuum. The residue was partitioned between methylene chloride and water and the pH of the aqueous layer was adjusted to 11 with solid sodium hydrogen carbonate and aqueous ammonia. The organic

layer was separated, dried (MgSO_4) and evaporated. The residue was triturated with ether, filtered, washed with ether and dried under vacuum to give 4-chloro-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline (1.83 g, 54 %).

MS-ESI : 336 $[\text{MH}]^+$

- 5 ^1H NMR Spectrum: (CDCl_3) 1.4-1.7 (m, 3H) ; 1.8 (d, 2H) ; 1.9 (dd, 2H) ; 2.05 (t, 2H) ; 2.35 (s, 3H) ; 2.95 (d, 2H) ; 4.05 (s, 3H) ; 4.25 (t, 2H) ; 7.3 (s, 1H) ; 7.4 (s, 1H) ; 8.88 (s, 1H)

Example 242

- 10 A solution of 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (213 mg, 0.662 mmol), (prepared as described for the starting material in Example 10), and 6-fluoro-5-hydroxyindole (120 mg, 0.794 mmol) in DMF (3 ml) containing potassium carbonate (137 mg, 0.994 mmol) was stirred at 95°C for 3.5 hours. After cooling, the mixture was poured onto water. The mixture was filtered and the solid was washed with water. The solid was dissolved in methylene chloride. The organic layer was dried (MgSO_4), and evaporated.
- 15 The residue was triturated with ether/ethyl acetate and the solid was filtered and dried under vacuum to give 4-(6-fluoroindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (135 mg, 46 %).

MS-ESI : 437 $[\text{MH}]^+$

- 20 ^1H NMR Spectrum: (DMSO-d_6) 1.3-1.45 (m, 2H) ; 1.8 (d, 2H) ; 1.9 (t, 2H) ; 1.7-1.9 (m, 1H) ; 2.17 (s, 3H) ; 2.8 (d, 2H) ; 4.0 (s, 3H) ; 4.1 (d, 2H) ; 6.48 (br s, 1H) ; 7.38 (d, 1H) ; 7.4 (s, 1H) ; 7.42 (t, 1H) ; 7.58 (d, 1H) ; 7.6 (s, 1H) ; 8.5 (s, 1H)

Elemental analysis Found C 65.0 H 5.8 N 12.7

$\text{C}_{24}\text{H}_{25}\text{FN}_4\text{O}_3 \cdot 0.4 \text{H}_2\text{O}$ Requires C 65.0 H 5.9 N 12.6%

- 25 The starting material was prepared as follows:

- A mixture of 2-fluoro-4-nitrophenol (15g, 95.5 mmol) and benzyl bromide (18g, 105 mmol) in acetone (125 ml) containing potassium carbonate (26.5 gr, 190 mmol) was refluxed for 2 hours. The volatiles were removed and the residue was partitioned between 2N hydrochloric acid and ethyl acetate. The organic layer was separated, washed with water, brine, dried (MgSO_4) and the volatiles were removed under vacuum. The solid was triturated with petroleum ether to give 2-fluoro-4-nitro-benzyloxybenzene (23g, 97%).
- 30

^1H NMR Spectrum: (CDCl_3) 5.3 (s, 2H) ; 7.1 (t, 1H) ; 7.35-7.55 (m, 5H) ; 8.0 (m, 2H)

To a solution of potassium *tert*-butoxide (1.72g, 15.4 mmol) in DMF (15 ml) cooled at -30°C, was added dropwise a solution of 2-fluoro-4-nitro-benzyloxybenzene (1.73g, 7 mmol) and 4-chlorophenoxyacetonitrile (1.29 g, 7.7 mmol) while maintaining the temperature below -25°C. After completion of addition, the mixture was stirred for 30 minutes at -20°C and then poured onto a mixture of cold 1N hydrochloric acid and ether. The organic layer was separated, washed with 1N sodium hydroxide, followed by water, brine, dried (MgSO₄). The volatiles were removed under vacuum and the residue was purified by column chromatography eluting with methylene chloride/petroleum ether (3/1) to give a mixture of 3-cyanomethyl-2-fluoro-4-nitrobenzyloxybenzene and 5-cyanomethyl-2-fluoro-4-nitrobenzyloxybenzene (1.2 g, 60%).

¹H NMR Spectrum: (DMSO-d₆) 4.22 (s, 2H, 3-cyanomethyl isomer) ; 4.3 (s, 2H, 5-cyanomethyl isomer); 5.32 (s, 2H, 5-cyanomethyl isomer) ; 5.36 (s, 2H, 3-cyanomethyl isomer); 7.3-7.7 (m, 6H); 8.1 (d, 1H, 3-cyanomethyl isomer); 8.2 (d, 1H, 5-cyanomethyl isomer)

A solution of a mixture of 3-cyanomethyl-2-fluoro-4-nitrobenzyloxybenzene and 5-cyanomethyl-2-fluoro-4-nitrobenzyloxybenzene (23g, 80.4 mmol) in ethanol (220ml) and acetic acid (30ml) containing 10% palladium on charcoal (600mg) was hydrogenated under 3 atmospheres pressure until hydrogen uptake ceased. The mixture was filtered and the filtrate was evaporated under vacuum. The residue was purified on column chromatography using a Prochrom® equipment eluting with methylene chloride/petroleum ether (20/80) to give 4-fluoro-5-hydroxyindole (2.48g) and 6-fluoro-5-hydroxyindole (3.5 g).


4-fluoro-5-hydroxyindole:

¹H NMR Spectrum: (DMSO-d₆) 6.32 (s, 1H) ; 6.75 (dd, 1H) ; 7.0 (d, 1H) ; 7.28 (dd, 1H) ; 8.8 (br s, 1H) ; 11.05 (br s, 1H)

6-fluoro-5-hydroxyindole:

¹H NMR Spectrum: (DMSO-d₆) 6.25 (s, 1H) ; 7.0 (d, 1H) ; 7.12 (d, 1H) ; 7.2 (dd, 1H) ; 9.0 (br s, 1H)

Example 243



A solution of 4-chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (213 mg, 0.662 mmol), (prepared as described for the starting material in Example 10), and 4-fluoro-5-hydroxyindole (120 mg, 0.794 mmol), (prepared as described for the starting material in Example 242), in DMF (3 ml) containing potassium carbonate (137 mg, 0.994 mmol) was stirred at 95°C for 3 hours. After cooling, the mixture was partitioned between ethyl acetate and water. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was triturated in cold ether. The solid was filtered and dried under vacuum to give 4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (77 mg, 26 %).

MS - ESI : 437 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5 (m, 2H) ; 1.8 (d, 2H) ; 1.9 (t, 2H) ; 1.7-1.95 (m, 1H) ; 2.2 (s, 3H) ; 2.8 (d, 2H) ; 4.02 (s, 3H) ; 4.1 (d, 2H) ; 6.55 (s, 1H) ; 7.1 (t, 1H) ; 7.3 (d, 1H) ; 7.4 (s, 1H) ; 7.48 (t, 1H) ; 7.62 (s, 1H) ; 8.5 (s, 1H)

Elemental analysis	Found	C 64.8	H 5.8	N 12.6
C ₂₄ H ₂₅ FN ₄ O ₃ 0.4 H ₂ O	Requires	C 65.0	H 5.9	N 12.6%

Example 244

A mixture of 4-chloro-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (282 mg, 0.662 mmol), 6-fluoro-5-hydroxyindole (120 mg, 0.794 mmol), (prepared as described for the starting material in Example 242), in DMF (3 ml) containing potassium carbonate (137 mg, 0.994 mmol) was heated at 95°C for 3 hours. After cooling, the residue was poured in water (12 ml) and the pH was adjusted to 8. The mixture was extracted with ethyl acetate. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by preparative column chromatography on C¹⁸ silica eluting with 60 % methanol in aqueous ammonium carbonate (2g ammonium carbonate/litre saturated with CO₂). The fractions containing the expected product were combined and evaporated. The residue was triturated with ether and the solid was filtered, dried under

vacuum to give 4-(6-fluoroindol-5-yloxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (147 mg, 48 %).

MS-ESI : 466 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.3-2.4 (m, 2H) ; 3.0 (s, 3H) ; 3.2-3.9 (m, 8H) ;
5 3.5 (t, 2H) ; 4.1 (s, 3H) ; 4.4 (t, 2H) ; 6.52 (d, 1H) ; 7.45 (d, 1H) ; 7.48 (s, 1H) ; 7.6 (s, 1H) ;
7.65 (d, 1H) ; 7.82 (s, 1H) ; 9.0 (s, 1H)

Elemental analysis	Found	C 62.1	H 6.4	N 14.2
C ₂₅ H ₂₈ FN ₅ O ₃ · 0.9 H ₂ O	Requires	C 62.3	H 6.2	N 14.5%

10 The starting material was prepared as follows:

To a suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (29g, 94.7 mmol), (prepared as described for the starting material in Example 12), in methylene chloride (280ml) cooled at 5°C was added triphenylphosphine (37.1g, 141.6 mmol) followed by 3-bromo-1-propanol (12.8 ml, 141.6 mmol) and diethyl
15 azodicarboxylate (2.4 ml, 141.6 mmol) dropwise. After stirring for 2 hours at ambient temperature, the volatiles were removed under vacuum and the residue was purified by column chromatography eluting with methylene chloride/methanol (98/2). The fractions containing the expected product were combined and evaporated and the solid was triturated with ether, filtered, washed with ether and dried under vacuum to give 7-(3-bromopropoxy)-6-
20 methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (37.22 g, 92%).

MS-ESI : 427-429 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.18 (s, 9H) ; 2.32 (m, 2H) ; 3.7 (t, 2H) ; 3.92 (s, 3H) ; 4.28 (t, 2H) ; 5.95 (s, 2H) ; 7.2 (s, 1H) ; 7.5 (s, 1H) ; 8.4 (s, 1H)

A suspension of 7-(3-bromopropoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (36.7 g, 86 mmol) in 1-methylpiperazine (370ml) was stirred at
25 100°C for 90 minutes. After removal of the volatiles under vacuum, the residue was partitioned between methylene chloride and aqueous ammonium chloride. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The solid was triturated with ether, filtered, washed with ether and dried under vacuum to give 7-(3-(4-
30 methylpiperazin-1-yl)propoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (31.9 gr, 83%).

MS-ESI : 447 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.15 (s, 9H) ; 2.25 (t, 2H) ; 2.5 (s, 3H) ; 3.45 (t, 2H) ; 3.2-4.0 (m, 8H); 3.9 (s, 3H) ; 4.25 (t, 2H) ; 5.95 (s, 2H) ; 7.22 (s, 1H) ; 7.55 (s, 1H) ; 8.6 (s, 1H)

A suspension of 7-(3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-3-

- 5 ((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (31.8 g, 71.3 mmol) in methanol saturated with ammonia was stirred at ambient temperature overnight. The volatiles were removed under vacuum. The solid was triturated with ether containing about 10% of methylene chloride, filtered, washed with ether containing about 10% methylene chloride and dried under vacuum to give 7-(3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-3,4-
- 10 dihydroquinazolin-4-one (22.63g, 95%).

MS-ESI : 333 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.92 (m, 2H) ; 2.15 (s, 3H) ; 2.2-2.5 (m, 10H) ; 3.88 (s, 3H) ; 4.15 (t, 2H) ; 7.1 (s, 1H) ; 7.45 (s, 1H) ; 7.98 (s, 1H)


A solution of 7-(3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-3,4-

- 15 dihydroquinazolin-4-one (22.6 g, 68 mmol) in thionyl chloride (300ml) containing DMF (5 ml) was refluxed for 2 hours. After cooling, the volatiles were removed under vacuum and the residue was azeotroped with toluene twice. The solid was dissolved in methylene chloride and water was added. The mixture was cooled to 0°C and the pH of the aqueous layer was adjusted to 7 with solid hydrogen carbonate and then raised to 10 with 6N Sodium hydroxide.
- 20 The organic layer was separated and the aqueous layer was extracted with methylene chloride. The organic layer was washed with brine, dried (MgSO₄), filtered and the volatiles were removed under vacuum. The residue was triturated with ether, filtered, washed with ether and dried under vacuum to give 4-chloro-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (16.3 gr, 68%).

- 25 MS-ESI : 351-353 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.98 (t, 2H) ; 2.18 (s, 3H) ; 2.45 (t, 2H) ; 2.22-2.5 (m, 8H) ; 4.05 (s, 3H) ; 4.28 (t, 2H) ; 7.4 (s, 3H) ; 7.45 (s, 1H) ; 8.9 (s, 1H)

Example 245



Using an analogous procedure to that described in Example 243, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (213 mg, 0.662 mmol), (prepared as described for the starting material in Example 9), was reacted with 6-fluoro-5-hydroxyindole (120 mg, 0.794 mmol), (prepared as described for the starting material in Example 242), in DMF (3 ml) containing potassium carbonate (137 mg, 0.993 mmol) to give **4-(6-fluoroindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (154 mg, 53 %).

MS-ESI : 437 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.7-1.8 (m, 4H) ; 2.0-2.1 (m, 2H) ; 2.48 (br s, 4H) ; 2.6 (t, 2H) ; 4.02 (s, 3H) ; 4.3 (t, 2H) ; 6.5 (s, 1H) ; 7.4 (d, 1H) ; 7.4 (s, 1H) ; 7.45 (t, 1H) ; 7.6 (d, 1H) ; 7.62 (s, 1H) ; 8.52 (s, 1H)

Elemental analysis Found C 65.4 H 6.0 N 12.9

C₂₄H₂₅N₄O₃ · 0.2 H₂O Requires C 65.5 H 5.8 N 12.7%

15 **Example 246**

To a solution of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline (500 mg, 1.2 mmol), (prepared as described in Example 70), in methanol (11.5 ml) containing potassium iodide (99 mg, 0.6 mmol) was added 4-(2-chloroethyl)morpholine hydrochloride (134 mg, 0.72 mmol) followed by sodium hydrogen carbonate (151 mg, 1.8 mmol). After stirring for 1 hour at reflux, 4-(2-chloroethyl)morpholine hydrochloride (134 mg, 0.72 mmol) and sodium hydrogen carbonate (151 mg, 1.8 mmol) were added. After stirring 1 hour at reflux, the mixture was cooled and the precipitate was filtered, washed with methanol followed by water and dried over phosphorus pentoxide to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-(2-morpholinoethyl)piperidin-4-ylmethoxy)quinazoline** (470 mg, 73 %).


MS-ESI : 532 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.45 (m, 2H) ; 1.8 (d, 2H) ; 1.7-1.9 (m, 1H) ; 2.0 (t, 2H) ; 2.3-2.45 (m, 8H) ; 2.4 (s, 3H) ; 2.95 (d, 2H) ; 3.6 (t, 4H) ; 4.0 (s, 3H) ; 4.08 (d, 2H) ; 6.18 (s,

1H) ; 6.9 (dd, 1H) ; 7.3 (s, 1H) ; 7.35 (d, 1H) ; 7.4 (s, 1H) ; 7.6 (s, 1H) ; 8.5 (s, 1H) ; 11.05 (s, 1H)

Elemental analysis	Found	C 65.3	H 7.1	N 12.6
C ₃₀ H ₃₇ N ₅ O ₄ 0.6 H ₂ O 0.6 Methanol	Requires	C 65.4	H 7.3	N 12.5%

5

Example 247

A solution of 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (1.76 g, 5.47 mmol), (prepared as described for the starting material in Example 9), 4-fluoro-5-hydroxyindole (0.992 g, 6.57 mmol), (prepared as described for the starting material in Example 242), in DMF (25 ml) containing potassium carbonate (1.14 g ; 8.21 mmol) was heated at 95°C for 1 hour. After cooling, the mixture was filtered and washed with DMF. The filtrate was evaporated and the residue was purified by column chromatography eluting with methanol/methylene chloride (1/9) followed by methanol/methanol chloride/methanol (containing ammonia) (16/80/4). The fractions containing the expected product were combined and evaporated. The residue was repurified by column chromatography eluting with a gradient of methylene chloride/methanol (80/20 to 40/60). The fractions containing the expected product were combined and evaporated. The residue was triturated in cold methanol and the solid was filtered, washed with ether and dried under vacuum to give 4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (1.24 g, 52 %).

MS-ESI : 437 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.7 (br s, 4H) ; 2.0 (m, 2H) ; 2.45 (br s, 4H) ; 2.6 (t, 2H) ; 4.05 (s, 3H) ; 4.28 (t, 2H) ; 6.58 (s, 1H) ; 7.1 (t, 2H) ; 7.35 (d, 1H) ; 7.4 (s, 1H) ; 7.5 (t, 1H) ; 7.65 (s, 1H) ; 8.52 (s, 1H)

Elemental analysis	Found	C 65.3	H 5.9	N 12.6
C ₂₄ H ₂₅ FN ₄ O ₃ 0.19 Methanol, 0.17 H ₂ O	Requires	C 65.2	H 5.9	N 12.6%

Example 248

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (222 mg, 0.662 mmol), (prepared as described for the starting material in Example 67), and 6-fluoro-5-hydroxyindole (120 mg, 0.794 mmol), (prepared as described for the starting material in Example 242), in DMF (3 ml) containing potassium carbonate (137 mg, 0.993 mmol) was heated at 95°C for 3.5 hours. After cooling the mixture was poured onto water and extracted with ethyl acetate. The organic layers were washed with water, brine, dried (MgSO₄) and evaporated. The residue was triturated with ether, filtered and dried under vacuum to give **4-(6-fluoroindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline** (139 mg, 46 %).

MS-ESI : 451 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.35-1.45 (m, 2H) ; 1.45-1.6 (m, 4H) ; 2.0 (m, 2H) ; 2.35 (br s, 4H) ; 2.42 (t, 2H) ; 4.05 (s, 3H) ; 4.25 (t, 2H) ; 6.5 (s, 1H) ; 7.4 (d, 1H) ; 7.42 (s, 1H) ; 7.44 (t, 1H) ; 7.6 (d, 1H) ; 7.65 (s, 1H) ; 8.5 (s, 1H)


Elemental analysis

Found C 65.9 H 6.2 N 12.3

C₂₅H₂₇FN₄O₃ · 0.3 H₂O

Requires C 65.9 H 6.1 N 12.3%

Example 249



Using an analogous procedure to that described in Example 244, 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (407 mg, 1.21 mmol), (prepared as described for the starting material in Example 67), 4-fluoro-5-hydroxyindole (220 mg, 1.45 mmol) (prepared as described for the starting material in Example 242), and potassium carbonate (251 mg, 1.82 mmol) in DMF (6 ml) were heated at 95°C for 90 minutes and purified to give **4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline** (367 mg, 67 %).

MS-ESI : 451 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.35-1.45 (m, 2H) ; 1.55 (m, 4H) ; 2.0 (m, 2H) ; 2.38 (br s, 4H) ; 2.45 (t, 2H) ; 4.02 (s, 3H) ; 4.25 (t, 2H) ; 6.55 (s, 1H) ; 7.12 (dd, 1H) ; 7.32 (d, 1H) ; 7.4 (s, 1H) ; 7.5 (s, 1H) ; 7.65 (s, 1H) ; 8.52 (s, 1H)

Elemental analysis

Found

C 66.0 H 6.2 N 12.4

$C_{25}H_{27}FN_4O_3 \cdot 0.2 H_2O$

Requires

C 66.1 H 6.1 N 12.3%

Example 250

Using an analogous procedure to that described in Example 248, 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (268 mg, 0.833 mmol), (prepared as described for the starting material in Example 9), was reacted with 6-fluoro-5-hydroxy-2-methylindole (165 mg, 1 mmol) in DMF (3.5 ml) containing potassium carbonate (173 mg, 1.25 mmol) to give **4-(6-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (215 mg, 57 %).

MS-ESI : 451 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.65-1.8 (br s, 4H) ; 2.02 (m, 2H) ; 2.4 (s, 3H) ; 2.48 (br s, 4H) ; 2.6 (t, 2H) ; 4.02 (s, 3H) ; 4.3 (t, 2H) ; 6.18 (s, 1H) ; 7.25 (d, 1H) ; 7.4 (s, 1H) ; 7.45 (d, 1H) ; 7.6 (s, 1H) ; 8.5 (s, 1H)

Elemental analysis

Found

C 65.6 H 6.1 N 12.2

15 $C_{25}H_{27}FN_4O_3 \cdot 0.4 H_2O$

Requires

C 65.6 H 6.1 N 12.2%

The starting material was prepared as follows:


To a solution of 6-fluoro-5-methoxy-2-methylindole (1.23 g, 6.86 mmol), (prepared as described for the starting material in Example 237), in methylene chloride (15 ml) cooled at -30°C was added a solution of boron tribromide (3.78 g, 15.1 mmol) in methylene chloride (2 ml). After stirring for 90 minutes at ambient temperature, the mixture was poured onto ice and diluted with methylene chloride. The pH of the aqueous layer was adjusted to 6. The organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with ethylacetate/petroleum ether (8/2) to give 6-fluoro-5-hydroxy-2-methylindole (905 mg, 80 %).

MS-ESI : 166 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.3 (s, 3H) ; 5.95 (s, 1H) ; 6.9 (d, 1H) ; 7.0 (d, 1H) ; 8.85 (s, 1H) ; 10.6 (s, 1H)

¹³C NMR Spectrum: (DMSO-d₆) 13.3 ; 97.4 (d) ; 98.3 ; 105.5 ; 124.5 ; 128.8 (d) ; 135.6 ; 138.5 (d) ; 148.3 (d).

Example 251



A mixture of 4-chloro-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline (232 mg, 0.662 mmol), (prepared as described for the starting material in Examples 176 or 244), and 4-fluoro-5-hydroxyindole (120 mg, 0.794 mmol), (prepared as described for the starting material in Example 242), in DMF (3 ml) containing potassium carbonate (137 mg, 1 mmol) was stirred at 95°C for 3 hours. After cooling, the residue was poured onto water (12 ml) and extracted with ethyl acetate. The organic layer was washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by reversed phase C₁₈ column chromatography eluting with methanol/ammonium carbonate (2g of ammonium carbonate/litre saturated with CO₂) (60/40 followed by 70/30). The fractions containing the expected product were combined and evaporated. The residue was dissolved in ethyl acetate, dried (MgSO₄) and the volatiles were removed under vacuum. The residue was triturated with ether, filtered and dried under vacuum to give **4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline** (130 mg, 42 %).

MS-ESI : 466 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.3-2.4 (m, 2H) ; 2.97 (s, 3H) ; 3.2-4.1 (m, 8H) ; 3.5 (t, 2H) ; 4.07 (s, 3H) ; 4.4 (t, 2H) ; 6.6 (d, 1H) ; 7.15 (t, 1H) ; 7.38 (d, 1H) ; 7.5 (d, 1H) ; 7.6 (s, 1H) ; 7.82 (s, 1H) ; 8.95 (s, 1H).

Elemental analysis

Found C 64.4 H 6.1 N 15.0

C₂₅H₂₈FN₅O₃

Requires C 64.5 H 6.1 N 15.0%

Example 252

A mixture of 6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline (600 mg, 1.43 mmol), (prepared as described in Example 70), 1-(2-chloroethyl)-pyrrolidine (292 mg, 1.72 mmol) in methanol (14 ml) containing sodium carbonate (262 mg, 4.3 mmol) and potassium iodide (48 mg, 0.29 mmol) was heated at 50°C for 20 hours. After cooling, the volatiles were removed under vacuum. The residue was purified by preparation HPLC on reverse C₁₈ silica eluting with methanol/aqueous ammonium

carbonate (2g ammonium carbonate per litre saturated with CO₂) (60/40 followed by 70/30). The fractions containing the expected product were combined and the volatiles were removed under vacuum. The residue was triturated with ether and the solid was filtered, washed with ether and dried under vacuum to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-(2-**

5 **(pyrrolidin-1-yl)ethyl)-piperidin-4-ylmethoxy)quinazoline** (102 mg, 20 %).

MS-ESI : 516 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.3-1.5 (m, 2H) ; 1.6-1.75 (m, 4H) ; 1.8 (d, 2H) ; 1.7-1.9 (m, 1H) ; 1.95 (t, 2H) ; 2.45 (s, 3H) ; 2.4-2.5 (m, 5H) ; 2.95 (d, 2H) ; 3.35 (d, 2H) ; 4.0 (s, 3H) ; 4.1 (d, 2H) ; 6.18 (s, 1H) ; 6.9 (d, 1H) ; 7.25 (s, 1H) ; 7.35 (d, 1H) ; 7.38 (s, 1H) ; 7.6 (s, 1H) ; 8.5 (s, 1H) ; 11.05 (s, 1H)

Elemental analysis	Found	C 68.6	H 7.2	N 13.3
C ₃₀ H ₃₇ N ₅ O ₃ 0.5 H ₂ O	Requires	C 68.7	H 7.3	N 13.4%

Example 253

15 A mixture of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (110 mg, 0.325 mmol), (prepared as described for the starting material in Example 1), and 6-fluoro-5-hydroxyindole (59 mg, 0.39 mmol), (prepared as described for the starting material in Example 242), in DMF (1.8 ml) containing potassium carbonate (67 mg, 0.487 mmol) was heated at 90°C for 2 hours. After cooling, water was added. The solid was separated and
20 triturated with methanol. Water was added and the solid was filtered, washed with water and dried under vacuum to give **4-(6-fluoroindol-5-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline** (55 mg, 41 %).

MS-ESI : 453 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.95-2.05 (m, 2H) ; 2.45 (br s, 4H) ; 2.5 (t, 2H) ; 3.62 (t, 4H) ; 4.02 (s, 3H) ; 4.3 (t, 2H) ; 6.5 (s, 1H) ; 7.4 (d, 1H) ; 7.45 (s, 1H) ; 7.47 (t, 1H) ; 7.58 (d, 1H) ; 7.62 (s, 1H) ; 8.5 (s, 1H)

Elemental analysis	Found	C 61.6	H 5.5	N 11.9
C ₂₄ H ₂₅ FN ₄ O ₄ 0.8 H ₂ O	Requires	C 61.7	H 5.7	N 12.0%

30 Example 254

To a solution of 7-hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (183 mg, 0.6 mmol), (prepared as described for the starting material in Example 107), triphenylphosphine

(235 mg, 0.89 mmol) and 4-(2-hydroxyethyl)morpholine (93 mg, 0.72 mmol) in methylene chloride (4 ml) cooled at 10°C was added diethyl azodicarboxylate (140 μ l, 0.89 mmol).

After stirring at ambient temperature for 3 hours, the mixture was left overnight at 5°C. The mixture was poured onto a column of silica and eluted with methylene chloride followed by

5 methanol/methylene chloride (2/98) followed by 3N ammonia methanol/methylene chloride (2/98). The fractions containing the expected products were combined and evaporated to give
4-(indol-5-yloxy)-6-methoxy-7-(2-morpholinoethoxy)quinazoline (137 mg, 55 %).


MS-ESI : 421 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 3.30 (t, 2H) ; 3.65 (d, 2H) ; 3.7-3.8 (m, 4H) ; 4.05
 10 (d, 2H) ; 4.1 (s, 3H) ; 4.7 (t, 2H) ; 6.5 (s, 1H) ; 7.05 (dd, 1H) ; 7.4-7.6 (m, 3H) ; 7.65 (s, 1H) ;
 7.82 (s, 1H) ; 9.0 (s, 1H)


Examples 255-257

Using an analogous procedure to that described in Example 254, 7-hydroxy-4-(indol-
 15 5-yloxy)-6-methoxyquinazoline (183 mg, 0.6 mmol), (prepared as described for the starting material in Example 107), was used to prepare the compounds in Table XVIII.

Table XVIII



Example number	Weight (mg)	Yield %	MS-ESI [MH] ⁺	R	Note
255	123	51	405		a
256	124	48	434		b

257	165	62	448		c
-----	-----	----	-----	--	---

a) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (183 mg, 0.6 mmol) was reacted with 1-(2-hydroxyethyl)pyrrolidine (82 mg) to give **4-(indol-5-yloxy)-6-methoxy-7-(2-(pyrrolidin-1-yl)ethoxy)quinazoline**.

5 ¹H NMR Spectrum: (DMSO-d₆) 1.72 (br s, 4H) ; 2.6 (br s, 4H) ; 2.9 (t, 2H) ; 4.0 (s, 3H) ; 4.3 (t, 2H) ; 6.48 (s, 1H) ; 7.0 (dd, 1H) ; 7.4-7.5 (m, 3H) ; 7.6 (s, 1H) ; 8.5 (s, 1H) ; 11.3 (br s, 1H)

b) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (183 mg, 0.6 mmol) was reacted with 4-(2-hydroxyethyl)-1-methylpiperazine (103 mg) to give **4-(indol-5-yloxy)-6-methoxy-7-(2-(4-methylpiperazin-1-yl)ethoxy)quinazoline**.

10 ¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.5 (s, 3H) ; 3.35 (t, 2H) ; 3.65 (d, 2H) ; 3.7-3.8 (m, 4H) ; 4.05 (d, 2H) ; 4.1 (s, 3H) ; 4.7 (t, 2H) ; 7.05 (dd, 1H) ; 7.45 (s, 1H) ; 7.5-7.6 (m, 2H) ; 7.65 (s, 1H) ; 7.82 (s, 1H) ; 9.0 (s, 1H)

15 The starting material was prepared as follows:-

2-Bromoethanol (2.36g, 19mmol) was added dropwise to a mixture of 1-methylpiperazine (1.26g, 13mmol) and potassium carbonate (5.0g, 36mmol) in absolute ethanol (150ml) and the mixture heated at reflux for 18 hours. The mixture was allowed to cool and the precipitates were removed by filtration and the solvent volatiles were removed by

20 evaporation. The residue was treated with acetone/methylene chloride, the insolubles were removed by filtration and the solvent was removed from the filtrate by evaporation to give 4-(2-hydroxyethyl)-1-methylpiperazine (870mg, 48%) as a light brown oil.

¹H NMR Spectrum: (CDCl₃) 2.18(s, 3H); 2.3-2.7(br m, 8H); 2.56(t, 2H); 3.61(t, 2H)

MS - ESI: 145 [MH]⁺

25 c) 7-Hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (183 mg, 0.6 mmol) was reacted with 1-(3-hydroxypropyl)-4-methylpiperazine (113 mg), (prepared as described for the starting material in Example 133), to give **4-(indol-5-yloxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline**.

¹H NMR Spectrum: (DMSO-d₆) 2.15 (s, 3H) ; 2.3-2.4 (br s, 4H) ; 2.5-2.6 (m, 4H) ; 2.8 (t, 2H) ; 4.0 (s, 3H) ; 4.35 (t, 2H) ; 6.45 (s, 1H) ; 7.0 (dd, 1H) ; 7.4-7.5 (m, 4H) ; 7.62 (s, 1H) ; 8.5 (s, 1H)

5 Example 258

A solution of (2*R*)-7-(2-acetoxy-3-(pyrrolidin-1-yl)propoxy)-4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxyquinazoline (570 mg, 1.12 mmol) in methanol saturated with ammonia (7 ml) was stirred overnight at ambient temperature. The volatiles were removed under vacuum and the residue was purified by column chromatography eluting with
10 methylene chloride/methanol containing ammonia (approximately 3*N*) to give (2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxyquinazoline (390mg ; 75%).

MS-ESI : 467 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.85-2.0 (m, 2H) ; 2.0-2.15 (m, 2H) ; 2.42 (s, 3H)
15 ; 3.15 (m, 2H) ; 3.4 (d, 2H) ; 3.65 (m, 2H) ; 4.1(s, 3H) ; 4.32 (d, 2H) ; 4.4 (m, 1H) ; 7.05 (dd, 1H) ; 7.22 (d, 1H) ; 7.6 (s, 1H) ; 7.85 (s, 1H) ; 9.02 (s, 1H)

The starting material was prepared as follows:

A suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.2 g, 3.91 mmol), (prepared as described for the starting material
20 in Example 12), and 2-(*R*)-(-)-Glycidyl tosylate (1.25 g, 5.47 mmol) in DMF (10 ml) containing potassium carbonate (1.61 g, 11.7 mmol) was stirred at 60°C for 4 hours. After cooling, the mixture was filtered and the solid was washed with DMA. The filtrate was evaporated and the residue was partitioned between ethyl acetate and aqueous ammonia. The
25 organic layer was separated, washed with water, brine, dried (MgSO₄) and evaporated. The residue was purified by column chromatography eluting with ethyl acetate. The fractions containing the expected product were combined and evaporated to give (2*R*)-7-(oxiran-2-ylmethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.21 g, 85 %).
MS-ESI : 363 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.12 (s, 9H) ; 2.75 (m, 1H) ; 2.9 (t, 1H) ; 3.4 (m, 1H) ; 3.93 (s, 3H) ; 4.0 (dd, 1H) ; 4.52 (dd, 1H) ; 5.9 (s, 2H) ; 7.2 (s, 1H) ; 7.52 (s, 1H) ; 8.35 (s, 1H)
30

A solution of (2*R*)-7-(oxiran-2-ylmethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (1.1 g, 3 mmol) and pyrrolidine (216 mg, 3 mmol) in trichloromethane (15 ml) was refluxed for 11 hours. The volatiles were removed under vacuum and the residue was purified by column chromatography eluting with methylene chloride/methanol (85/15 followed by 70/30) to give (2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-3-((pivaloyloxymethyl)-3,4-dihydroquinazolin-4-one (1.18 g, 90 %).
¹H NMR Spectrum: (DMSO-d₆) 1.15 (s, 9H) ; 1.7 (br s, 4H) ; 2.48 (m, 1H) ; 2.5 (br s, 4H) ; 2.65 (dd, 1H) ; 3.9 (s, 3H) ; 4.0 (br s, 1H) ; 4.05 (dd, 1H) ; 4.18 (dd, 1H) ; 4.95 (br s, 1H) ; 5.9 (s, 2H) ; 7.2 (s, 1H) ; 7.5 (s, 1H) ; 8.35 (s, 1H)

A solution of (2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-3-((pivaloyloxymethyl)-3,4-dihydroquinazolin-4-one (778 mg, 1.8 mmol) in methanol saturated with ammonia (20 ml) was stirred for 24 hours at ambient temperature. The volatiles were removed under vacuum. The residue was triturated with ether and the residue was filtered, washed with ether and dried under vacuum to give (2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (800 mg, quant.).
¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.92 (m, 2H) ; 2.05 (m, 2H) ; 3.15 (m, 2H) ; 3.35 (d, 2H) ; 3.62 (m, 2H) ; 3.98 (s, 3H) ; 4.18 (d, 2H) ; 4.32 (m, 1H) ; 7.35 (s, 1H) ; 7.6 (s, 1H) ; 9.2 (s, 1H)

A mixture of (2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (803 mg, 2.51 mmol) in acetic anhydride (1.2 ml, 12.5 mmol) was stirred at ambient temperature for 1 hour. Water (360 µl, 20 mmol) was added and stirring was continued for 90 minutes. The mixture was partitioned between aqueous sodium hydrogen carbonate and methylene chloride. The organic layer was separated, washed with brine, dried (MgSO₄) and evaporated. The residue was triturated with ether, filtered and dried under vacuum to give (2*R*)-7-(2-acetoxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (595 mg, 65%).

MS-ESI : 362 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.7 (br s, 4H) ; 2.05 (s, 3H) ; 2.5 (br s, 4H) ; 2.72 (m, 2H) ; 3.9 (s, 3H) ; 4.3 (m, 2H) ; 5.25 (m, 1H) ; 7.2 (s, 1H) ; 7.45 (s, 1H) ; 8.0 (s, 1H)

A solution of (2*R*)-7-(2-acetoxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (556mg, 1.54 mmol) in thionyl chloride (6 ml) containing DMF (3 drops) was heated at 80°C for 4 hours. The volatiles were removed under vacuum. The

residue was dissolved in methylene chloride and the organic layer was washed with aqueous sodium hydrogen carbonate, brine, dried (MgSO₄) and evaporated to give (2*R*)-7-(2-acetoxy-3-(pyrrolidin-1-yl)propoxy)-4-chloro-6-methoxyquinazoline (530mg, 90%).

¹H NMR Spectrum: (DMSO-d₆) 1.7 (br s, 4H) ; 2.05 (s, 3H) ; 2.55 (br s, 4H) ; 2.75 (br s, 2H) ;
5 4.02 (s, 3H) ; 4.35-4.5 (m, 2H) ; 5.3 (m, 1H) ; 7.4 (s, 1H) ; 7.5 (s, 1H) ; 7.9 (s, 1H)

A suspension of (2*R*)-7-(2-acetoxy-3-(pyrrolidin-1-yl)propoxy)-4-chloro-6-methoxyquinazoline (530 mg, 1.4 mmol) and 4-fluoro-5-hydroxy-2-methylindole (277 mg, 1.68 mmol), (prepared as described for the starting material in Example 237), in DMF (8 ml) containing potassium carbonate (290 mg, 2.1 mmol) was stirred at 90°C for 2 hours. After
10 cooling, the volatiles were removed under vacuum and the residue was purified by column chromatography eluting with methylene chloride/methanol (95/5) to give (2*R*)-7-(2-acetoxy-3-(pyrrolidin-1-yl)propoxy)-4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxyquinazoline (580 mg, 81 %).

¹H NMR Spectrum: (DMSO-d₆) 1.7 (br s, 4H) ; 2.05 (s, 3H) ; 2.4 (s, 3H) ; 2.52 (br s, 4H) ;
15 2.65-2.82 (m, 2H) ; 4.0 (s, 3H) ; 4.4 (m, 2H) ; 5.3 (m, 1H) ; 6.25 (s, 1H) ; 7.0 (dd, 1H) ; 7.18 (d, 1H) ; 7.48 (s, 1H) ; 7.62 (s, 1H) ; 8.5 (s, 1H)

Example 259


A solution of 4-chloro-6-methoxy-7(3-(pyrrolidin-1-yl)propoxy)quinazoline (61 mg, 0.19 mmol), (prepared as described for the starting material in Example 9), and 5-aminoindole
20 (30 mg, 0.23 mmol) in isopropanol (2 ml) containing 6.2 N hydrogen chloride in isopropanol (33 µl) was heated at 80°C for 6 hours. After cooling, the precipitate was filtered, washed with ether and dried under vacuum to give 4-(indol-5-ylamino)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline hydrochloride (80 mg, 72 %).

25 MS - ESI : 418 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 1.9 (m, 2H) ; 2.05 (m, 2H) ; 2.3 (m, 2H) ; 3.1 (m, 2H) ; 3.4 (t, 2H) ; 3.65 (m, 2H) ; 4.05 (s, 3H) ; 4.35 (t, 2H) ; 6.5 (s, 0.5H, partly exchanged) ;
7.3 (d, 1H) ; 7.4 (s, 1H) ; 7.45 (s, 1H) ; 7.55 (d, 1H) ; 7.8 (s, 1H) ; 8.25 (s, 1H) ; 8.8 (s, 1H)

Example 260-265

Using an analogous procedure to that described in Example 259, 5-aminoindole (30 mg, 0.23 mmol) was used in the synthesis of the compounds described in Table XIX.

Table XIX

Example number	Weight (mg)	Yield (%)	MS-ESI [MH] ⁺	Note	R
260	101	76	510	a	
261	92	83	418	b	
262	92	80	434	c	
263	84	80	427	d	
264	78	79	401	e	
265	72	70	416	f	

- 5 a) 4-Chloro-6-methoxy-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)quinazoline (78 mg), (prepared as described for the starting material in Example 12), was reacted with 5-aminoindole to give 4-(indol-5-ylamino)-6-methoxy-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)quinazoline hydrochloride.

¹H NMR Spectrum: (DMSO-d₆) 1.65-1.8 (m, 2H) ; 2.05 (d, 2H) ; 2.2 (br s, 1H) ; 3.1 (br s, 2H)
 10 ; 3.2 (s, 3H) ; 3.5 (br s, 2H) ; 3.6 (d, 2H) ; 3.8 (m, 2H) ; 4.05 (s, 3H) ; 4.1 (d, 2H) ; 6.5 (s, 1H)

; 7.3 (d, 1H) ; 7.42 (m, 2H) ; 7.5 (d, 1H) ; 7.8 (s, 1H) ; 8.4 (s, 1H) ; 8.7 (s, 1H) ; 11.15 (br s, 1H) ; 11.32 (s, 1H) . 11.5 (s, 1H).

b) 4-Chloro-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline (61 mg), (prepared as described for the starting material in Example 10), was reacted with 5-aminoindole to give **4-(indol-5-ylamino)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline hydrochloride**.

¹H NMR Spectrum: (DMSO-d₆) 1.6-1.8 (m, 2H) ; 2.02 (d, 2H) ; 2.15 (br s, 1H) ; 2.75 (s, 3H) ; 3.0 (br s, 2H) ; 3.45 (d, 2H) ; 4.02 (s, 3H) ; 4.1 (d, 2H) ; 6.5 (s, 1H) ; 7.3 (d, 1H) ; 7.4 (m, 2H) ; 7.5 (d, 1H) ; 7.8 (s, 1H) ; 8.3 (s, 1H) ; 8.7 (s, 1H) ; 10.4 (br s, 1H) ; 11.3 (s, 1H)

The presence of a second form of the piperidine ring (due to protonation effects) is detectable in the NMR Spectrum as a doublet at 4.3 ppm (approximately 20 % of parent compound).

c) 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (64 mg), (prepared as described for the starting material in Example 1), was reacted with 5-aminoindole to give **4-(indol-5-ylamino)-6-methoxy-7-(3-morpholinopropoxy)quinazoline hydrochloride**.

¹H NMR Spectrum (DMSO-d₆ ; CF₃COOD) : 2.35 (m, 2H) ; 3.15 (t, 2H) ; 3.3 (t, 2H) ; 3.57 (d, 2H) ; 3.8 (m, 2H) ; 4.02 (d, 2H) ; 4.03 (s, 3H) ; 4.3 (t, 2H) ; 6.5 (d, 1H) ; 7.3 (dd, 1H) ; 7.4 (s, 1H) ; 7.45 (s, 1H) ; 7.52 (d, 1H) ; 7.8 (s, 1H) ; 8.25 (s, 1H) ; 8.78 (s, 1H)

d) 4-Chloro-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline (62 mg), (prepared as described for the starting material in Example 50), was reacted with 5-aminoindole in the presence of 6.2N hydrogen chloride in isopropanol (4 µl) to give **4-(indol-5-ylamino)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline hydrochloride**.

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.2-2.4 (m, 2H) ; 3.07 (s, 3H) ; 3.35 (t, 2H) ; 4.05 (s, 3H) ; 4.35 (t, 2H) ; 6.5 (d, 0.5 H, partly exchanged) ; 7.2-7.35 (m, 2H) ; 7.45 (s, 1H) ; 7.5 (d, 1H) ; 7.8 (s, 1H) ; 8.2 (s, 1H) ; 8.75 (s, 1H)

e) 4-Chloro-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline (58 mg) was reacted with 5-aminoindole in the presence of 6.2N hydrogen chloride in isopropanol (4 µl) to give **4-(indol-5-ylamino)-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline hydrochloride**.

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 4.03 (s, 3H) ; 4.65 (t, 2H) ; 4.8 (t, 2H) ; 6.5 (d, 1H; partly exchanged) ; 7.30 (d, 1H) ; 7.4 (s, 1H) ; 7.45 (s, 1H) ; 7.52 (d, 1H) ; 7.75 (s, 1H) ; 7.8 (s, 1H) ; 7.9 (s, 1H) ; 8.25 (s, 1H) ; 8.75 (s, 1H) ; 9.25 (s, 1H)

5 The starting material was prepared as follows:

Diethyl azodicarboxylate (435mg, 2.5mmol) was added dropwise to a suspension of 7-hydroxy-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (612mg, 2mmol), (prepared as described for the starting material in Example 12), 2-(imidazol-1-yl)ethanol (280mg, 2.5mmol), (J. Med. Chem. 1993, **25** 4052-4060), and triphenylphosphine (655mg, 10 2.5mmol) in methylene chloride (10ml) at 5°C. The mixture was stirred for 10 minutes at 5°C and then 1 hour at ambient temperature. The mixture was poured directly on to a silica column and eluted with methylene chloride/methanol (95/5) to give 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (640mg, 80%).

¹H NMR Spectrum: (CDCl₃) 1.19(s, 9H); 3.98(s, 3H); 4.34(m, 2H); 4.45(m, 2H); 5.94(s, 15 2H); 7.02(s, 1H); 7.07(s, 1H); 7.11(s, 1H); 7.64(s, 1H); 7.67(s, 1H); 8.17(s, 1H)

MS - ESI: 423 [MNa]⁺

Elemental Analysis:	Found	C 58.3	H 6.4	N 13.9
---------------------	-------	--------	-------	--------

C ₂₀ H ₂₄ N ₄ O ₅ 0.7H ₂ O	Requires	C 58.2	H 6.2	N 13.6%
---	----------	--------	-------	---------

A solution of 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-3-((pivaloyloxy)methyl)-3,4-dihydroquinazolin-4-one (640mg, 1.6mmol) in saturated methanolic ammonia (10ml) was 20 stirred for 15 hours at ambient temperature. The volatiles were removed by evaporation, the solid was triturated with ether, collected by filtration and dried under vacuum to give 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-3,4-dihydroquinazolin-4-one (412mg, 90%).

¹H NMR Spectrum: (DMSO-d₆) 3.89(s, 3H); 4.4-4.5(m, 4H); 6.9(s, 1H); 7.16(s, 1H); 7.28(s, 25 1H); 7.47(s, 1H); 7.7(s, 1H); 7.99(s, 1H)

MS - ESI: 287 [MH]⁺

Elemental Analysis:	Found	C 57.8	H 5.2	N 19.3
---------------------	-------	--------	-------	--------

C ₁₄ H ₁₄ N ₄ O ₃ 0.3H ₂ O	Requires	C 57.7	H 5.1	N 19.2%
---	----------	--------	-------	---------

A mixture of 7-(2-(imidazol-1-yl)ethoxy)-6-methoxy-3,4-dihydroquinazolin-4-one 30 (412mg, 1.44mmol), thionyl chloride (5 ml) and DMF (0.2ml) was heated at reflux for 1 hour. The mixture was diluted with toluene and the volatiles were removed by evaporation. The

residue was suspended in methylene chloride, cooled to 0°C and aqueous sodium hydrogen carbonate solution was added. The resulting precipitate was collected by filtration and dried under vacuum to give 4-chloro-7-(2-(imidazol-1-yl)ethoxy)-6-methoxyquinazoline (258mg, 59%).

5 ¹H NMR Spectrum: (DMSO-d₆) 4.01(s, 3H); 4.47(m, 2H); 4.53(m, 2H); 6.89(s, 1H); 7.27(s, 1H); 7.41(s, 1H); 7.49(s, 1H); 7.70(s, 1H); 8.88(s, 1H)

MS - ESI: 327 [MNa]⁺

10 f) 4-Chloro-6-methoxy-7-(3-(1*H*-1,2,4-triazol-1-yl)propoxy)quinazoline (61 mg) was reacted with 5-aminoindole in the presence of 6.2N hydrogen chloride in isopropanol (4 µl) to give 4-(indol-5-ylamino)-6-methoxy-7-(3-(1*H*-1,2,4-triazol-1-yl)propoxy)quinazoline hydrochloride.

¹H NMR Spectrum: (DMSO-d₆, CF₃COOD) 2.5 (m, 2H) ; 4.0 (s, 3H) ; 4.3 (t, 2H) ; 4.6 (t, 2H) ; 6.52 (d, 0.5H partly exchanged) ; 7.3 (s, 1H) ; 7.35 (d, 1H) ; 7.45 (s, 1H) ; 7.55 (d, 1H) ; 7.8 (s, 1H) ; 8.16 (s, 1H) ; 8.66 (s, 1H) ; 8.77 (s, 1H) ; 9.43 (s, 1H)

Example 266

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (144mg, 0.43mmol), (prepared as described for the starting material in Example 67), potassium carbonate (91mg, 0.66mmol) and 3-fluoro-7-hydroxyquinoline (77mg, 0.47mmol), (prepared as described for the starting material in Example 157), in DMF (3ml) was stirred at 100°C for 2 hours and then allowed to cool to ambient temperature. The reaction mixture was evaporated to dryness and the residue chromatographed on silica eluting with methanol/dichloromethane/aqueous ammonia (0.880) (5/100/1). The relevant fractions were combined and evaporated to dryness to give 4-(3-fluoro-quinolin-7-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline (87mg, 44%).

¹H NMR Spectrum: (DMSO-d₆) 1.37(m, 2H); 1.49(m, 4H); 1.96(m, 2H); 2.34(m, 4H); 2.43(t, 2H); 4.00(s, 3H); 4.23(t, 2H); 7.38(s, 1H); 7.62(s, 1H); 7.69(dd, 1H); 8.00(d, 1H); 8.12(d, 1H); 8.34(dd, 1H); 8.54(s, 1H); 8.98(d, 1H)

30 MS (ESI): 463 (MH)⁺

Example 267

A mixture of 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (218mg, 0.68mmol), (prepared as described for the starting material in Example 9), potassium carbonate (138mg, 1.13mmol) and 3-fluoro-7-hydroxyquinoline (117mg, 0.72mmol),
5 (prepared as described for the starting material in Example 157), in DMF (4.5ml) was stirred at 100°C for 4 hours and then allowed to cool to ambient temperature. The reaction mixture was evaporated to dryness and the residue taken up in dichloromethane, washed with water, brine and dried (MgSO₄). The organic fractions were evaporated to dryness and the residue recrystallised from acetonitrile to give **4-(3-fluoro-quinolin-7-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (86mg, 28%).

¹H NMR Spectrum: (DMSO-d₆) 1.90(m, 2H); 2.00(m, 2H); 2.27(m, 2H); 3.02(m, 2H); 3.32(m, 2H); 3.59(m, 2H); 4.00(s, 3H); 4.33(t, 2H); 7.43(s, 1H); 7.62(s, 1H); 7.70(dd, 1H); 7.99(d, 1H); 8.11(d, 1H); 8.35(dd, 1H); 8.54(s, 1H); 8.97(d, 1H)

MS (ESI): 449 (MH)⁺

Example 268

A mixture of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (280 mg, 0.87 mmol), (prepared as described in Example 49), potassium carbonate (370 mg, 2.68 mmol) and 4-(1-methyl-2-oxopiperidin-4-yl)methyl-4-toluene sulphonate (260 mg, 0.87 mmol) in DMF (8 ml) was stirred at 95°C for 4 hours and allowed to cool to ambient temperature. The reaction mixture was diluted with acetone, filtered and the filtrate evaporated 'in vacuo' to give a residue which was purified by column chromatography, eluting with dichloromethane/methanol/0.88 ammonia (100/8/1). The relevant fractions were combined and evaporated 'in vacuo' to give an oil which crystallised on trituration with diethyl ether to give **6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-methyl-2-oxopiperidin-4-ylmethoxy)quinazoline** (66 mg, 17%).

m. p. 250 – 251°C

¹H NMR Spectrum: (DMSO-d₆) 1.66 (m, 1H), 2.10 (m, 2H), 2.40 (s, 3H), 2.50 (m, 2H), 2.84 (s, 3H), 3.34 (m, 2H), 3.99 (s, 3H), 4.12 (d, 2H), 6.12 (s, 1H), 6.86 (m, 1H),

7.25 (d, 1H), 7.30 (d, 1H), 7.38 (s, 1H), 7.59 (s, 1H), 8.48 (s, 1H) and 10.98 (br s, 1H).

MS (ESI) : 447 (MH)⁺

Elemental analysis Found C 66.8 H 5.9 N 12.4

$C_{25}H_{26}N_4O_4 \cdot 0.2 H_2O$ Requires C 66.7 H 5.9 N 12.5%

The starting material was prepared as follows:-

- A solution of 4-hydroxymethyl-1-methyl-2-piperidone (120 mg, 0.84 mmol),
 5 (Yakugaku Zasshi 88, (5), 573 - 582, (1968)), in dichloromethane was treated with triethylamine (187 mg, 1.85 mmol) followed by p-toluenesulphonyl chloride (176 mg, 0.92 mmol) and the mixture stirred at ambient temperature overnight. The reaction mixture was diluted with dichloromethane and washed successively with aqueous sodium hydrogen carbonate, water and brine. The dichloromethane solution was dried over magnesium
 10 sulphate, filtered and the filtrate evaporated '*in vacuo*' to give a dark oily residue. This was washed several times with diethyl ether to remove the product from insoluble impurities, the washings combined and evaporated '*in vacuo*' to give 4-(1-methyl-2-oxopiperidin-4-yl)methyl-4-toluene sulphonate as a light brown oil (130 mg, 52%). This was used without further purification.
 15 MS (ESI) : 298 (MH)⁺ and impurities

Example 269

- A mixture of (2*R*)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.79 mmol), and 1-methylpiperazine (0.26 ml, 2.38 mmol)
 20 in DMF (10 ml) was stirred at 70° C for 24 hours and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography, gradient elution (dichloromethane, 5% methanol/95% dichloromethane, dichloromethane/methanol/0.88 ammonia (100/8/1) and evaporated *in vacuo* to give (2*R*)-7-(2-hydroxy-3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-
 25 yloxy)quinazoline (344 mg, 91 %).
¹H NMR Spectrum: (DMSO-*d*₆) 2.10 (s, 3H), 2.4 (m, 13H), 3.98 (s, 3H), 4.06 (m, 3H), 4.90 (br s, 1H), 6.12 (s, 1H), 6.85 (dd, 1H), 7.3 (m, 2H), 7.58 (s, 1H), 8.42 (s, 1H) and 10.98 (br s, 1H)
 MS (ESI) : 478 (MH)⁺
 30 Elemental analysis: Found C 61.3 H 6.3 N 13.8
 $C_{26}H_{30}N_4O_4 \cdot 0.2H_2O \cdot 0.5dichloromethane$ Requires C 61.9 H 6.2 N 13.4%

The starting material was prepared as follows:

A mixture of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (300 mg, 0.93 mmol), (prepared as described in Example 49), potassium carbonate (385 mg, 2.79 mmol) and (2R)-(-)- glycidyl tosylate (426 mg, 2.79 mmol) in DMF (15 ml) was stirred at 60°C for 2 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate exaporated *in vacuo*. The residue was dissolved in dichloromethane and washed with saturated sodium hydrogen carbonate solution. The organic layer was then dried (MgSO₄), filtered and the solvent removed *in vacuo* to give a yellow solid. This was triturated with ether, filtered off and dried to give (2R)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline as a yellow solid (185 mg, 53 %).

¹H NMR Spectrum: (DMSOd₆) 2.40 (s, 3H), 2.75 (m, 1H), 2.90 (m, 1H), 3.40 (m, 1H), 3.98 (s, 3H), 4.05 (m, 1H), 4.60 (m, 1H), 6.15 (s, 1H), 6.85 (dd, 1H), 7.30 (m, 2H) 7.40 (s, 1H), 7.60 (s, 1H), 8.45 (s, 1H) and 10.98 (s, 1H)

MS (ESI) : 378 (MH)⁺

Example 270

A mixture of (2R)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.79 mmol), (prepared as described for the starting material in Example 269), and diethylamine (0.25 ml, 2.38 mmol) in DMF (10 ml) was stirred at 70 °C for 24 hours and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography, gradient elution (dichloromethane, 5% methanol/95% dichloromethane, dichloromethane/methanol/0.88 ammonia (100/8/1)) to give (2R)-7-(3-(N,N-diethylamino)-2-hydroxypropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (288 mg, 81 %).

¹H NMR Spectrum: (DMSO-d₆) 0.95 (t, 6H), 2.10 (s, 3H), 2.4 (m, 6H), 3.98 (s, 3H), 4.14 (m, 3H), 4.84 (br s, 1H), 6.12 (s, 1H), 6.85 (dd, 1H), 7.3 (m, 3H), 7.58 (s, 1H), 8.42 (s, 1H) and 10.98 (br s, 1H)

MS (ESI) : 448 (MH)⁺

Elemental analysis:	Found	C 64.3	H 6.6	N 12.0
C ₂₅ H ₃₀ N ₄ O ₄ 0.4dichloromethane	Requires	C 64.0	H 6.4	N 11.6%

Example 271

A mixture of 7-benzyloxy-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline (7.76 g, 18.9 mmol), ammonium formate (17.82 g, 282 mmol) and 10% palladium on charcoal (800 mg) in DMF (350 ml) was stirred at ambient temperature for 1 hour. The catalyst was filtered off through celite and the cake washed with DMF. The solvent was removed *in vacuo* and the residue stirred with a saturated solution of sodium hydrogen carbonate for 2 hours. The suspension was then filtered, washed with water and dried to give **7-hydroxy-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (5.49 g, 91%).

¹H NMR Spectrum: (DMSO-d₆) 2.20 (s, 3H), 3.98 (s, 3H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.20 (s, 1H), 7.35 (m, 3H), 7.58 (s, 1H), 8.40 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 322 (MH)⁺

The starting material was prepared as follows:

A mixture of 7-benzyloxy-4-chloro-6-methoxyquinazoline (7.859 g, 26.1 mmol), (prepared as described for the starting material in Example 1), potassium carbonate (18.03 g, 130 mmol) and 5-hydroxy-3-methylindole (5.00 g, 34.0 mmol), (Journal of Organic Chemistry 1993, 58, 3757), in DMA (600 ml) was stirred at 75 °C for 2 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*. The crude solid was purified by silica column chromatography, eluting with 2.5% methanol/97.5% dichloromethane to give 7-benzyloxy-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline (7.791 g, 73 %).

¹H NMR Spectrum: (CDCl₃) 2.30 (s, 3H), 4.10 (s, 3H), 5.36 (s, 2H), 7.04 (m, 2H), 7.43 (m, 8H), 7.62 (s, 1H), 8.02 (s, 1H), and 8.60 (s, 1H)

MS (ESI) : 412 (MH)⁺

Example 272

A mixture of 7-hydroxy-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline (800 mg, 2.49 mmol), (prepared as described in Example 271), potassium carbonate (687 mg, 4.98 mmol) and 1-chloro-3-morpholinopropane (448 mg, 2.74 mmol), (prepared as described for the starting material in Example 1), in DMF (20 ml) was stirred at 80 °C for 2 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by silica column chromatography, gradient

elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) and the product was recrystallised from ethanol to give **6-methoxy-4-(3-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline** (570 mg, 51 %).

5 ¹H NMR Spectrum: (DMSO-d₆) 1.98 (m, 2H), 2.20 (s, 3H), 2.40 (t, 4H), 2.50 (m, 2H), 3.60 (t, 4H), 3.98 (s, 3H), 4.20 (t, 2H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.45 (s, 1H), and 10.82 (br s, 1H)

MS (ESI) : 449 (MH)⁺

Elemental analysis:

Found C 64.2 H 6.0 N 11.8

10 C₂₅H₂₈N₄O₄ 0.7H₂O 0.7ethanol

Requires C 64.2 H 6.9 N 11.4%

Example 273

A mixture of 7-hydroxy-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline (800 mg, 2.49 mmol), (prepared as described for the starting material in Example 271), potassium carbonate (1.031 g, 7.47 mmol) and 4-(2-chloroethyl)morpholine hydrochloride (510 mg, 2.74 mmol) in DMF (25 ml) was stirred at 80 °C for 2 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate removed *in vacuo*. The residue was purified by silica column chromatography, gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) and the product recrystallised from ethanol to give **6-methoxy-4-(3-methylindol-5-yloxy)-7-(2-morpholinoethoxy)quinazoline** (510 mg, 47 %).

15 ¹H NMR Spectrum: (DMSO-d₆) 2.20 (s, 3H), 2.55 (t, 4H), 2.80 (t, 2H), 3.60 (t, 4H), 3.98 (s, 3H), 4.30 (t, 2H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 2H), 7.40 (s, 1H), 7.60 (s, 1H), 8.45 (s, 1H), and 10.82 (br s, 1H)

25 MS (ESI) : 449 (MH)⁺

Elemental analysis:

Found C 64.1 H 6.3 N 12.2

C₂₄H₂₆N₄O₄ 0.4H₂O 0.8ethanol

Requires C 64.3 H 6.1 N 11.7%

Example 274

30 A mixture of 7-hydroxy-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline (1.00 g, 3.11 mmol), (prepared as described for the starting material in Example 271), potassium carbonate (1.288 g, 9.33 mmol) and 4-(4-methylphenylsulphonyloxymethyl)-1-*tert*-

butoxycarbonylpiperidine (1.264 g, 3.42 mmol), (prepared as described for the starting material in Example 10), in DMF (35 ml) was stirred at 80 °C for 2 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the solvent removed *in vacuo*. The residue was purified by silica column chromatography, 5% methanol/95%

5 dichloromethane and the product was recrystallised from ethanol to give **6-methoxy-4-(3-methylindol-5-yloxy)-7-(1-*tert*-butoxycarbonylpiperidin-4-ylmethoxy)quinazoline** (1.011 g, 63 %).

¹H NMR Spectrum: (DMSO-d₆) 1.3 (m, 4H), 1.42 (s, 9H), 1.90 (d, 2H), 2.10 (m, 1H), 2.28 (s, 3H), 2.80 (m, 2H), 3.98 (s, 3H), 4.08 (d, 2H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.45 (s, 1H), and 10.82 (br s, 1H)

10 MS (ESI) : 519 (MH)⁺

Example 275

A mixture of 7-hydroxy-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline (600 mg, 1.87 mmol), (prepared as described for the starting material in Example 271), potassium carbonate (773 mg, 5.60 mmol) and 3-(1,1-dioxothiomorpholino)propoxy tosylate (1.296 g, 3.74 mmol) in DMF (30 ml) was stirred at 75 °C overnight and allowed to cool to ambient temperature. The reaction mixture was filtered and the solvent removed *in vacuo*. The residue was purified by silica column chromatography, gradient elution (dichloromethane, 5% 20 methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) and the product recrystallised from ethanol to give **7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (525 mg, 56%).

¹H NMR Spectrum: (DMSO-d₆) 1.98 (m, 2H), 2.17 (s, 3H), 2.65 (t, 2H), 2.90 (t, 4H), 3.10 (t, 4H), 3.98 (s, 3H), 4.25 (t, 2H), 6.95 (dd, 1H), 7.15 (s, 1H), 7.30 (d, 1H), 7.35 (m, 2H), 7.60 (s, 1H), 8.45 (s, 1H), and 10.82 (br s, 1H)

25 MS (ESI) : 497 (MH)⁺

Elemental analysis: Found C 58.4 H 5.5 N 11.1

C₂₅H₂₈N₄O₅S 0.8H₂O Requires C 58.8 H 5.8 N 11.0%

Example 276

A mixture of 6-methoxy-4-(3-methylindol-5-yloxy)-7-(1-*tert*-butoxycarbonylpiperidin-4-ylmethoxy)quinazoline (1.290 g, 2.49 mmol), (prepared as described in Example 274), in 25% trifluoroacetic acid/75% dichloromethane solution (75 ml) was stirred at ambient temperature for 2 hours. The solvents were then removed *in vacuo* and the dark yellow gum triturated with concentrated ammonia. The resulting solid was filtered off and dried to give **6-methoxy-4-(3-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline** (648 mg, 62%).

¹H NMR Spectrum: (DMSO-d₆) 1.35 (m, 2H), 1.80 (m, 2H), 2.05 (m, 1H), 2.10 (s, 3H), 2.70 (m, 2H), 3.10 (m, 2H), 3.98 (s, 3H), 4.05 (d, 2H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.34 (m, 3H), 7.60 (s, 1H), 8.45 (s, 1H), and 10.82 (br s, 1H)

MS (ESI) : 419 (MH)⁺

Example 277

A mixture of 6-methoxy-4-(3-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline (460 mg, 1.10 mmol), (prepared as described in Example 276), triethylamine (5 ml) and chloroacetonitrile (0.38 ml, 6.05 mmol) in methanol (5 ml) was stirred at ambient temperature for 24 hours. The solvents were removed *in vacuo* and the residue purified by silica column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) and the product recrystallised from acetonitrile to give **7-(1-cyanomethylpiperidin-4-ylmethoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (178 mg, 35 %).

¹H NMR Spectrum: (DMSO-d₆) 1.40 (m, 2H), 1.80 (m, 4H), 2.20 (m, 4H), 2.81 (m, 2H), 3.65 (s, 2H), 3.98 (s, 3H), 4.05 (d, 2H), 6.98 (dd, 1H), 7.15 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.45 (s, 1H), and 10.83 (br s, 1H)

MS (ESI) : 458 (MH)⁺

Elemental analysis: Found C 66.3 H 6.1 N 14.8

C₂₆H₂₇N₃O₃ · 0.7H₂O Requires C 66.4 H 6.1 N 14.9%

Example 278

A mixture of 7-hydroxy-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline (1.35 g, 4.2 mmol), (prepared as described for the starting material in Example 271), potassium carbonate

(1.74 g, 12.6 mmol) and (2*R*)-(-)- glycidyl tosylate (1.92 g, 8.4 mmol) in DMF (25 ml) was stirred at 60 °C for 2 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the solvent removed *in vacuo*. The residue was dissolved in dichloromethane and washed with saturated sodium hydrogen carbonate solution. The organic layer was then dried (MgSO₄), filtered and solvent removed *in vacuo* to give a solid. This was triturated with ether and the solid filtered off and dried to give (2*R*)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (842 mg, 53 %).

¹H NMR Spectrum: (DMSO-d₆) 2.20 (s, 3H), 2.80 (m, 1H), 2.90 (m, 1H), 3.42 (m, 1H), 3.98 (s, 3H), 4.02 (m, 1H), 4.60 (m, 1H), 6.98 (dd, 1H), 7.18 (s, 1H) 7.35 (m, 3H), 7.60 (s, 1H), 8.45 (s, 1H) and 10.82 (s, 1H)

MS (ESI) : 378 (MH)⁺

Example 279

A mixture of (2*R*)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.65 mmol), (prepared as described in Example 278), and piperidine (0.2 ml, 2.04 mmol) in DMF (5 ml) was stirred at 60 °C for 24 hours and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography, gradient elution (dichloromethane, 5% methanol/95% dichloromethane, 1% 0.880 saturated aqueous ammonia/10% methanol/89% dichloromethane) (2*R*)-7-(2-hydroxy-3-piperidinopropoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline (237 mg, 78 %).

¹H NMR Spectrum: (DMSO-d₆) 1.38 (m, 2H), 1.50 (m, 4H), 2.34 (m, 9H), 3.98 (s, 3H), 4.16 (m, 3H), 4.85 (br s, 1H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.42 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 464 (MH)⁺

Elemental analysis:	Found	C 66.3	H 6.6	N 12.1
C ₂₆ H ₃₀ N ₄ O ₄ 0.5methanol	Requires	C 66.5	H 6.7	N 11.7%

Example 280

A mixture of (2*R*)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.65 mmol), (prepared as described in Example 278), and pyrrolidine (0.17 ml, 2.04 mmol) in DMF (5 ml) was stirred at 60 °C for 24 hours and allowed

to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) to give **(2R)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (257 mg, 88 %).

¹H NMR Spectrum: (DMSO-d₆) 1.65 (m, 4H), 1.98 (m, 2H), 2.20 (s, 3H), 2.50 (m, 2H), 2.62 (m, 2H), 3.98 (s, 3H), 4.17 (m, 3H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.42 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 449 (MH)⁺

10	Elemental analysis:	Found	C 64.1	H 6.4	N 12.6
	C ₂₅ H ₂₈ N ₄ O ₄ 1.0H ₂ O	Requires	C 64.4	H 6.5	N 12.0%

Example 281

A mixture of **(2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline** (350 mg, 0.93 mmol), (prepared as described in Example 278), and 1-methylpiperazine (0.31 ml, 2.78 mmol) in DMF (5 ml) was stirred at 60 °C for 24 hours and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) to give **(2R)-7-(2-hydroxy-3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (352 mg, 80 %).

¹H NMR Spectrum: (DMSO-d₆) 2.10 (s, 3H), 2.20 (s, 3H), 2.40 (m, 10H), 3.98 (s, 3H), 4.13 (m, 3H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.42 (s, 1H) and 10.82 (br s, 1H)

25 MS (ESI) : 478 (MH)⁺

	Elemental analysis:	Found	C 61.6	H 6.4	N 14.4
	C ₂₆ H ₃₁ N ₅ O ₄ 1.0H ₂ O.0.25Methanol	Requires	C 61.6	H 6.8	N 13.9%

Example 282

30 A mixture of **(2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline** (350 mg, 0.93 mmol), (prepared as described in Example 278), and morpholine (0.24 ml, 2.78 mmol) in DMF (5 ml) was stirred at 60 °C for 24 hours and

allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) to give **(2R)-7-(2-hydroxy-3-morpholinopropoxy)-6-methoxy-4-(3-**

5 **methylindol-5-yloxy)quinazoline** (398 mg, 93 %).

¹H NMR Spectrum: (DMSO-d₆) 2.20 (s, 3H), 2.44 (m, 6H), 3.48 (t, 4H), 3.98 (s, 3H), 4.13 (m, 3H), 4.98 (br s, 1H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.42 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 465 (MH)⁺

10	Elemental analysis:	Found	C 58.5	H 6.0	N 11.2
	C ₂₅ H ₂₈ N ₄ O ₅ · 2.5H ₂ O.	Requires	C 58.9	H 6.5	N 11.0%

Example 283

A mixture of **(2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-**
15 **ylmethoxy)quinazoline** (350 mg, 0.93 mmol), (prepared as described in Example 278), and 2.0 M dimethylamine in ethanol (4.60 ml, 9.30 mmol) in DMF (5 ml) was stirred at 60 °C for 24 hours and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography, gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous
20 ammonia (100/8/1)) to give **(2R)-7-(2-hydroxy-3-dimethylaminopropoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (308 mg, 78 %).

¹H NMR Spectrum: (DMSO-d₆) 2.10 (m, 9H), 2.20 (m, 2H), 3.98 (s, 3H), 4.13 (m, 3H), 4.98 (br s, 1H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.42 (s, 1H) and 10.82 (br s, 1H)

25 MS (ESI) : 423 (MH)⁺

Elemental analysis:	Found	C 65.5	H 6.2	N 13.2
C ₂₃ H ₂₀ N ₄ O ₄	Requires	C 65.4	H 6.2	N 13.3%

Example 284

30 A mixture of **(2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline** (350 mg, 0.93 mmol), (prepared as described in Example 278), and diethylamine (0.29 ml, 2.78 mmol) in DMF (5 ml) was stirred at 60 °C for 24 hours and

allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) to give **(2R)-7-(2-hydroxy-3-((N,N-diethylamino)propoxy))-6-**

5 **methoxy-4-(3-methylindol-5-yloxy)quinazoline** (338 mg, 81 %).

¹H NMR Spectrum: (DMSO-d₆) 0.95 (t, 6H), 2.11 (s, 3H), 2.40 (m, 6H), 3.98 (s, 3H), 4.13 (m, 3H), 4.84 (br s, 1H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.42 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 451 (MH)⁺

10	Elemental analysis:	Found	C 64.4	H 6.6	N 12.0
	C ₂₅ H ₃₀ N ₄ O ₄ 1.0H ₂ O.	Requires	C 64.1	H 6.9	N 12.0%

Example 285

A mixture of **(2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-**
15 **ylmethoxy)quinazoline** (350 mg, 0.93 mmol), (prepared as described in Example 278), and isopropylamine (0.29 ml, 4.65 mmol) in DMF (5 ml) was stirred at 100 °C for 24 hours and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous
20 ammonia (100/8/1)) to give **(2R)-7-(2-hydroxy-3-(isopropylamino)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (307 mg, 75 %).

¹H NMR Spectrum: (DMSO-d₆) 0.98 (d, 6H), 2.20 (s, 3H), 2.55-2.80 (m, 3H), 3.98 (s, 3H), 4.02-4.20 (m, 3H), 4.98 (br s, 1H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.30-7.40 (m, 3H), 7.60 (s, 1H), 8.42 (s, 1H) and 10.82 (br s, 1H)

25 MS (ESI) : 437 (MH)⁺

	Elemental analysis:	Found	C 63.3	H 6.3	N 12.4
	C ₂₄ H ₂₈ N ₄ O ₄ 1.0H ₂ O.	Requires	C 63.4	H 6.7	N 12.3%

Example 286

30 A mixture of **(2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline** (350 mg, 0.93 mmol), (prepared as described in Example 278), and diisopropylamine (0.78 ml, 5.58 mmol) in DMF (10 ml) was stirred at 130 °C for 24 hours

and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by silica column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, methanol/dichloromethane/0.880 saturated aqueous ammonia (100/8/1)) to give **(2R)-7-(2-hydroxy-3-((N,N-diisopropyl)amino)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (398 mg, 93 %).

¹H NMR Spectrum: (DMSO-d₆) 0.98 (d, 12H), 2.20 (s, 3H), 2.72 (m, 2H), 3.00 (m, 2H), 3.98 (s, 3H), 4.11 (m, 3H), 6.98 (dd, 1H), 7.18 (s, 1H), 7.35 (m, 3H), 7.60 (s, 1H), 8.42 (s, 1H) and 10.82 (br s, 1H)

MS (ESI) : 479 (MH)⁺

10	Elemental analysis:	Found	C 65.4	H 6.8	N 11.3
	C ₂₇ H ₃₄ N ₄ O ₄ 0.8H ₂ O.	Requires	C 55.8	H 7.2	N 11.4%

Example 287

A mixture of (2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (100 mg, 0.28 mmol), (prepared as described in Example 278), and 4-(3-aminopropyl)morpholine (0.12 ml, 0.84 mmol) in DMF (5 ml) was heated to 70 °C for 3 hours. The solvents were removed *in vacuo* and the residue taken up in dichloromethane. This was washed with water, dried (MgSO₃), filtered and evaporated. The residue was purified by silica column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, 20% methanolic ammonia (7M)/80% dichloromethane) to give **(2R)-7-(2-hydroxy-3-(3-morpholinopropylamino)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (67 mg, 46 %).

¹H NMR Spectrum: (DMSO-d₆) 1.28 (m, 2H), 2.30 (t, 4H), 2.56 (t, 2H), 2.650 (m, 4H), 3.55 (t, 4H), 3.98 (s, 3H), 4.15 (m, 3H), 6.42 (s, 1H), 6.98 (dd, 1H), 7.42 (m, 4H), 7.60 (s, 1H), 8.45 (s, 1H), and 11.19 (br s, 1H)

MS (ESI) : 508 (MH)⁺

Elemental analysis:	Found	C 59.7	H 6.6	N 13.4
C ₂₇ H ₃₃ N ₅ O ₅ 1.8H ₂ O	Requires	C 60.1	H 6.8	N 13.0%

Example 288

A mixture of (2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (100 mg, 0.28 mmol), (prepared as described in Example 278), and 1-

(3-aminopropyl)-4-methylpiperazine (132 mg, 0.84 mmol) in DMF (5 ml) was heated to 70 °C for 3 hours. The solvents were removed *in vacuo* and the residue taken up in dichloromethane. This was washed with water, dried (MgSO₄), filtered and evaporated. The residue was purified by silica column chromatography using gradient elution

5 (dichloromethane, 5% methanol/95% dichloromethane, 20% methanolic ammonia (7M)/80% dichloromethane) to give **(2R)-7-(2-hydroxy-3-(3-(4-methylpiperazin-1-yl)propylamino)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (44 mg, 31 %).

¹H NMR Spectrum: (DMSO-d₆) 1.55 (m, 2H), 2.10 (s, 3H), 2.30 (t, 8H), 2.62 (m, 6H), 3.98 (s, 10 3H), 4.12 (m, 3H), 6.42 (s, 1H), 6.98 (dd, 1H), 7.42 (m, 4H), 7.60 (s, 1H), 8.45 (s, 1H), and 11.19 (br s, 1H)

MS (ESI) : 521 (MH)⁺

Elemental analysis: Found C 61.3 H 7.3 N 16.1

C₂₈H₃₆N₆O₄ 1.6H₂O Requires C 61.2 H 7.2 N 16.3%

15

Example 289

A mixture of (2R)-6-methoxy-4-(3-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (70 mg, 0.19 mmol), (prepared as described in Example 278), and 1-(3-aminopropyl)pyrrolidine (74 mg, 0.58 mmol) in DMF (5 ml) was heated to 60 °C 20 overnight. The solvents were removed *in vacuo* and the residue purified by column chromatography using gradient elution (dichloromethane, 5% methanol/95% dichloromethane, 20% methanolic ammonia (7M)/80% dichloromethane) to give **(2R)-7-(2-hydroxy-3-(3-(pyrrolidin-1-yl)propylamino)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline** (64 mg, 68 %).

25 ¹H NMR Spectrum: (DMSO-d₆) 1.60 (m, 6H), 2.25 (m, 4H), 2.60 (m, 4H), 3.08 (m, 2H), 3.98 (s, 3H), 4.12 (m, 3H), 6.42 (s, 1H), 6.98 (dd, 1H), 7.34 (m, 4H), 7.58 (s, 1H), 8.42 (s, 1H), and 11.80 (br s, 1H)

MS (ESI) : 492 (MH)⁺

30 Example 290

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (380 mg, 1.13 mmol), (prepared as described for the starting material in Example 67), potassium carbonate

(469 mg, 3.4 mmol), 4-bromo-5-hydroxyindole (240 mg, 1.13 mmol) and DMA (4.0 ml) were stirred at 90°C for 3 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated under vacuum. The residue was purified by column chromatography eluting with dichloromethane/methanolic ammonia (7M) (95/5) to give an oil. This oil was further purified by column chromatography eluting with dichloromethane/methanol (60/40) to give **4-(4-bromoindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline** (256 mg, 44%).

¹H NMR Spectrum: (CDCl₃) 1.47 (m, 2H), 1.60 (m, 4H); 2.14 (m, 2H), 2.44 (m, 4H), 2.54 (t, 2H), 4.08 (s, 3H), 4.27 (t, 2H), 6.67 (m, 1H), 7.15 (d, 1H), 7.32 (t, 1H), 7.36 (s, 1H), 7.42 (d, 1H), 7.69 (s, 1H) 8.55 (br s, 1H) and 8.62 (s, 1H)

MS (ESI) : 511, 513 (MH)⁺

Elemental analysis	Found	C 58.2	H 5.3	N 10.8
C ₂₅ H ₂₇ BrN ₄ O ₃ 0.25 H ₂ O,	Requires	C 58.2	H 5.4	N 10.9%

The starting material was prepared as follows:

Ethyl 4-bromo-5-hydroxyindole-2-carboxylate (1.49 g, 5 mmol.), (Jnl. Org. Chem. 1984, 49, 4761), was dissolved in ethanol (10 ml) and water (3.5 ml). Potassium hydroxide (840 mg) was added and the mixture stirred at 50 °C under an atmosphere of nitrogen for 1 hour then cooled to ambient temperature. The solvent was evaporated and the residue redissolved in water (25 ml). 2M Aqueous hydrochloric acid was added until the reaction mixture was at pH4, giving a precipitate which was filtered off, washed with water and dried under vacuum to give 4-bromo-5-methoxyindole-2-carboxylic acid (1.30, 96%).

¹H NMR Spectrum: (DMSO-d₆) 3.83 (s, 3H), 6.90 (d, 1H), 7.16 (d, 1H), 7.40 (d, 1H). 11.88 (br s, 1H) and 13.19 (br s, 1H)

MS (ESI) : 268, 270 (M-H)

4-Bromo-5-methoxyindole-2-carboxylic acid (1.25 g, 4.19 mmol), quinoline (15ml) and copper chromite (313 mg) were mixed together. Nitrogen was gently bubbled through the mixture for 5 minutes, then the mixture heated quickly to 245°C under an atmosphere of nitrogen. After 90 minutes the mixture was cooled to ambient temperature diluted with ethyl acetate (100ml) and washed with 2M aqueous hydrochloric acid (60ml). The ethyl acetate layer was filtered, the filtrate dried (MgSO₄) and the solvent evaporated. The residue was

purified by silica column chromatography eluting with dichloromethane/hexane (1/1) to give 4-bromo-5-methoxyindole (635mg, 60%).

¹H NMR Spectrum (CDCl₃) 3.94 (s, 3H), 6.55 (m, 1H), 6.93 (d, 1H), 7.27 (m, 2H). 8.18 (br s, 1H)

5 MS (ESI) : 224, 226 (M-H)⁻

A solution of 4-bromo-5-methoxyindole (540 mg, 2.4 mmol) in dichloromethane (12ml) was cooled to - 40°C under an atmosphere of nitrogen. Boron tribromide (4.8 ml of a 1M solution in dichloromethane, 4.8 mmol) was added dropwise then the mixture warmed to ambient temperature and stirred for 1 hour. The mixture was diluted with dichloromethane
10 (5ml) and washed with 2M aqueous hydrochloric acid (3ml). The organic layer was separated, dried (MgSO₄) and evaporated to give a dark oil. This was purified by silica column chromatography eluting with dichloromethane/ethyl acetate (8/2) to give 4-bromo-5-hydroxyindole (295 mg, 55%).

¹H NMR Spectrum: (CDCl₃) 6.46 (m, 1H), 7.92 (d, 1H), 7.22 (m, 2H), 8.80 (br s, 1H)

15 MS (ESI) : 210, 212 (M-H)⁻

Example 291

Nitrogen was bubbled through a mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (335 mg, 0.68 mmol), (prepared as described for the starting
20 material in Example 67), potassium carbonate (281.5 mg, 2.04 mmol), 5-hydroxy-1-methylindole (100 mg, 0.68 mmol) and DMA (4.0 ml) for 5 minutes. The mixture was then stirred at 90°C for 4 hours under an atmosphere of nitrogen and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated under vacuum. The residue was purified by trituration with methanol then water to give **6-methoxy-4-(1-methylindol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline** (148 mg, 49%).

¹H NMR Spectrum: (DMSO-d₆) 1.38 (m, 2H), 1.51 (m, 4H), 1.93 (m, 2H), 2.35 (m, 4H), 2.41 (t, 2H), 3.83 (s, 3H), 3.97 (s, 3H), 4.24 (t, 2H), 6.42 (d, 1H), 7.06 (dd, 1H), 7.33 (s, 1H), 7.42 (m, 2H), 7.50 (d, 1H), 7.59 (s, 1H) and 8.47 (s, 1H)

MS (ESI) : 447 (MH)⁺

30 Elemental analysis

Found	C 69.5	H 6.8	N 12.5
Requires	C 69.9	H 6.8	N 12.6%

C₂₆H₃₀N₄O₃

The starting material was prepared as follows:

A solution of 5-benzyloxy-1-methylindole (3.5 g, 15.7 mmol), in ethanol (100 ml) was hydrogenated at ambient temperature and 1 atmosphere pressure hydrogen for 4 hours using 10% palladium on carbon (0.5 g) as catalyst. The catalyst was filtered off and the filtrate evaporated *in vacuo*. The residue was purified by silica column chromatography eluting with ethyl acetate/dichloromethane (10/90) to give 5-hydroxy-1-methylindole (2.1 g, 97%).

MS (ESI) : 146 (M-H)⁻

¹H NMR Spectrum: (CDCl₃) 3.74 (s, 3H), 4.50 (s, 1H), 6.33 (d, 1H), 6.79 (dd, 1H), 7.00 (m, 2H), 7.17 (d, 1H)

Example 292

A mixture of (2*R*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.83 mmol), and pyrrolidine (176 mg, 2.48 mmol) in DMF (5 ml) was stirred at 75°C for 3 hours under an atmosphere of nitrogen and allowed to cool to ambient temperature.

The solvents were removed *in vacuo* and the residue purified on silica gel, gradient elution with dichloromethane, dichloromethane/methanol (95/5), dichloromethane/methanolic ammonia (7M) (98/2 to 90/10), to give (2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (326 mg, 87%).

¹H NMR Spectrum: (CDCl₃) 1.80 (m, 4H), 2.56 (m, 3H), 2.71 (m, 2H), 2.87 (m, 1H), 4.04 (s, 3H), 4.23 (m, 3H), 6.59 (m, 1H), 7.07 (dd, 1H), 7.25 (m, 1H), 7.32 (s, 1H), 7.45 (d, 1H), 7.50 (d, 1H), 7.61 (s, 1H), 8.30 (br s, 1H) and 8.60 (s, 1H)

MS (ESI) : 435(MH)⁺

Elemental analysis

Found C 63.4 H 5.9 N 12.3

C₂₄H₂₆N₄O₄·1H₂O

Requires C 63.7 H 6.2 N 12.4%

The starting material was prepared as follows:

Nitrogen was bubbled through a mixture of 7-hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (3.07 g, 10 mmol), (prepared as described for the starting material in Example 107), potassium carbonate (4.14 g, 30 mmol) and (2*R*)-(-)-glycidyl tosylate (4.57 g, 20 mmol) in DMA (35 ml) for 5 minutes. The mixture was then stirred at 60 °C for 2 hours under an atmosphere of nitrogen and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by column

chromatography on silica by gradient elution with dichloromethane/methanol (100/0 to 95/5), to give (2*R*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline as a yellow solid (1.92g, 53%).

¹H NMR Spectrum: (DMSO-d₆) 2.75 (m, 1H), 2.89 (m, 1H), 3.44 (m, 1H), 3.97 (s, 3H), 4.06 (m, 1H), 4.58 (dd, 1H), 6.44 (m, 1H), 6.95 (dd, 1H), 7.40 (m, 4H) 7.62 (s, 1H), 8.47 (s, 1H), 11.19 (br s 1H)

MS (ESI) : 364 (MH)⁺

Example 293

Using an analogous procedure to that described in Example 292, (2*R*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.83 mmol), (prepared as described for the starting material in Example 292), was reacted with morpholine (211 mg, 2.49 mmol) to give (2*R*)-7-(2-hydroxy-3-morpholinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (338 mg, 85%).

¹H NMR Spectrum: (CDCl₃) 2.48 (m, 2H), 2.624 (m, 2H), 2.68 (m, 2H), 3.78 (m, 4H), 4.04 (s, 3H), 4.24 (m, 3H), 6.58 (m, 1H), 7.08 (dd, 1H), 7.29 (m, 1H), 7.34 (s, 1H), 7.46 (d, 1H), 7.50 (d, 1H), 7.62 (s, 1H), 8.31 (br s, 1H) and 8.62 (s, 1H)

MS (ESI) : 451(MH)⁺

Elemental analysis	Found	C	60.3	H	5.9	N	12.3
--------------------	-------	---	------	---	-----	---	------

C ₂₄ H ₂₆ N ₄ O ₅ ·1.5H ₂ O	Requires	C	60.4	H	6.1	N	11.7%
--	----------	---	------	---	-----	---	-------

Example 294

Using an analogous procedure to that described in Example 292, (2*R*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.83 mmol), (prepared as described for the starting material in Example 292), was reacted with piperidine (211 mg, 2.49 mmol) to give (2*R*)-7-(2-hydroxy-3-piperidinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (325 mg, 86%).

¹H NMR Spectrum: (CDCl₃) 1.47 (m, 2H), 1.61 (m, 4H), 2.39 (m, 2H), 2.54 (d, 2H), 2.64 (m, 2H), 4.04 (s, 3H), 4.24 (m, 3H), 6.58 (m, 1H), 7.08 (dd, 1H), 7.29 (m, 1H), 7.32 (s, 1H), 7.45 (d, 1H), 7.48 (d, 1H), 7.62 (s, 1H), 8.28 (br s, 1H) and 8.60 (s, 1H)

MS (ESI) : 449 (MH)⁺

Elemental analysis	Found	C	65.9	H	6.3	N	12.3
--------------------	-------	---	------	---	-----	---	------

$C_{25}H_{28}N_4O_4 \cdot 0.5H_2O$

Requires C 65.6 H 6.4 N 12.3%

Example 295

A mixture of (2*R*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.83 mmol), (prepared as described for the starting material in Example 292), and dimethylamine (1.24 ml of a 2M solution in THF, 2.48 mmol) in DMF (5 ml) was stirred at 75°C for 3 hours under an atmosphere of nitrogen then allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by trituration with methanol to give (2*R*)-7-(2-hydroxy-3-dimethylaminopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (265 mg, 63%).

¹H NMR Spectrum: (DMSO-d₆) 2.21 (s, 6H), 2.38 (m, 2H), 3.97 (s, 3H), 4.073 (m, 2H), 4.21 (m, 1H), 4.96 (d, 1H), 6.43 (m, 1H), 6.97 (dd, 1H), 7.37 (s, 1H), 7.43 (m, 3H), 7.62 (s, 1H), 8.48 (s, 1H) and 11.20 (br s, 1H)

MS (ESI) : 409(MH)⁺

Elemental analysis	Found	C 62.8	H 5.8	N 13.2
--------------------	-------	--------	-------	--------

$C_{22}H_{24}N_4O_4 \cdot 0.7H_2O$	Requires	C 62.8	H 6.1	N 13.3%
------------------------------------	----------	--------	-------	---------

Example 296

A mixture of (2*R*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (300 mg, 0.83 mmol), (prepared as described for the starting material in Example 292), and diisopropylamine (1.35 ml, 9.7 mmol) in DMF (5 ml) was stirred at 70°C for 19 hours under an atmosphere of nitrogen then allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified on silica gel using gradient elution with dichloromethane, dichloromethane/methanol (95/5), dichloromethane/methanolic ammonia (7M) (98/2 to 90/10) to give (2*R*)-7-(2-hydroxy-3-((*N,N*-diisopropyl)amino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (343 mg, 86%).

¹H NMR Spectrum: (CDCl₃) 1.08 (m, 12H), 1.57 (m, 1H), 1.75 (m, 1H), 3.10 (m, 2H), 4.04 (s, 3H), 4.16 (m, 3H), 6.58 (m, 1H), 7.08 (dd, 1H), 7.26 (m, 1H), 7.32 (s, 1H), 7.45 (d, 1H), 7.50 (d, 1H), 7.61 (s, 1H), 8.32 (br s, 1H) and 8.61 (s, 1H)

MS (ESI) : 465(MH)⁺

Elemental analysis	Found	C 64.8	H 6.8	N 11.9
--------------------	-------	--------	-------	--------

$C_{26}H_{32}N_4O_4 \cdot 1.0H_2O$	Requires	C 64.6	H 7.0	N 11.6%
------------------------------------	----------	--------	-------	---------

Example 297

A mixture of (2*S*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (100 mg, 0.28 mmol), and pyrrolidine (60 mg, 0.84 mmol) in DMF (5 ml) was stirred at 75°C for 3 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified on silica gel, gradient elution with dichloromethane, dichloromethane/methanol (95/5), dichloromethane/methanolic ammonia (7M) (98/2 to 90/10), to give (2*S*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (114 mg, 92%).

¹H NMR Spectrum: (CDCl₃) 1.80 (m, 4H), 2.56 (m, 3H), 2.71 (m, 2H), 2.86 (m, 1H), 4.04 (s, 3H), 4.23 (m, 3H), 6.59 (m, 1H), 7.07 (dd, 1H), 7.25 (m, 1H), 7.32 (s, 1H), 7.45 (d, 1H), 7.50 (d, 1H), 7.61 (s, 1H), 8.30 (br s, 1H) and 8.60 (s, 1H)

MS (ESI) : 435(MH)⁺

Elemental analysis

Found C 64.7 H 6.0 N 12.6

C₂₄H₂₆N₄O₄·0.5H₂O

Requires C 64.9 H 6.1 N 12.7%

The starting material was prepared as follows:

Nitrogen was bubbled through a mixture of 7-hydroxy-4-(indol-5-yloxy)-6-methoxyquinazoline (3.07 g, 10 mmol), (prepared as described for the starting material in Example 107), potassium carbonate (4.14 g, 30 mmol) and (2*S*)-(+)-glycidyl tosylate (4.57 g, 20 mmol) in DMA (35 ml) for 5 minutes. This mixture was then stirred at 60 °C for 2 hours under an atmosphere of nitrogen and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by column chromatography on silica by gradient elution with dichloromethane/methanol (100/0 to 95/5), to give after removal of the solvents *in vacuo* and trituration of the residue with ether, (2*S*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (1.88g, 52%) as a yellow solid.

¹H NMR Spectrum: (DMSO-d₆) 2.75 (m, 1H), 2.89 (m, 1H), 3.44 (m, 1H), 3.97 (s, 3H), 4.06 (m, 1H), 4.58 (dd, 1H), 6.44 (m, 1H), 6.95 (dd, 1H), 7.46 (m, 4H) 7.62 (s, 1H), 8.47 (s, 1H) and 11.19 (br s 1H)

MS (ESI) : 364 (MH)⁺

Example 298

Using an analogous procedure to that described in Example 297, (2*S*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (100 mg, 0.28 mmol), (prepared as described for the starting material in Example 297), was reacted with morpholine (73.2 mg, 0.84 mmol) to give (2*S*)-7-(2-hydroxy-3-morpholinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (82 mg, 63%).

¹H NMR Spectrum: (CDCl₃) 2.48 (m, 2H), 2.62 (m, 2H), 2.68 (m, 2H), 3.78 (m, 4H), 4.04 (s, 3H), 4.29 (m, 3H), 6.58 (m, 1H), 7.08 (dd, 1H), 7.29 (m, 1H), 7.34 (s, 1H), 7.46 (d, 1H), 7.50 (d, 1H), 7.62 (s, 1H), 8.31 (br s, 1H) and 8.62 (s, 1H)

MS (ESI) : 451(MH)⁺

Elemental analysis	Found	C 61.7	H 5.7	N 11.8
C ₂₄ H ₂₆ N ₄ O ₅ ·1.0H ₂ O	Requires	C 61.5	H 6.0	N 12.0%

Example 299

Using an analogous procedure to that described in Example 297, (2*S*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (100 mg, 0.28 mmol), (prepared as described for the starting material in Example 297), was reacted with piperidine (70 mg, 0.83 mmol), to give (2*S*)-7-(2-hydroxy-3-piperidinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (93 mg, 73%).

¹H NMR Spectrum: (CDCl₃) 1.47 (m, 2H), 1.61 (m, 4H), 2.39 (m, 2H), 2.54 (d, 2H), 2.64 (m, 2H), 4.04 (s, 3H), 4.29 (m, 3H), 6.58 (m, 1H), 7.08 (dd, 1H), 7.29 (m, 1H), 7.32 (s, 1H), 7.45 (d, 1H), 7.48 (d, 1H), 7.62 (s, 1H), 8.28 (br s, 1H) and 8.60 (s, 1H)

MS (ESI) : 449 (MH)⁺

Elemental analysis	Found	C 65.8	H 6.2	N 12.2
C ₂₅ H ₂₈ N ₄ O ₄ ·0.5H ₂ O	Requires	C 65.6	H 6.4	N 12.3%

Example 300

A mixture of (2*S*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (100 mg, 0.28 mmol), (prepared as described for the starting material in Example 297), and dimethylamine (0.42 ml of a 2M solution in THF, 0.84 mmol) in DMF (5 ml) was stirred at 75°C for 3 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified by trituration with

methanol to give **(2S)-7-(2-hydroxy-3-dimethylaminopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline** (100 mg, 85%).

¹H NMR Spectrum: (DMSO-d₆) 2.21 (s, 6H), 2.38 (m, 2H), 3.97 (s, 3H), 4.083 (m, 2H), 4.21 (m, 1H), 4.96 (d, 1H), 6.43 (m, 1H), 6.97 (dd, 1H), 7.37 (s, 1H), 7.43 (m, 3H), 7.62 (s, 1H), 8.48 (s, 1H) and 11.20 (br s, 1H)

MS (ESI) : 409(MH)⁺

Elemental analysis	Found	C 63.6 H 6.0 N 13.3
C ₂₂ H ₂₄ N ₄ O ₄ ·0.5H ₂ O	Requires	C 63.3 H 6.0 N 13.4%

10 **Example 301**

A mixture of **(2S)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline** (100 mg, 0.28 mmol), (prepared as described for the starting material in Example 297), and diisopropylamine (0.45 ml, 3.2 mmol) in DMF (5 ml) was stirred at 70°C for 19 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified on silica gel, using gradient elution with dichloromethane/methanol (100/0 to 95/5), dichloromethane/methanolic ammonia (7M) (98/2 to 90/10) to give **(2S)-7-(2-hydroxy-3-((N,N)-diisopropyl)amino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline** (43 mg, 33%).

¹H NMR Spectrum: (CDCl₃) 1.08 (m, 12H), 1.57 (m, 1H), 1.759 (m, 1H), 3.10 (m, 2H), 4.04 (s, 3H), 4.16 (m, 3H), 6.58 (m, 1H), 7.08 (dd, 1H), 7.26 (m, 1H), 7.32 (s, 1H), 7.45 (d, 1H), 7.50 (d, 1H), 7.61 (s, 1H), 8.32 (br s, 1H) and 8.61 (s, 1H)

MS (ESI) : 465(MH)⁺

Elemental analysis	Found	C 67.2 H 7.0 N 11.9
C ₂₆ H ₃₂ N ₄ O ₄	Requires	C 67.2 H 6.9 N 12.1%

25 **Example 302**

A mixture of **(2R)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline** (100 mg, 0.28 mmol), (prepared as described for the starting material in Example 292), and isopropylamine (1.0 ml) in THF (10 ml) was stirred at 75°C for 18 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by silica gel chromatography, gradient elution with dichloromethane/methanolic ammonia (7M) (100/0 to 90/10) to give **(2R)-7-(2-**

hydroxy-3-(isopropylamino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (82 mg, 68%).

¹H NMR Spectrum: (DMSO-d₆) 0.98 (m, 6H), 2.68 (m, 3H), 3.96 (m, 4H), 4.13 (m, 2H), 5.06 (br s, 1H), 6.44 (s, 1H), 6.98 (dd, 1H), 7.439 (m, 4H), 7.60 (s, 1H), 8.46 (s, 1H) and 11.22 (s, 1H)

MS (ESI) : 423(MH)⁺

Elemental analysis	Found	C 63.6	H 6.4	N 12.9
C ₂₃ H ₂₆ N ₄ O ₄ ·0.6H ₂ O	Requires	C 63.8	H 6.3	N 12.9%

10 **Example 303**

A mixture of (2*S*)-4-(indol-5-yloxy)-6-methoxy-7-(oxiran-2-ylmethoxy)quinazoline (100 mg, 0.28 mmol), (prepared as described for the starting material in Example 297), and isopropylamine (1.0 ml) in THF (10 ml) was stirred at 75°C for 18 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by silica gel chromatography using gradient elution with dichloromethane/methanolic ammonia (7M) (100/0 to 90/10) to give (2*S*)-7-(2-hydroxy-3-(isopropylamino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline (66 mg, 56%).

¹H NMR Spectrum: (DMSO-d₆) 0.985 (m, 6H), 2.68 (m, 3H), 3.96 (m, 4H), 4.13 (m, 2H), 5.06 (br s, 1H), 6.44 (s, 1H), 6.98 (dd, 1H), 7.43 (m, 4H), 7.60 (s, 1H), 8.46 (s, 1H) and 11.22 (s, 1H)

MS (ESI) : 423(MH)⁺

Elemental analysis	Found	C 63.1	H 6.3	N 12.7
C ₂₃ H ₂₆ N ₄ O ₄ ·0.9 H ₂ O	Requires	C 63.0	H 6.4	N 12.8%

25 **Example 304**

A mixture of (2*S*)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (250 mg, 0.66 mmol), and pyrrolidine (1.5 ml) in THF (10 ml) was stirred at 75°C for 3 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by silica gel chromatography using gradient elution with dichloromethane/methanolic ammonia (7M) (100/0 to 90/10) to give (2*S*)-7-(2-hydroxy-3-

(pyrrolidin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (106 mg, 36%).

¹H NMR Spectrum: (DMSO-d₆) 1.60 (s, 4H), 2.38 (s, 3H), 2.57 (m, 6H), 4.11 (m, 6H), 4.95 (d, 1H), 6.14 (s, 1H), 6.88 (dd, 1H), 7.29 (m, 2H), 7.37 (s, 1H), 7.59 (s, 1H), 8.48 (s, 1H) and 11.00 (s, 1H)

MS (ESI) : 450 (MH)⁺

Elemental analysis	Found	C 67.0	H 6.5	N 12.0
C ₂₅ H ₂₈ N ₄ O ₄ ·0.1 H ₂ O	Requires	C 66.7	H 6.3	N 12.4%

The starting material was prepared as follows:

A mixture of 7-hydroxy-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline (300 mg, 0.93 mmol), (prepared as described in Example 49), potassium carbonate (385 mg, 2.79 mmol) and (2S)-(-)- glycidyl tosylate (426 mg, 2.79 mmol) in DMF (15 ml) was stirred at 60°C for 2 hours and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*. The residue was dissolved in dichloromethane and washed with saturated sodium hydrogen carbonate solution. The organic layer was then dried (MgSO₄), filtered and the solvent removed *in vacuo* to give a yellow solid. This was triturated with ether, filtered off and dried to give (2S)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline as a yellow solid (277 mg, 78 %).

¹H NMR Spectrum: (DMSO) 2.40 (s, 3H), 2.75 (m, 1H), 2.90 (m, 1H), 3.40 (m, 1H), 3.98 (s, 3H), 4.05 (m, 1H), 4.60 (m, 1H), 6.15 (s, 1H), 6.85 (dd, 1H), 7.30 (m, 2H) 7.40 (s, 1H), 7.60 (s, 1H), 8.45 (s, 1H) and 10.98 (s, 1H)

MS (ESI) : 378 (MH)⁺

Example 305

A mixture of the (2R)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (250 mg, 0.66 mmol), (prepared as described for the starting material in Example 269), and pyrrolidine (1.5 ml) in THF (10 ml) was stirred at 75°C for 3 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by silica gel chromatography using gradient elution with dichloromethane/methanolic ammonia (7M)

(100/0 to 90/10) to give **(2R)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (165 mg, 55%).

¹H NMR Spectrum: (DMSO-d₆) 1.60 (s, 4H), 2.38 (s, 3H), 2.57 (m, 6H), 4.11 (m, 6H), 4.95 (d, 1H), 6.14 (s, 1H), 6.88 (dd, 1H), 7.29 (m, 2H), 7.37 (s, 1H), 7.59 (s, 1H), 8.48 (s, 1H) and 11.00 (s, 1H)

MS (ESI) : 450 (MH)⁺

Elemental analysis	Found	C 66.8	H 6.3	N 12.4
C ₂₅ H ₂₈ N ₄ O ₄ ·0.1 H ₂ O	Requires	C 66.7	H 6.3	N 12.4%

10 Example 306

A mixture of (2S)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (250 mg, 0.66 m.mol), (prepared as described for the starting material in Example 304), and isopropylamine (1.5 ml) in THF (10 ml) was stirred at 75°C for 18 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by silica gel chromatography using gradient elution with dichloromethane/methanolic ammonia (7M) (100/0 to 90/10) to give **(2S)-7-(2-hydroxy-3-(isopropylamino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (210 mg, 73%).

¹H NMR Spectrum: (DMSO-d₆) 0.99 (d, 6H), 2.39 (s, 3H), 2.66 (m, 3H), 4.07 (m, 6H), 5.08 (d, 1H), 6.14 (s, 1H), 6.88 (dd, 1H), 7.29 (m, 2H), 7.37 (s, 1H), 7.58 (s, 1H), 8.49 (s, 1H) and 11.03 (s, 1H)

MS (ESI) : 437 (MH)⁺

Elemental analysis	Found	C 64.3	H 6.4	N 12.3
C ₂₄ H ₂₈ N ₄ O ₄ ·0.5 H ₂ O	Requires	C 64.7	H 6.6	N 12.6%

Example 307

A mixture of (2R)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (250 mg, 0.66 mmol), (prepared as described for the starting material in Example 269), and isopropylamine (1.5 ml) in THF (10 ml) was stirred at 75°C for 18 hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by silica gel chromatography using gradient elution with dichloromethane/methanolic ammonia (7M)

(100/0 to 90/10) to give **(2R)-7-(2-hydroxy-3-(isopropylamino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline** (243 mg, 84%).

¹H NMR Spectrum: (DMSO-d₆): 0.99 (d, 6H), 2.39 (s, 3H), 2.66 (m, 3H), 4.07 (m, 6H), 5.08 (d, 1H), 6.14 (s, 1H), 6.88 (dd, 1H), 7.29 (m, 2H), 7.37 (s, 1H), 7.58 (s, 1H), 8.49 (s, 1H) and 11.03 (s, 1H)

MS (ESI) : 437 (MH)⁺

Elemental analysis	Found	C 64.3	H 6.5	N 12.3
C ₂₄ H ₂₈ N ₄ O ₄ .0.5 H ₂ O	Requires	C 64.7	H 6.6	N 12.6%

10 Example 308

Nitrogen was bubbled through a mixture of 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (400 mg, 1.19 mmol), (prepared as described for the starting material in Example 1), potassium carbonate (476 mg, 3.45 mmol), 5-hydroxy-1-methylindole (220 mg, 1.5 mmol), (prepared as described for the starting material in Example 291), and DMA (5.0 ml) for 5 minutes. The mixture was then stirred at 90°C for 3 hours under an atmosphere of nitrogen and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by trituration with methanol to give **6-methoxy-4-(1-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline** (312 mg, 59%).

¹H NMR Spectrum: (CDCl₃) 2.13 (m, 2H), 1.48 (t, 4H), 1.57 (t, 2H), 3.72 (t, 4H), 3.84 (s, 3H), 4.05 (s, 3H), 4.3 (t, 2H), 6.50 (d, 1H), 7.08-7.13 (m, 2H), 7.32 (s, 1H), 7.37 (s, 1H), 7.47 (d, 1H), 7.62 (s, 1H) 8.59 (s, 1H)

MS (ESI) : 449 (MH)⁺

Elemental analysis	Found	C 66.5	H 6.4	N 12.3
C ₂₅ H ₂₈ N ₄ O ₄ .0.1 H ₂ O	Requires	C 66.7	H 6.3	N 12.4%

Example 309

Nitrogen was bubbled through a mixture of 4-chloro-6-methoxy-7-(2-piperidinoethoxy)quinazoline (400 mg, 1.24 mmol), (prepared as described for the starting material in Example 180), potassium carbonate (500 mg, 3.62 mmol), 5-hydroxy-1-methylindole (231 mg, 1.57 mmol), (prepared as described for the starting material in Example 291), and DMA (5.0 ml) for 5 minutes. The mixture was then stirred at 90°C for 3

hours under an atmosphere of nitrogen and then allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by trituration with methanol to give **6-methoxy-4-(1-methylindol-5-yloxy)-7-(2-piperidinopropoxy)quinazoline** (447 mg, 83%).

5 ¹H NMR Spectrum: (CDCl₃) 1.47 (m, 2H), 1.64 (m, 4H), 2.57 (t, 4H) 2.94 (t, 2H), 3.83 (s, 3H), 4.05 (s, 3H), 4.34 (t, 2H), 6.49 (d, 1H), 7.10 (m, 2H), 7.32 (s, 1H), 7.38 (d, 1H), 7.45 (d, 1H), 7.62 (s, 1H) 8.60 (s, 1H)

MS (ESI) : 433 (MH)⁺

Elemental analysis

Found C 69.2 H 6.7 N 12.7

10 C₂₅H₂₈N₄O₃

Requires C 69.4 H 6.5 N 13.0%

Example 310

Nitrogen was bubbled through a mixture of 4-chloro-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline (400 mg, 1.24 mmol), (prepared as described for the starting material
15 in Example 9), potassium carbonate (500 mg, 3.62 mmol), 5-hydroxy-1-methylindole (231 mg, 1.57 mmol), (prepared as described for the starting material in Example 291), and DMA (5.0 ml) for 5 minutes. The mixture was then stirred at 90°C for 3 hours under an atmosphere of nitrogen and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*. The residue was purified by column chromatography,
20 gradient elution, with dichloromethane/methanolic ammonia (7M), (100/0 to 90/10) to give **6-methoxy-4-(1-methylindol-5-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline** (247 mg, 44%).

¹H NMR Spectrum: (CDCl₃) 1.81 (m, 4H), 2.18 (m, 2H), 2.56 (m, 4H), 2.69 (t, 2H), 3.82 (s, 3H), 4.05 (s, 3H), 4.30 (t, 2H), 6.45 (d, 1H), 7.09 (dd, 2H), 7.31 (s, 1H), 7.38 (d, 1H), 7.47 (d,
25 1H), 7.62 (s, 1H) and 8.59 (s, 1H)

MS (ESI) : 433 (MH)⁺

Elemental analysis

Found C 66.5 H 6.3 N 12.4

C₂₅H₂₈N₄O₃ 0.1 dichloromethane+0.7 H₂O

Requires C 66.7 H 6.6 N 12.4%

30 Example 311

Nitrogen was bubbled through a mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (114 mg, 0.34 mmol), (prepared as described for the starting

material in Example 67), potassium carbonate (141 mg, 1.02 mmol), 5-hydroxy-4-nitroindole (60.5 mg, 0.34 mmol) and DMA (8.5 ml) for 5 minutes at ambient temperature. This mixture was then stirred at 90°C for 4 hours under an atmosphere of nitrogen and allowed to cool to ambient temperature. The reaction mixture was filtered and the filtrate evaporated *in vacuo*.

5 The residue was purified by silica column chromatography using gradient elution with dichloromethane/methanol (100/0 to 95/5) followed by dichloromethane/methanolic ammonia (7M) (95/5) to give a partially purified oil. This oil was further purified by silica column chromatography, gradient elution with ethyl acetate/methanolic ammonia (95/5 to 80/20) to give **6-methoxy-(4-nitroindol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline** (63 mg, 39%).

10 ¹H NMR Spectrum: (CDCl₃) 1.46 (m, 2H), 1.60 (m, 4H), 2.16 (m, 2H), 2.43 (m, 4H), 2.54 (t, 2H), 3.85 (s, 3H), 4.33 (t, 2H), 7.04 (d, 1H), 7.10 (s, 1H), 7.47 (s, 1H), 7.57 (d, 1H), 7.83 (d, 1H), 7.95 (d, 1H) and 9.09 (s, 1H)

MS (ESI) : 478 (MH)⁺

Elemental analysis

Found

C 62.5 H 5.8 N 14.7

15 C₂₅H₂₇N₅O₅

Requires

C 62.9 H 5.7 N 14.7%

The starting material was prepared as follows:

A mixture of ethyl 5-methoxyindole-2-carboxylate (8.15 g, 37.2 mmol), (prepared by the method described in Heterocycles Vol. 43, No. 2, p. 263-266), nitric acid adsorbed on
20 silica gel (24g) and dichloromethane (150 ml) was stirred at ambient temperature for 18 hours. The dichloromethane was removed *in vacuo* and the product washed off the silica with acetone. The acetone was evaporated *in vacuo*. The residue was treated again with nitric acid on silica (1g) as above and the work up procedure repeated to give ethyl 5-hydroxy-4-nitroindole-2-carboxylate (5.8 g, 59%).

25 ¹H NMR Spectrum: (DMSO-d₆) 1.33 (t, 3H), 3.95 (s, 3H), 4.35 (q, 2H), 7.19 (d, 1H), 7.35 (d, 1H), 7.75 (d, 1H) and 12.45 (br s, 1H)

Ethyl 5-hydroxy-4-nitroindole-2-carboxylate (1.0 g, 3.8 mmol.) was suspended in a mixture of ethanol (20 ml) and water (5 ml). Potassium hydroxide (840 mg) was added and the mixture stirred at 50 °C under an atmosphere of nitrogen for 1 hour then cooled to ambient
30 temperature. The solvent was evaporated *in vacuo* and the residue re-dissolved in water (25 ml). The pH was adjusted to pH2 using aqueous hydrochloric acid (2M). The resulting

precipitate was filtered off, washed with water and dried *in vacuo* to give 5-methoxy-4-nitroindole-2-carboxylic acid (790 mg). This was used without further purification.

The crude 5-methoxy-4-nitroindole-2-carboxylic acid (720 mg, 3.05 mmol), quinoline (9 ml) and copper chromite (180 mg) were stirred together. Nitrogen was gently bubbled through the mixture for 5 minutes, then the mixture was heated quickly to 225°C, and stirred at this temperature for 40 minutes under an atmosphere of nitrogen. The mixture was cooled to ambient temperature diluted with ethyl acetate (80ml) and the insoluble material filtered off. The filtrate was extracted twice with aqueous hydrochloric acid (2M) and then with saturated aqueous sodium hydrogen carbonate solution. The ethyl acetate layer was dried (MgSO₄), evaporated and the residue purified by silica column chromatography eluting with dichloromethane to give 5-methoxy-4-nitroindole (129mg, 22%).

¹H NMR Spectrum: (CDCl₃) 3.99 (s, 3H), 6.88 (t, 1H), 6.97 (d, 1H), 7.37 (t, 1H), 7.55 (d, 1H) and 8.38 (br s, 1H)

MS (ESI) : 193 (MH)⁺

A solution of 5-methoxy-4-nitroindole (110 mg, 0.57 mmol) in dichloromethane (12 ml) was cooled to - 30°C under an atmosphere of nitrogen. Boron tribromide (0.74 ml of a 1M solution in dichloromethane, 0.74 mmol) was added dropwise then the mixture warmed to ambient temperature and stirred for 1 hour. The mixture was cooled to 5°C, diluted with dichloromethane (5ml), and water (10 ml). After stirring for 5 minutes the insoluble material was filtered off and the dichloromethane layer separated, dried (MgSO₄), and evaporated to give a dark oil which was and purified by silica column chromatography eluting with dichloromethane to give 5-hydroxy-4-nitroindole (68 mg, 67%).

¹H NMR Spectrum: (CDCl₃) 6.95 (d, 1H), 7.29 (m, 1H), 7.43 (t, 1H), 7.63 (d, 1H) and 11.60 (br s, 1H)

MS (ESI) : 177 (M-H)⁻

Example 312

6-Methoxy-(4-nitroindol-5-yl)-7-(3-piperidinopropoxy)quinazoline (45 mg 0.094 mmol), (prepared as described in Example 311), ethanol (20 ml) and 10% palladium on charcoal were hydrogenated at 45°C and 1 atmosphere pressure of hydrogen for 3.5 hours. The mixture was cooled to ambient temperature, the catalyst filtered off and the filtrate evaporated *in vacuo*. The residue was purified

by silica column chromatography using gradient elution with dichloromethane/methanolic ammonia (7M) (100/0 to 95/5), to give **4-(4-amino-indol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline** (39 mg, 87%).

¹H NMR Spectrum: (CDCl₃) 1.39(m, 2H), 1.50 (m, 4H), 1.96 (m, 2H), 2.35 (m, 4H), 2.43 (t, 2H), 3.80 (s, 3H), 4.28 (t, 2H), 4.84 (br s, 2H), 6.68 (d, 1H), 6.78 (d, 1H), 6.94 (s, 1H), 7.28 (s, 1H), 7.45 (s, 1H), 7.69 (s, 1H), 8.45 (br s, 1H) and 8.98 (s, 1H)

MS (ESI) : 448 (MH)⁺

Elemental analysis

Found C 64.0 H 6.4 N 14.4

C₂₅H₂₉N₅O₃·0.3 H₂O+ 0.4 dichloromethane Requires C 63.6 H 6.3 N 14.4%

Example 313

A mixture of 4-chloro-6-methoxy-7-(3-piperidinopropoxy)quinazoline (227 mg, 0.68 mmol), (prepared as described for the starting material in Example 67), 5-hydroxy-1*H*-pyrrolo[2,3-*b*]pyridine (100 mg, 0.75 mmol), (prepared as described for the starting material in Example 182), and potassium carbonate (350 mg, 2.5mmol) in DMF (4 ml) was stirred at 95°C for 6 hours and allowed to cool to ambient temperature. The reaction mixture was treated with 1.0 N aqueous sodium hydroxide solution and allowed to stir at ambient temperature for a few minutes. The resulting precipitate was filtered off, washed with water and air dried to give a crude product. This was purified by column chromatography, eluting initially with dichloromethane/methanol (85/15) to isolate a less polar impurity and then with dichloromethane/methanol/0.88 ammonia (100/8/1) to isolate the target compound. The relevant fractions were combined and evaporated *in vacuo* to give a white solid which was triturated with acetone, filtered and dried to give **6-methoxy-7-(3-piperidinopropoxy)-4-(1*H*-pyrrolo[2,3-*b*]pyridin-5-yloxy)quinazoline** (58 mg, 20%).

¹H NMR Spectrum: (DMSO-*d*₆) 1.38 (m, 2H), 1.50 (m, 4H), 1.95 (m, 2H), 2.15 (m, 4H), 2.42 (t, 2H), 3.99 (s, 3H), 4.22 (t, 2H), 6.47 (m, 1H), 7.36 (s, 1H), 7.55 (m, 1H), 7.60 (s, 1H), 7.90 (d, 1H), 8.18 (d, 1H), 8.49 (s, 1H) and 11.76 (br s, 1H)

MS (ESI) : 434 (MH)⁺

Elemental analysis

Found

C 63.9

H 6.4 N 15.4

C₂₄H₂₇N₅O₃ 1.0 H₂O

Requires

C 63.8

H 6.5 N 15.5%

Example 314

To a solution of 7-(3-bromopropoxy)-4-(1*H*-indol-5-yloxy)-6-methoxyquinazoline (200mg, 0.47 mmol) in methylene chloride was added 4-piperidinopiperidine (237mg, 1.41mmol) and the reaction heated at 40° C for 1 hour. A further portion of 4-piperidinopiperidine (100mg, 0.59mmol) was added and reaction heated for a further 2 hours.

- 5 The reaction was purified by flash chromatography eluting from methylene chloride to 15% methanol/methylene chloride (+ 1% ammonium hydroxide). The product was evaporated, triturated with ether and filtered to give **4-(indol-5-yloxy)-6-methoxy-7-(3-(4-piperidino)piperidinopropoxy)quinazoline** (200mg, 83%) as a yellow solid.

¹H NMR Spectrum: (CDCl₃) 1.48-2.18 (m, 19H), 2.58 (t, 2H), 3.06 (d, 2H), 4.05 (s, 3H), 4.26 (t, 2H), 6.59 (s, 1H), 7.08 (dd, 1H), 7.28 (d, 1H), 7.36 (s, 1H), 7.50 (d, 1H), 7.63 (s, 1H), 8.30 (s, 1H), 8.59 (s, 1H)

M S: 516 [MH]⁺

The starting material was prepared as follows:

- 15 To a solution of 7-hydroxy-4-(1*H*-indol-5-yloxy)-6-methoxyquinazoline (1g, 3.2 mmol), (prepared as described for the starting material in Example 107), in DMF (50ml) was added powdered potassium carbonate (1.32g, 9.6mmol) and 1,3-dibromopropane (6.43g, 32mmol). The reaction was heated at 50°C for 2 hours. The inorganic material was filtered off and then the DMF removed. The residue was partitioned between methylene
- 20 chloride/water. The organics were separated, dried over MgSO₄, filtered, evaporated in vacuo and purified by flash chromatography eluting from methylene chloride to 5% methanol/95% methylene chloride. The product was concentrated *in vacuo*, triturated with ether and the resulting solid filtered to give 7-(3-bromopropoxy)-4-(1*H*-indol-5-yloxy)-6-methoxyquinazoline (900mg, 66%) as a white solid.

25 ¹H NMR Spectrum: (CDCl₃) 2.46-2.57 (m, 2H), 3.68 (t, 2H), 4.08 (s, 3H), 4.38 (t, 2H), 6.58 (s, 1H), 7.09 (d, 1H), 7.27 (s, 1H), 7.35 (s, 1H), 7.46 (d, 1H), 7.50 (s, 1H), 7.63 (s, 1H), 8.30 (s, 1H), 8.62 (s, 1H)

M S: 428 [MH]⁺

30 **Example 315**

To a solution of 7-hydroxy-6-methoxy-4-(2-methyl-1*H*-indol-5-yloxy)quinazoline (225mg, 0.7 mmol), (prepared as described in Example 49), in DMF was added powdered

potassium carbonate (290mg, 2.1 mmol) and (5*S*)-5-(*p*-toluenesulphonylmethyl)-1-methyl-2-pyrrolidinone (340mg, 1.2mmol). The reaction was then heated at 95°C for 5 hours. The inorganic material was filtered off and the DMF removed by evaporation. The residue was then purified by chromatography eluting from methylene chloride to 12 % methanol/88% methylene chloride (+1% ammonium hydroxide). The product was evaporated, triturated with ether and filtered to give (5*S*)-6-methoxy-4-(2-methyl-1*H*-indol-5-yloxy)-7-(1-methyl-2-oxopyrrolidin-5-ylmethoxy)quinazoline (100mg, 33%) as a white solid.

¹H NMR Spectrum: (DMSO-*d*₆) 1.84-1.96 (m, 1H), 2.10-2.30 (m, 2H), 2.39 (s, 3H), 2.43-2.53 (m, 1H), 2.80 (s, 3H), 3.98 (s, 4H), 4.22 (dd, 1H), 4.40 (dd, 1H), 6.10 (s, 1H), 6.84 (dd, 1H), 7.23 (d, 1H), 7.30 (d, 1H), 7.40 (s, 1H), 7.59 (s, 1H), 8.49 (s, 1H), 10.98 (br s, 1H)

M S: 429 [MH]⁺

Elemental Analysis:	Found	C 64.4	H 5.4	N 12.6
C ₂₄ H ₂₄ N ₄ O ₄ 0.8H ₂ O	Requires	C 64.5	H 5.8	N 12.5%

The starting material was prepared as follows:

(5*S*)-5-(*p*-Toluenesulphonylmethyl)-2-pyrrolidinone (0.8g, 3mmol) was dissolved in dry THF and cooled to -70°C. Lithium diisopropylamide was slowly added and the reaction stirred for 20 minutes before addition of methyl iodide (2ml, excess). The reaction was allowed to warm to ambient temperature for over 2 hours. The reaction was partitioned between ethyl acetate and water, the organic layer separated, dried over MgSO₄, filtered, and evaporated *in vacuo*. The residue was purified by flash chromatography eluting from methylene chloride to 5% methanol/95% methylene chloride and the product evaporated to give (5*S*)-5-(*p*-toluenesulphonyl-methyl)-1-methyl-2-pyrrolidinone (340 mg, 40%) as a brown oil.

¹H NMR Spectrum: (CDCl₃) 2.10-2.44 (m, 4H), 2.48 (s, 3H), 2.76 (s, 3H), 3.30-3.54 (m, 1H), 4.04 (dd, 1H), 4.15 (dd, 1H), 7.38 (d, 2H), 7.78 (d, 2H)

M S: 284 [MH]⁺

Example 316

To a solution of 7-hydroxy-4-(1*H*-indol-5-yloxy)-6-methoxyquinazoline (600mg, 1.95 mmol), (prepared as described for the starting material in Example 107), in DMF (20ml) was added powdered potassium carbonate (540mg, 3.9 mmol) and (5*S*)-5-(*p*-toluene-

5 sulphonylmethyl)-2-pyrrolidinone (580mg, 2.16mmol). The reaction was then heated at 100°C for 4 hours. The inorganic material was filtered off and the DMF removed by evaporation. The residue was then purified by chromatography eluting from methylene chloride to 12 % methanol/88% methylene chloride (+1% ammonium hydroxide). The product was evaporated, triturated with ether, and filtered to give **(5*S*)-4-(1*H*-indol-5-yloxy)-6-methoxy-7-(2-oxopyrrolidin-5-ylmethoxy)quinazoline** (240mg, 31%) as a white solid. ¹H NMR Spectrum: (DMSO-*d*₆) 1.87-2.48 (m, 4H), 3.97 (s, 3H), 4.17 (m, 2H), 6.45 (s, 1H), 6.96 (dd, 1H), 7.38-7.49 (m, 4H), 7.60 (s, 1H), 7.81 (s, 1H), 8.50 (s, 1H)
M S: 405 [MH]⁺

Example 317

10 To a solution of 7-hydroxy-4-(1*H*-indol-5-yloxy)-6-methoxyquinazoline (800mg, 2.6 mmol), (prepared as described for the starting material in Example 107), in DMF (20ml) was added powdered potassium carbonate (1.08g, 7.8mmol) and (5*R*)-5-(*p*-toluenesulphonylmethyl)-2-pyrrolidinone (1.13g, 4.2mmol). The reaction was then heated at 90°C for 4 hours. The inorganic material was filtered off and the DMF removed by evaporation. The residue was then purified by chromatography eluting from methylene chloride to 12% methanol/88% methylene chloride (+1% ammonium hydroxide). A small portion was recolumned using the same gradient. The product was evaporated, triturated with ether and filtered to give **(5*R*)-4-(1*H*-indol-5-yloxy)-6-methoxy-7-(2-oxopyrrolidin-5-ylmethoxy)quinazoline** (70mg, 6.5%) as a white solid.
20 ¹H NMR Spectrum: (DMSO-*d*₆) 1.64-2.45 (m, 4H), 3.78 (m, 1H), 3.99 (s, 3H), 4.18 (t, 2H), 6.42 (s, 1H), 6.97 (dd, 1H), 7.38 -7.48 (m, 3H), 7.60 (s, 1H), 7.73 (s, 2H), 8.48 (s, 1H), 11.18 (br s, 1H)
25 M S: 405 [MH]⁺

The starting material was prepared as follows:

To a solution of (5*R*)-5-hydroxymethyl-2-pyrrolidinone (5.0g, 43mmol) in methylene chloride (100 ml) was added 4-dimethylaminopyridine (15.7g, 129mmol) and *p*-toluenesulphonyl chloride (9.0g, 47mmol). The reaction was stirred at ambient temperature for 16 hours. The reaction was then washed with 1M hydrochloric acid and the organic layer

separated. This was then dried over MgSO_4 , filtered and evaporated to give (5*R*)-5-(*p*-toluenesulphonylmethyl)-2-pyrrolidinone (10.3g, 89%) as a white solid.

^1H NMR Spectrum: (CDCl_3) 1.68-1.86 (m, 1H), 2.16-2.38 (m, 3H), 2.48 (s, 3H), 3.86-3.96 (m, 2H), 4.08 (dd, 1H), 6.20 (br s, 1H), 7.38 (d, 2H), 7.80 (d, 2H)

5 M S: 270 [MH]⁺

Example 318

To a suspension of 7-hydroxy-6-methoxy-4-(2-methyl-1*H*-indol-5-yloxy)quinazoline (1.36g, 4.24mmol), (prepared as described in Example 49), in DMF (70ml), was added
10 potassium carbonate (2.34g, 17.0mmol, 4eq.) followed by (5*R*)-5-(*p*-toluenesulphonylmethyl)-2-pyrrolidinone (1.25g, 4.66mmol, 1.1eq.), (prepared as described for the starting material in Example 317), and the resulting yellow suspension heated at reflux. After 4 hours, some starting material remained, and a further addition of (5*R*)-5-(*p*-toluenesulphonylmethyl)-2-pyrrolidinone (0.57g, 2.12mmol, 0.5eq.) was made. The reaction
15 was heated at reflux for a further 2 hours resulting in consumption of starting material. The reaction was cooled to ambient temperature, the inorganic residue filtered off and the filtrate evaporated *in vacuo* to leave a brown oil which was purified by column chromatography (methylene chloride/methanol, (100/0 to 90/10)) to give a light brown oil. Trituration with ether afforded a thick oil, which upon chromatography eluting as above gave a yellow oil.
20 Trituration of this oil with ether gave an initial crop of (5*R*)-6-methoxy-4-(2-methyl-1*H*-indol-5-yloxy)-7-(2-oxopyrrolidin-5-ylmethoxy)quinazoline (5mg) as an off-white solid (ca. 90% pure by nmr). Chromatography of the residues (eluting as above) followed by ether trituration gave further crops of (5*R*)-6-methoxy-4-(2-methyl-1*H*-indol-5-yloxy)-7-(2-oxopyrrolidin-5-ylmethoxy)quinazoline as a white solid (180mg, >95% pure by nmr), as an
25 off-white solid (800mg, ca. 95% pure by nmr).

^1H NMR Spectrum: (DMSO-d_6) 1.8-2.2 (m, 5H), 2.4 (s, 3H), 4.0 (br s, 3H), 4.1-4.2 (m, 2H), 6.1 (br s, 1H), 6.9 (dd, 1H), 7.2 (d, 1H), 7.3 (d, 1H), 7.4 (s, 1H), 7.6 (s, 1H), 7.8 (s, 1H), 8.5 (s, 1H), 11.0 (br s, 1H)

M S: 419 [MH]⁺

30	Elemental analysis:	Found	C 60.8	H 5.3	N 12.1
	$\text{C}_{23}\text{H}_{22}\text{N}_4\text{O}_4 \cdot 2\text{H}_2\text{O}$	Requires	C 60.8	H 5.7	N 12.3%

Example 319

To a solution of 7-hydroxy-6-methoxy-4-(2-methyl-1*H*-indol-5-yloxy)quinazoline (4.8g, 15.7mmol), (prepared as described in Example 49), in DMF (100 ml), was added potassium carbonate (6.5g, 47mmol) and 3-chloropropyl piperidine (3.3g, 20.4mmol). The reaction was then heated to 100°C for 4 hours. The inorganic material was filtered off and the DMF removed by evaporation. The residue was then purified by chromatography eluting from methylene chloride to 10 % methanol/90% methylene chloride (+1% ammonium hydroxide). The relevant fractions were concentrated and the residue dissolved in ethyl acetate. Hexane was added and the precipitate was filtered off. The filtrate was evaporated and the residue was triturated with ether and filtered to give **6-methoxy-4-(1-(3-piperidinopropyl)-1*H*-indol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline** (170mg, 1.9%) as a white solid.

¹H NMR Spectrum: (DMSO-*d*₆) 1.38 (br s, 4H), 1.50 (br s, 8H), 1.92 (m, 4H), 2.14-2.48 (m, 12H), 3.98 (s, 3H), 4.24 (t, 4H), 6.43 (s, 1H), 7.02 (d, 1H), 7.38 (s, 1H), 7.42 (s, 2H), 7.53 (d, 1H), 7.58 (s, 1H), 8.44 (s, 1H)

MS: 558 [MH]⁺

Example 320

A mixture of (2*R*)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (6.201g, 16.4 mmol), (prepared as described for the starting material in Example 269), and piperidine (4.8 ml, 49.3 mmol) in DMF (100 ml) was stirred at 60°C for 24 hours and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified on silica gel, eluting with dichloromethane, dichloromethane/methanol (95/5) then dichloromethane/methanol/0.880 aqueous ammonia (89:10:1). The product was then recrystallised from acetonitrile to give **(2*R*)-6-methoxy-(2-methyl-1*H*-indol-5-yloxy)-7-(2-hydroxy-3-piperidinopropoxy)quinazoline** (3.33g, 44 %) as an off-white solid.

¹H NMR Spectrum: (DMSO-*d*₆) 1.35 (m, 2H), 1.51 (m, 4H), 2.30-2.40 (m, 9H), 3.98 (s, 3H), 4.08 (m, 2H), 4.21 (m, 1H), 4.86 (m, 1H), 6.10 (s, 1H), 6.87 (dd, 1H), 7.25 (d, 1H), 7.30 (d, 1H), 7.40 (s, 1H), 7.60 (s, 1H), 8.45 (s, 1H) and 10.98 (br s, 1H)

MS (ESI) : 463 (MH)⁺

Elemental analysis: Found C 66.5 H 6.6 N 12.0

$C_{26}H_{30}N_4O_4 \cdot 0.4H_2O$

Requires

C 66.5 H 6.6 N 11.9%

Example 321


A mixture of (2*S*)-6-methoxy-4-(2-methylindol-5-yloxy)-7-(oxiran-2-ylmethoxy)quinazoline (175 mg 0.46 mmol), (prepared as described for the starting material in Example 304), and piperidine (0.14 ml, 1.39 mmol) in DMF (5 ml) was stirred at 60°C for 24 hours and allowed to cool to ambient temperature. The solvents were removed *in vacuo* and the residue purified on silica gel, gradient elution eluting with dichloromethane, dichloromethane/methanol (95/5) then dichloromethane/methanol/0.880 aqueous ammonia (89:10:1). The product was then recrystallised from acetonitrile to give (2*S*)-6-methoxy-(2-methyl-1*H*-indol-5-yloxy)-7-(2-hydroxy-3-piperidinopropoxy)quinazoline (88 mg, 41 %) as an off-white solid.

¹H NMR Spectrum: (DMSO₆) 1.35 (m, 2H), 1.51 (m, 4H), 2.30-2.40 (m, 9H), 3.98 (s, 3H), 4.08 (m, 2H), 4.21 (m, 1H), 4.86 (m, 1H), 6.10 (s, 1H), 6.87 (dd, 1H), 7.25 (d, 1H) 7.30 (d, 1H), 7.40 (s, 1H), 7.60 (s, 1H), 8.45 (s, 1H) and 10.98 (br s, 1H)

MS (ESI) : 463 (MH)⁺

Elemental analysis:	Found	C 66.2	H 6.8	N 11.9
---------------------	-------	--------	-------	--------

$C_{26}H_{30}N_4O_4 \cdot 0.5H_2O$	Requires	C 66.2	H 6.6	N 11.9%
------------------------------------	----------	--------	-------	---------

Example 322

A solution of 4-chloro-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline (1.22g, 3.65 mmol), (prepared as described for the starting material in Example 241), 4-fluoro-5-hydroxy-2-methylindole (723mg, 4.38 mmol), (prepared as described for the starting material in Example 237), in DMF (20ml) containing potassium carbonate (756mg, 5.48 mmol) was stirred at 95°C for 3 hours. After cooling, the mixture was filtered and the filtrate was evaporated. The residue was purified by column chromatography eluting with methylene chloride/methanol (9/1) followed by methylene chloride/methanol/methanol saturated with ammonia (90/5/5). The fractions containing the expected product were combined and evaporated. The residue was triturated with ether, filtered, washed with ether and dried

under vacuum. The solid was dissolved in methylene chloride/ethyl acetate and the minimum of methanol, filtered and the volatiles were removed under vacuum. The solid was triturated with ether, filtered, washed with ether and dried under vacuum at 50°C to give **4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline** (1.06g, 62%).

MS-ESI : 465 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 1.1-1.3 (m, 2H) ; 1.35-1.5 (m, 1H) ; 1.6-1.9 (m, 6H) ; 2.12 (s, 3H) ; 2.4 (s, 3H) ; 2.75 (d, 2H) ; 3.95 (s, 3H) ; 4.22 (t, 2H) ; 6.2 (s, 1H) ; 6.95 (dd, 1H) ; 7.15 (d, 1H) ; 7.4 (s, 1H) ; 7.6 (s, 1H) ; 8.5 (s, 1H)

Example 323

Sodium hydride (71 mg, 1.8 mmol) was added to 5-hydroxy-2-methylbenzimidazole (204 mg, 0.89 mmol) in anhydrous DMF (2.5 ml) under an argon atmosphere. The mixture was stirred at ambient temperature for 10 minutes. 4-Chloro-6,7-dimethoxyquinazoline (200 mg, 0.89 mmol) was added and the reaction mixture stirred at 95 °C for 2 hours. Upon cooling to ambient temperature the mixture was poured in water and extracted with ethyl acetate. The organic phase was washed with brine, dried (MgSO₄), silica was added and the solvent evaporated off. The obtained powder was placed on the top of a disposable silica column (ISOLUTE) and the product eluted off using a gradient of methanol/dichloromethane (3/97, 5/95, 8/92). Evaporation of the solvent gave **6,7-dimethoxy-4-(2-methyl-1H-benzimidazol-6-yloxy)quinazoline** (145 mg, 48%).

¹H NMR Spectrum: (DMSO-d₆) 2.50 (s, 3H); 3.95 (s, 3H); 4.0 (s, 3H); 7.05 (d, 1H); 7.38 (s, 1H); 7.39 (d, 1H); 7.51 (d, 1H); 7.60 (s, 1H); 8.50 (s, 1H)

MS (ESI) : 337 [MH]⁺

Example 324




- 7-Hydroxyquinazoline (87 mg, 0.6 mmol) and potassium carbonate (110 mg, 0.8 mmol) were added to 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (180 mg, 0.53 mmol), (prepared as described for the starting material in Example 1), in suspension in DMF (3 ml) under an argon atmosphere. The reaction mixture was heated to 100 °C for 90 minutes. Upon cooling to ambient temperature the reaction was diluted with ethyl acetate and a saturated ammonium chloride solution. The aqueous phase was re-extracted with ethyl acetate, the organic phases combined, dried (MgSO₄) and the solvent evaporated. The residue was purified by flash chromatography using a gradient of methanol/dichloromethane (3/97, 4/96, 5/95). Evaporation of the solvent and trituration of the solid with ether gave 6-methoxy-7-(3-morpholinopropoxy)-4-(quinazolin-7-yloxy)quinazoline (197 mg, 83%).
- ¹H NMR Spectrum (DMSO-d₆) 2.01 (t, 2H); 2.47 (m, 4H); 2.49 (m, 2H); 3.60 (m, 4H); 4.01 (s, 3H); 4.29 (t, 2H); 7.45 (s, 1H); 7.65 (s, 1H); 7.80 (d, 1H); 8.01 (d, 1H); 8.32 (d, 1H); 8.60 (s, 1H); 9.34 (s, 1H); 9.69 (s, 1H)
- MS (ESI) : 448 [MH]⁺
- Elemental analysis: Found C 63.4 H 5.7 N 15.6
C₂₄H₂₅N₅O₄ · 0.4 H₂O Requires C 63.4 H 5.7 N 15.4%

The starting material was prepared as follows:

- Raney Nickel (about 200 mg), (prewashed several times with ethanol), was added to a solution of 7-hydroxy-4-thiomethylquinazoline (400mg, 2.08 mmol), (Tet. Lett. 1999, 40, 3881), and the solution was refluxed for 1 hour. Raney Nickel (100 mg) was added and the mixture was refluxed for a further 1 hour. The mixture was filtered, washed with ethanol and the volatiles were removed under vacuum. The residue was purified by column chromatography eluting with methylene chloride/methanol (97/3 followed by 96/4) to give 7-hydroxyquinazoline (62mg, 20%).

Example 325



Using an analogous procedure to that described in Example 201, 7-hydroxy-4-(indol-6-ylamino)-6-methoxyquinazoline (98 mg, 0.32 mmol), (prepared as described for the starting material in Example 217), was reacted with 5-(2-hydroxyethyl)-4-methylthiazole (69 mg, 0.48 mmol) to give **6-methoxy-4-(indol-6-ylamino)-7-(2-(4-methylthiazol-5-**

5 **yl)ethoxy)quinazoline** (47 mg, 34 %).

MS - ESI : 432 [MH]⁺

¹H NMR Spectrum: (DMSO-d₆) 2.4 (s, 3H) ; 3.3 (t, 2H) ; 4.0 (s, 3H) ; 4.35 (t, 2H) ; 6.45 (s, 1H) ; 7.2 (s, 1H) ; 7.25-7.4 (m, 2H) ; 7.55 (d, 1H) ; 7.9 (s, 1H) ; 8.05 (s, 1H) ; 8.45 (s, 1H) ; 8.87 (s, 1H) ; 9.45 (s, 1H)

Example 326

The following illustrate representative pharmaceutical dosage forms containing the compound of formula I, or a pharmaceutically acceptable salt thereof (hereafter compound X), for therapeutic or prophylactic use in humans:

(a)	<u>Tablet I</u>	<u>mg/tablet</u>
	Compound X	100
	Lactose Ph.Eur.....	182.75
	Croscarmellose sodium	12.0
20	Maize starch paste (5% w/v paste)	2.25
	Magnesium stearate	3.0
(b)	<u>Tablet II</u>	<u>mg/tablet</u>
	Compound X	50
25	Lactose Ph.Eur.....	223.75
	Croscarmellose sodium	6.0
	Maize starch.....	15.0
	Polyvinylpyrrolidone (5% w/v paste).....	2.25
	Magnesium stearate	3.0
30	(c)	<u>Tablet III</u>
		<u>mg/tablet</u>
	Compound X	1.0

- 313 -

		Lactose Ph.Eur.....	93.25
		Croscarmellose sodium	4.0
		Maize starch paste (5% w/v paste)	0.75
		Magnesium stearate	1.0
5	(d)	<u>Capsule</u>	<u>mg/capsule</u>
		Compound X	10
		Lactose Ph.Eur.....	488.5
		Magnesium stearate	1.5
10	(e)	<u>Injection I</u>	<u>(50 mg/ml)</u>
		Compound X	5.0% w/v
		1N Sodium hydroxide solution.....	15.0% v/v
		0.1N Hydrochloric acid (to adjust pH to 7.6)	
15		Polyethylene glycol 400	4.5% w/v
		Water for injection to 100%	
	(f)	<u>Injection II</u>	<u>10 mg/ml</u>
		Compound X	1.0% w/v
20		Sodium phosphate BP.....	3.6% w/v
		0.1N Sodium hydroxide solution.....	15.0% v/v
		Water for injection to 100%	
	(g)	<u>Injection III</u>	<u>(1mg/ml,buffered to pH6)</u>
25		Compound X	0.1% w/v
		Sodium phosphate BP.....	2.26% w/v
		Citric acid	0.38% w/v
		Polyethylene glycol 400	3.5% w/v
		Water for injection to 100%	

30


Note

- 314 -

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

Claims:

1. The use of a compound of the formula I:



(I)

wherein:

ring C is an 8, 9, 10, 12 or 13-membered bicyclic or tricyclic moiety which moiety may be saturated or unsaturated, which may be aromatic or non-aromatic, and which optionally may contain 1-3 heteroatoms selected independently from O, N and S;

Z is -O-, -NH-, -S-, -CH₂- or a direct bond;

n is an integer from 0 to 5;

m is an integer from 0 to 3;

R² represents hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C₁₋₃alkyl, C₁₋₃alkoxy, C₁₋₃alkylsulphanyl, -NR³R⁴ (wherein R³ and R⁴, which may be the same or different, each represents hydrogen or C₁₋₃alkyl), or R⁵X¹- (wherein X¹ represents a direct bond, -O-, -CH₂-, -OC(O)-, -C(O)-, -S-, -SO-, -SO₂-, -NR⁶C(O)-, -C(O)NR⁷-, -SO₂NR⁸-, -NR⁹SO₂- or -NR¹⁰- (wherein R⁶, R⁷, R⁸, R⁹ and R¹⁰ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), and R⁵ is selected from one of the following twenty-two groups:

1) hydrogen, oxiranylC₁₋₄alkyl or C₁₋₅alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, chloro, bromo and amino;

2) C₁₋₅alkylX²C(O)R¹¹ (wherein X² represents -O- or -NR¹²- (in which R¹² represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹¹ represents C₁₋₃alkyl, -NR¹³R¹⁴ or -OR¹⁵

(wherein R¹³, R¹⁴ and R¹⁵ which may be the same or different each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl));

- 3) $C_{1,3}alkylX^3R^{16}$ (wherein X^3 represents -O-, -S-, -SO-, -SO₂-, -OC(O)-, -NR¹⁷C(O)-, -C(O)NR¹⁸-, -SO₂NR¹⁹-, -NR²⁰SO₂- or -NR²¹- (wherein R¹⁷, R¹⁸, R¹⁹, R²⁰ and R²¹ each independently represents hydrogen, C_{1,3}alkyl or C_{1,3}alkoxyC_{2,3}alkyl) and R¹⁶ represents hydrogen, C_{1,3}alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C_{1,3}alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C_{1,4}alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1,4}cyanoalkyl, C_{1,4}alkyl, C_{1,4}hydroxyalkyl, C_{1,4}alkoxy, C_{1,4}alkoxyC_{1,4}alkyl, C_{1,4}alkylsulphonylC_{1,4}alkyl, C_{1,4}alkoxycarbonyl, C_{1,4}aminoalkyl, C_{1,4}alkylamino, di(C_{1,4}alkyl)amino, C_{1,4}alkylaminoC_{1,4}alkyl, di(C_{1,4}alkyl)aminoC_{1,4}alkyl, C_{1,4}alkylaminoC_{1,4}alkoxy, di(C_{1,4}alkyl)aminoC_{1,4}alkoxy and a group $-(O-)_f(C_{1,4}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C_{1,4}alkyl));
- 4) $C_{1,3}alkylX^4C_{1,3}alkylX^5R^{22}$ (wherein X^4 and X^5 which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR²³C(O)-, -C(O)NR²⁴-, -SO₂NR²⁵-, -NR²⁶SO₂- or -NR²⁷- (wherein R²³, R²⁴, R²⁵, R²⁶ and R²⁷ each independently represents hydrogen, C_{1,3}alkyl or C_{1,3}alkoxyC_{2,3}alkyl) and R²² represents hydrogen, C_{1,3}alkyl or C_{1,3}alkoxyC_{2,3}alkyl);
- 5) R²⁸ (wherein R²⁸ is a 5-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1,4}cyanoalkyl, C_{1,4}alkyl, C_{1,4}hydroxyalkyl, C_{1,4}alkoxy, C_{1,4}alkoxyC_{1,4}alkyl, C_{1,4}alkylsulphonylC_{1,4}alkyl, C_{1,4}alkoxycarbonyl, C_{1,4}aminoalkyl, C_{1,4}alkylamino, di(C_{1,4}alkyl)amino, C_{1,4}alkylaminoC_{1,4}alkyl, di(C_{1,4}alkyl)aminoC_{1,4}alkyl, C_{1,4}alkylaminoC_{1,4}alkoxy, di(C_{1,4}alkyl)aminoC_{1,4}alkoxy and a group $-(O-)_f(C_{1,4}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C_{1,4}alkyl));
- 6) C_{1,3}alkylR²⁸ (wherein R²⁸ is as defined herein);
- 7) C_{2,3}alkenylR²⁸ (wherein R²⁸ is as defined herein);
- 8) C_{2,3}alkynylR²⁸ (wherein R²⁸ is as defined herein);

- 9) R^{29} (wherein R^{29} represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} hydroxyalkyl, C_{1-4} aminoalkyl, C_{1-4} alkylamino, C_{1-4} hydroxyalkoxy, carboxy, trifluoromethyl, cyano, -
 5 $C(O)NR^{30}R^{31}$, $-NR^{32}C(O)R^{33}$ (wherein R^{30} , R^{31} , R^{32} and R^{33} , which may be the same or different, each represents hydrogen, C_{1-4} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and a group $-(O)_f(C_{1-4}alkyl)_gringD$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which
 10 cyclic group may bear one or more substituents selected from C_{1-4} alkyl));
- 10) $C_{1-5}alkylR^{29}$ (wherein R^{29} is as defined herein);
- 11) $C_{2-5}alkenylR^{29}$ (wherein R^{29} is as defined herein);
- 12) $C_{2-5}alkynylR^{29}$ (wherein R^{29} is as defined herein);
- 13) $C_{1-5}alkylX^6R^{29}$ (wherein X^6 represents -O-, -S-, -SO-, -SO₂-, -NR³⁴C(O)-, -C(O)NR³⁵-, -
 15 SO₂NR³⁶-, -NR³⁷SO₂- or -NR³⁸- (wherein R^{34} , R^{35} , R^{36} , R^{37} and R^{38} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{29} is as defined herein);
- 14) $C_{2-5}alkenylX^7R^{29}$ (wherein X^7 represents -O-, -S-, -SO-, -SO₂-, -NR³⁹C(O)-, -C(O)NR⁴⁰-, -
 SO₂NR⁴¹-, -NR⁴²SO₂- or -NR⁴³- (wherein R^{39} , R^{40} , R^{41} , R^{42} and R^{43} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{29} is as defined herein);
- 20 15) $C_{2-5}alkynylX^8R^{29}$ (wherein X^8 represents -O-, -S-, -SO-, -SO₂-, -NR⁴⁴C(O)-, -C(O)NR⁴⁵-, -
 SO₂NR⁴⁶-, -NR⁴⁷SO₂- or -NR⁴⁸- (wherein R^{44} , R^{45} , R^{46} , R^{47} and R^{48} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{29} is as defined herein);
- 16) $C_{1-4}alkylX^9C_{1-4}alkylR^{29}$ (wherein X^9 represents -O-, -S-, -SO-, -SO₂-, -NR⁴⁹C(O)-, -
 C(O)NR⁵⁰-, -SO₂NR⁵¹-, -NR⁵²SO₂- or -NR⁵³- (wherein R^{49} , R^{50} , R^{51} , R^{52} and R^{53} each
 25 independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{29} is as defined herein);
- 17) $C_{1-4}alkylX^9C_{1-4}alkylR^{28}$ (wherein X^9 and R^{28} are as defined herein);
- 18) $C_{2-5}alkenyl$ which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, $C_{1-4}alkylamino$, N,N-di($C_{1-4}alkyl$)amino, aminosulphonyl, N- $C_{1-4}alkylaminosulphonyl$ and N,N-di($C_{1-4}alkyl$)aminosulphonyl;
 30

- 19) $C_{2,5}$ alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, $C_{1,4}$ alkylamino, $\underline{N,N}$ -di($C_{1,4}$ alkyl)amino, aminosulphonyl, \underline{N} - $C_{1,4}$ alkylaminosulphonyl and $\underline{N,N}$ -di($C_{1,4}$ alkyl)aminosulphonyl;
- 20) $C_{2,5}$ alkenyl $X^9C_{1,4}$ alkyl R^{28} (wherein X^9 and R^{28} are as defined herein);
- 5 21) $C_{2,5}$ alkynyl $X^9C_{1,4}$ alkyl R^{28} (wherein X^9 and R^{28} are as defined herein); and
- 22) $C_{1,4}$ alkyl $R^{54}(C_{1,4}alkyl)_q(X^9)_rR^{55}$ (wherein X^9 is as defined herein, q is 0 or 1, r is 0 or 1, and R^{54} and R^{55} are each independently selected from hydrogen, $C_{1,3}$ alkyl, cyclopentyl, cyclohexyl and a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which $C_{1,3}$ alkyl group may bear 1 or 2 substituents selected
- 10 from oxo, hydroxy, halogeno and $C_{1,4}$ alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, $C_{1,4}$ cyanoalkyl, $C_{1,4}$ alkyl, $C_{1,4}$ hydroxyalkyl, $C_{1,4}$ alkoxy, $C_{1,4}$ alkoxy $C_{1,4}$ alkyl, $C_{1,4}$ alkylsulphonyl $C_{1,4}$ alkyl, $C_{1,4}$ alkoxycarbonyl, $C_{1,4}$ aminoalkyl, $C_{1,4}$ alkylamino, di($C_{1,4}$ alkyl)amino, $C_{1,4}$ alkylamino $C_{1,4}$ alkyl, di($C_{1,4}$ alkyl)amino $C_{1,4}$ alkyl, $C_{1,4}$ alkylamino $C_{1,4}$ alkoxy, di($C_{1,4}$ alkyl)amino $C_{1,4}$ alkoxy and a
- 15 group $-(-O-)_f(C_{1,4}alkyl)_g\text{ringD}$ (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from $C_{1,4}$ alkyl), with the proviso that R^{54} cannot be hydrogen);
- and additionally wherein any $C_{1,3}$ alkyl, $C_{2,5}$ alkenyl or $C_{2,5}$ alkynyl group in R^5X^1 may bear one
- 20 or more substituents selected from hydroxy, halogeno and amino);
- R^1 represents hydrogen, oxo, halogeno, hydroxy, $C_{1,4}$ alkoxy, $C_{1,4}$ alkyl, $C_{1,4}$ alkoxymethyl, $C_{1,4}$ alkanoyl, $C_{1,4}$ haloalkyl, cyano, amino, $C_{2,5}$ alkenyl, $C_{2,5}$ alkynyl, $C_{1,3}$ alkanoyloxy, nitro, $C_{1,4}$ alkanoylamino, $C_{1,4}$ alkoxycarbonyl, $C_{1,4}$ alkylsulphanyl, $C_{1,4}$ alkylsulphinyl, $C_{1,4}$ alkylsulphonyl, carbamoyl, \underline{N} - $C_{1,4}$ alkylcarbamoyl, $\underline{N,N}$ -di($C_{1,4}$ alkyl)carbamoyl,
- 25 aminosulphonyl, \underline{N} - $C_{1,4}$ alkylaminosulphonyl, $\underline{N,N}$ -di($C_{1,4}$ alkyl)aminosulphonyl, \underline{N} -($C_{1,4}$ alkylsulphonyl)amino, \underline{N} -($C_{1,4}$ alkylsulphonyl)- \underline{N} -($C_{1,4}$ alkyl)amino, $\underline{N,N}$ -di($C_{1,4}$ alkylsulphonyl)amino, a $C_{3,7}$ alkylene chain joined to two ring C carbon atoms, $C_{1,4}$ alkanoylamino $C_{1,4}$ alkyl, carboxy or a group $R^{56}X^{10}$ (wherein X^{10} represents a direct bond, -O-, -CH₂-, -OC(O)-, -C(O)-, -S-, -SO-, -SO₂-, -NR⁵⁷C(O)-, -C(O)NR⁵⁸-, -SO₂NR⁵⁹-, -NR⁶⁰SO₂- or -
- 30 NR⁶¹- (wherein R^{57} , R^{58} , R^{59} , R^{60} and R^{61} each independently represents hydrogen, $C_{1,3}$ alkyl or $C_{1,3}$ alkoxy $C_{2,3}$ alkyl), and R^{56} is selected from one of the following twenty-two groups:

- 1) hydrogen, oxiranylC₁₋₄alkyl or C₁₋₅alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, chloro, bromo and amino;
- 2) C₁₋₅alkylX¹¹C(O)R⁶² (wherein X¹¹ represents -O- or -NR⁶³- (in which R⁶³ represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁶² represents C₁₋₃alkyl, -NR⁶⁴R⁶⁵ or -OR⁶⁶ (wherein R⁶⁴, R⁶⁵ and R⁶⁶ which may be the same or different each represents hydrogen, C₁₋₅alkyl or C₁₋₃alkoxyC₂₋₃alkyl));
- 3) C₁₋₅alkylX¹²R⁶⁷ (wherein X¹² represents -O-, -S-, -SO-, -SO₂-, -OC(O)-, -NR⁶⁸C(O)-, -C(O)NR⁶⁹-, -SO₂NR⁷⁰-, -NR⁷¹SO₂- or -NR⁷²- (wherein R⁶⁸, R⁶⁹, R⁷⁰, R⁷¹ and R⁷² each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁶⁷ represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group -(O-)_f(C₁₋₄alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));
- 4) C₁₋₅alkylX¹³C₁₋₅alkylX¹⁴R⁷³ (wherein X¹³ and X¹⁴ which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR⁷⁴C(O)-, -C(O)NR⁷⁵-, -SO₂NR⁷⁶-, -NR⁷⁷SO₂- or -NR⁷⁸- (wherein R⁷⁴, R⁷⁵, R⁷⁶, R⁷⁷ and R⁷⁸ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁷³ represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl);
- 5) R⁷⁹ (wherein R⁷⁹ is a 5-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₄cyanoalkyl, C₁₋₄alkyl, C₁₋₄hydroxyalkyl, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, C₁₋₄alkylsulphonylC₁₋₄alkyl, C₁₋₄alkoxycarbonyl, C₁₋₄aminoalkyl, C₁₋₄alkylamino, di(C₁₋₄alkyl)amino, C₁₋₄alkylaminoC₁₋₄alkyl, di(C₁₋₄alkyl)aminoC₁₋₄alkyl, C₁₋₄alkylaminoC₁₋₄alkoxy, di(C₁₋₄alkyl)aminoC₁₋₄alkoxy and a group -(O-)_f(C₁₋₄alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected

independently from O, S and N, which cyclic group may bear one or more substituents selected from C₁₋₄alkyl));

6) C₁₋₅alkylR⁷⁹ (wherein R⁷⁹ is as defined herein);

7) C₂₋₅alkenylR⁷⁹ (wherein R⁷⁹ is as defined herein);

5 8) C₂₋₅alkynylR⁷⁹ (wherein R⁷⁹ is as defined herein);

9) R⁸⁰ (wherein R⁸⁰ represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄hydroxyalkyl, C₁₋

10 ₄aminoalkyl, C₁₋₄alkylamino, C₁₋₄hydroxyalkoxy, carboxy, trifluoromethyl, cyano, -C(O)NR⁸¹R⁸², -NR⁸³C(O)R⁸⁴ (wherein R⁸¹, R⁸², R⁸³ and R⁸⁴, which may be the same or different, each represents hydrogen, C₁₋₄alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and a group -(O)_f(C₁₋₄alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which
15 cyclic group may bear one or more substituents selected from C₁₋₄alkyl));

10) C₁₋₅alkylR⁸⁰ (wherein R⁸⁰ is as defined herein);

11) C₂₋₅alkenylR⁸⁰ (wherein R⁸⁰ is as defined herein);

12) C₂₋₅alkynylR⁸⁰ (wherein R⁸⁰ is as defined herein);

13) C₁₋₅alkylX¹⁵R⁸⁰ (wherein X¹⁵ represents -O-, -S-, -SO-, -SO₂-, -NR⁸⁵C(O)-, -C(O)NR⁸⁶-, -
20 SO₂NR⁸⁷-, -NR⁸⁸SO₂- or -NR⁸⁹- (wherein R⁸⁵, R⁸⁶, R⁸⁷, R⁸⁸ and R⁸⁹ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁸⁰ is as defined herein);

14) C₂₋₅alkenylX¹⁶R⁸⁰ (wherein X¹⁶ represents -O-, -S-, -SO-, -SO₂-, -NR⁹⁰C(O)-, -C(O)NR⁹¹-, -SO₂NR⁹²-, -NR⁹³SO₂- or -NR⁹⁴- (wherein R⁹⁰, R⁹¹, R⁹², R⁹³ and R⁹⁴ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁸⁰ is as defined herein);


15) C₂₋₅alkynylX¹⁷R⁸⁰ (wherein X¹⁷ represents -O-, -S-, -SO-, -SO₂-, -NR⁹⁵C(O)-, -C(O)NR⁹⁶-, -SO₂NR⁹⁷-, -NR⁹⁸SO₂- or -NR⁹⁹- (wherein R⁹⁵, R⁹⁶, R⁹⁷, R⁹⁸ and R⁹⁹ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁸⁰ is as defined herein);

16) C₁₋₄alkylX¹⁸C₁₋₄alkylR⁸⁰ (wherein X¹⁸ represents -O-, -S-, -SO-, -SO₂-, -NR¹⁰⁰C(O)-, -C(O)NR¹⁰¹-, -SO₂NR¹⁰²-, -NR¹⁰³SO₂- or -NR¹⁰⁴- (wherein R¹⁰⁰, R¹⁰¹, R¹⁰², R¹⁰³ and R¹⁰⁴ each
30 independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁸⁰ is as defined herein);

17) C₁₋₄alkylX¹⁸C₁₋₄alkylR⁷⁹ (wherein X¹⁸ and R⁷⁹ are as defined herein);

- 18) $C_{2,5}$ alkenyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, $C_{1,4}$ alkylamino, N,N -di($C_{1,4}$ alkyl)amino, aminosulphonyl, N - $C_{1,4}$ alkylaminosulphonyl and N,N -di($C_{1,4}$ alkyl)aminosulphonyl;
- 19) $C_{2,5}$ alkynyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, $C_{1,4}$ alkylamino, N,N -di($C_{1,4}$ alkyl)amino, aminosulphonyl, N - $C_{1,4}$ alkylaminosulphonyl and N,N -di($C_{1,4}$ alkyl)aminosulphonyl;
- 20) $C_{2,5}$ alkenyl $X^{18}C_{1,4}$ alkyl R^{79} (wherein X^{18} and R^{79} are as defined herein);
- 21) $C_{2,5}$ alkynyl $X^{18}C_{1,4}$ alkyl R^{79} (wherein X^{18} and R^{79} are as defined herein); and
- 22) $C_{1,4}$ alkyl $R^{105}(C_{1,4}alkyl)_x(X^{18})_yR^{106}$ (wherein X^{18} is as defined herein, x is 0 or 1, y is 0 or 1, and R^{105} and R^{106} are each independently selected from hydrogen, $C_{1,3}$ alkyl, cyclopentyl, cyclohexyl and a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which $C_{1,3}$ alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and $C_{1,4}$ alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, $C_{1,4}$ cyanoalkyl, $C_{1,4}$ alkyl, $C_{1,4}$ hydroxyalkyl, $C_{1,4}$ alkoxy, $C_{1,4}$ alkoxy $C_{1,4}$ alkyl, $C_{1,4}$ alkylsulphonyl $C_{1,4}$ alkyl, $C_{1,4}$ alkoxycarbonyl, $C_{1,4}$ aminoalkyl, $C_{1,4}$ alkylamino, di($C_{1,4}$ alkyl)amino, $C_{1,4}$ alkylamino $C_{1,4}$ alkyl, di($C_{1,4}$ alkyl)amino $C_{1,4}$ alkyl, $C_{1,4}$ alkylamino $C_{1,4}$ alkoxy, di($C_{1,4}$ alkyl)amino $C_{1,4}$ alkoxy and a group $-(O-)_f(C_{1,4}alkyl)_g$ ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from $C_{1,4}$ alkyl) with the proviso that R^{105} cannot be hydrogen);
- and additionally wherein any $C_{1,5}$ alkyl, $C_{2,5}$ alkenyl or $C_{2,5}$ alkynyl group in $R^{56}X^{10}$ may bear one or more substituents selected from hydroxy, halogeno and amino);
- or a salt thereof in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals such as humans.

2. The use of a compound of the formula I according to claim 1:



wherein:

ring C is a 9-10-membered bicyclic moiety which may be saturated or unsaturated, which may
 10 be aromatic or non-aromatic, and which optionally may contain 1-3 heteroatoms selected independently from O, N and S;

Z is -O-, -NH-, -S-, -CH₂- or a direct bond;

R¹ represents hydrogen, oxo, halogeno, hydroxy, C₁₋₄alkoxy, C₁₋₄alkyl, C₁₋₄alkoxymethyl, C₁₋₄alkanoyl, C₁₋₄haloalkyl, cyano, amino, C₂₋₅alkenyl, C₂₋₅alkynyl, C₁₋₃alkanoyloxy, nitro, C₁₋₄alkanoylamino, C₁₋₄alkoxycarbonyl, C₁₋₄alkylsulphanyl, C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, carbamoyl, N-C₁₋₄alkylcarbamoyl, N,N-di(C₁₋₄alkyl)carbamoyl, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl, N,N-di(C₁₋₄alkyl)aminosulphonyl, N-(C₁₋₄alkylsulphonyl)amino, N-(C₁₋₄alkylsulphonyl)-N-(C₁₋₄alkyl)amino, N,N-di(C₁₋₄alkylsulphonyl)amino or a C₃₋₇alkylene chain joined to two ring C carbon atoms;

20 n is an integer from 0 to 5;

m is an integer from 0 to 3;

R² represents hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C₁₋₃alkyl, C₁₋₃alkoxy, C₁₋₃alkylsulphanyl, -NR³R⁴ (wherein R³ and R⁴, which may be the same or different, each represents hydrogen or C₁₋₃alkyl), or R⁵X¹- (wherein X¹ represents a direct bond, -O-, -CH₂-, -OC(O)-, -C(O)-, -S-, -SO-, -SO₂-, -NR⁶C(O)-, -C(O)NR⁷-, -SO₂NR⁸-, -NR⁹SO₂- or -NR¹⁰- (wherein R⁶, R⁷, R⁸, R⁹ and R¹⁰ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl), and R⁵ is selected from one of the following twenty-one groups:

1) hydrogen or C₁₋₅alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino;

2) C₁₋₅alkylX²C(O)R¹¹ (wherein X² represents -O- or -NR¹²- (in which R¹² represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹¹ represents C₁₋₃alkyl, -NR¹³R¹⁴ or -OR¹⁵

(wherein R^{13} , R^{14} and R^{15} which may be the same or different each represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));

3) C_{1-5} alkyl X^3R^{16} (wherein X^3 represents -O-, -S-, -SO-, -SO₂-, -OC(O)-, -NR¹⁷C(O)-, -C(O)NR¹⁸-, -SO₂NR¹⁹-, -NR²⁰SO₂- or -NR²¹- (wherein R^{17} , R^{18} , R^{19} , R^{20} and R^{21} each

independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{16} represents hydrogen, C_{1-3} alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C_{1-3} alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C_{1-4} alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);

4) C_{1-5} alkyl X^4C_{1-5} alkyl X^5R^{22} (wherein X^4 and X^5 which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR²³C(O)-, -C(O)NR²⁴-, -SO₂NR²⁵-, -NR²⁶SO₂- or -NR²⁷- (wherein R^{23} , R^{24} , R^{25} , R^{26} and R^{27} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{22} represents hydrogen or C_{1-3} alkyl);

5) R^{28} (wherein R^{28} is a 5-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-4} cyanoalkyl, C_{1-4} alkyl, C_{1-4} hydroxyalkyl, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl and C_{1-4} alkylsulphonyl C_{1-4} alkyl);

6) C_{1-5} alkyl R^{28} (wherein R^{28} is as defined herein);

7) C_{2-5} alkenyl R^{28} (wherein R^{28} is as defined herein);

8) C_{2-5} alkynyl R^{28} (wherein R^{28} is as defined herein);

9) R^{29} (wherein R^{29} represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents on an available carbon atom selected from hydroxy, halogeno, amino, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} hydroxyalkyl, C_{1-4} aminoalkyl, C_{1-4} alkylamino, C_{1-4} hydroxyalkoxy, carboxy, trifluoromethyl, cyano, -C(O)NR³⁰R³¹ and -NR³²C(O)R³³ (wherein R^{30} , R^{31} , R^{32} and R^{33} , which may be the same or different, each represents hydrogen, C_{1-4} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));

10) C_{1-5} alkyl R^{29} (wherein R^{29} is as defined herein);

11) C_{2-5} alkenyl R^{29} (wherein R^{29} is as defined herein);

12) C_{2-5} alkynyl R^{29} (wherein R^{29} is as defined herein);

- 13) $C_{1,3}\text{alkylX}^6\text{R}^{29}$ (wherein X^6 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{34}\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^{35}-$, $-\text{SO}_2\text{NR}^{36}-$, $-\text{NR}^{37}\text{SO}_2-$ or $-\text{NR}^{38}-$ (wherein R^{34} , R^{35} , R^{36} , R^{37} and R^{38} each independently represents hydrogen, $C_{1,3}\text{alkyl}$ or $C_{1,3}\text{alkoxyC}_{2,3}\text{alkyl}$) and R^{29} is as defined herein);
- 14) $C_{2,5}\text{alkenylX}^7\text{R}^{29}$ (wherein X^7 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{39}\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^{40}-$, $-\text{SO}_2\text{NR}^{41}-$, $-\text{NR}^{42}\text{SO}_2-$ or $-\text{NR}^{43}-$ (wherein R^{39} , R^{40} , R^{41} , R^{42} and R^{43} each independently represents hydrogen, $C_{1,3}\text{alkyl}$ or $C_{1,3}\text{alkoxyC}_{2,3}\text{alkyl}$) and R^{29} is as defined herein);
- 15) $C_{2,5}\text{alkynylX}^8\text{R}^{29}$ (wherein X^8 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{44}\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^{45}-$, $-\text{SO}_2\text{NR}^{46}-$, $-\text{NR}^{47}\text{SO}_2-$ or $-\text{NR}^{48}-$ (wherein R^{44} , R^{45} , R^{46} , R^{47} and R^{48} each independently represents hydrogen, $C_{1,3}\text{alkyl}$ or $C_{1,3}\text{alkoxyC}_{2,3}\text{alkyl}$) and R^{29} is as defined herein);
- 16) $C_{1,3}\text{alkylX}^9\text{C}_{1,3}\text{alkylR}^{29}$ (wherein X^9 represents $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}^{49}\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^{50}-$, $-\text{SO}_2\text{NR}^{51}-$, $-\text{NR}^{52}\text{SO}_2-$ or $-\text{NR}^{53}-$ (wherein R^{49} , R^{50} , R^{51} , R^{52} and R^{53} each independently represents hydrogen, $C_{1,3}\text{alkyl}$ or $C_{1,3}\text{alkoxyC}_{2,3}\text{alkyl}$) and R^{29} is as defined herein);
- 17) $C_{1,3}\text{alkylX}^9\text{C}_{1,3}\text{alkylR}^{28}$ (wherein X^9 and R^{28} are as defined herein);
- 18) $C_{2,5}\text{alkenyl}$ which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, $C_{1,4}\text{alkylamino}$, $\text{N,N-di}(C_{1,4}\text{alkyl})\text{amino}$, aminosulphonyl, $\text{N-C}_{1,4}\text{alkylaminosulphonyl}$ and $\text{N,N-di}(C_{1,4}\text{alkyl})\text{aminosulphonyl}$;
- 19) $C_{2,5}\text{alkynyl}$ which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro, amino, $C_{1,4}\text{alkylamino}$, $\text{N,N-di}(C_{1,4}\text{alkyl})\text{amino}$, aminosulphonyl, $\text{N-C}_{1,4}\text{alkylaminosulphonyl}$ and $\text{N,N-di}(C_{1,4}\text{alkyl})\text{aminosulphonyl}$;
- 20) $C_{2,5}\text{alkenylX}^9\text{C}_{1,4}\text{alkylR}^{28}$ (wherein X^9 and R^{28} are as defined herein); and
- 21) $C_{2,5}\text{alkynylX}^9\text{C}_{1,4}\text{alkylR}^{28}$ (wherein X^9 and R^{28} are as defined herein);
- and salts thereof, and prodrugs thereof for example esters, amides and sulphides, in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals such as humans.

3. The use of a compound of the formula I according to claim 1, wherein R^2 represents hydroxy, halogeno, cyano, nitro, trifluoromethyl, $C_{1,3}\text{alkyl}$, amino or R^5X^1 - [wherein X^1 is as defined in claim 1 and R^5 is selected from one of the following twenty-two groups:

- 1) C₁₋₄alkyl which may be unsubstituted or which may be substituted with one or more groups selected from fluoro, chloro and bromo, or C₂₋₅alkyl which may be unsubstituted or substituted with one or more groups selected from hydroxy and amino;
- 2) C₂₋₃alkylX²C(O)R¹¹ (wherein X² is as defined in claim 1 and R¹¹ represents -NR¹³R¹⁴ or -OR¹⁵ (wherein R¹³, R¹⁴ and R¹⁵ which may be the same or different are each C₁₋₄alkyl or C₁₋₃alkoxyethyl));
- 3) C₂₋₄alkylX³R¹⁶ (wherein X³ is as defined in claim 1 and R¹⁶ is a group selected from C₁₋₃alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny and tetrahydropyranyl, which C₁₋₃alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C₁₋₂alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny or tetrahydropyranyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group -(O-)_f(C₁₋₃alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl));
- 4) C₂₋₃alkylX⁴C₂₋₃alkylX⁵R²² (wherein X⁴ and X⁵ are as defined in claim 1 and R²² represents hydrogen or C₁₋₃alkyl);
- 5) R²⁸ (wherein R²⁸ is as defined in claim 1);
- 6) C₁₋₄alkylR¹¹⁰ (wherein R¹¹⁰ is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidin-1-yl, azetidiny, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to C₁₋₄alkyl through a carbon atom and which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group -(O-)_f(C₁₋₃alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋

- ₃alkyl)) or C₂₋₄alkylR¹¹¹ (wherein R¹¹¹ is a group selected from morpholino, thiomorpholino, azetidin-1-yl, pyrrolidin-1-yl, piperazin-1-yl and piperidino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group -(O-)_f(C₁₋₃alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl));
- 7) C₃₋₄alkenylR¹¹² (wherein R¹¹² represents R¹¹⁰ or R¹¹¹ as defined herein);
- 8) C₃₋₄alkynylR¹¹² (wherein R¹¹² represents R¹¹⁰ or R¹¹¹ as defined herein);
- 9) R²⁹ (wherein R²⁹ is as defined in claim 1);
- 10) C₁₋₄alkylR²⁹ (wherein R²⁹ is as defined in claim 1);
- 11) 1-R²⁹prop-1-en-3-yl or 1-R²⁹but-2-en-4-yl (wherein R²⁹ is as defined in claim 1 with the proviso that when R⁵ is 1-R²⁹prop-1-en-3-yl, R²⁹ is linked to the alkenyl group via a carbon atom);
- 12) 1-R²⁹prop-1-yn-3-yl or 1-R²⁹but-2-yn-4-yl (wherein R²⁹ is as defined in claim 1 with the proviso that when R⁵ is 1-R²⁹prop-1-yn-3-yl, R²⁹ is linked to the alkynyl group via a carbon atom);
- 13) C₁₋₅alkylX⁶R²⁹ (wherein X⁶ and R²⁹ are as defined in claim 1);
- 14) 1-(R²⁹X⁷)but-2-en-4-yl (wherein X⁷ and R²⁹ are as defined in claim 1);
- 15) 1-(R²⁹X⁸)but-2-yn-4-yl (wherein X⁸ and R²⁹ are as defined in claim 1);
- 16) C₂₋₃alkylX⁹C₁₋₃alkylR²⁹ (wherein X⁹ and R²⁹ are as defined in claim 1);
- 17) C₂₋₃alkylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined in claim 1);
- 18) C₂₋₅alkenyl which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;
- 19) C₂₋₅alkynyl which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, fluoro, amino, C₁₋

₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;

20) C₂₋₄alkenylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined in claim 1);

21) C₂₋₄alkynylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined in claim 1); and

5 22) C₁₋₃alkylR⁵⁴(C₁₋₃alkyl)_q(X⁹)_rR⁵⁵ (wherein X⁹, q, r, R⁵⁴ and R⁵⁵ are as defined in claim 1);
and additionally wherein any C₁₋₅alkyl, C₂₋₅alkenyl or C₂₋₅alkynyl group in R⁵X¹- may bear one or more substituents selected from hydroxy, halogeno and amino].

10 4. The use of a compound of the formula I according to any one of the preceding claims wherein Z is -O-, -NH- or -S-.


15 5. The use of a compound of the formula I according to any one of the preceding claims wherein ring C is a 9-10-membered heteroaromatic bicyclic moiety which contains 1-3 heteroatoms selected independently from O, N and S.

20 6. The use of a compound of the formula I according to any one of the preceding claims wherein R¹ represents oxo, halogeno, hydroxy, C₁₋₂alkoxy, C₁₋₂alkyl, C₁₋₂alkoxymethyl, C₂₋₃alkanoyl, C₁₋₂haloalkyl, cyano, amino, C₂₋₄alkenyl, C₂₋₄alkynyl, C₂₋₃alkanoyloxy, nitro, C₂₋₃alkanoylamino, C₁₋₂alkoxycarbonyl, C₁₋₂alkylsulphanyl, C₁₋₂alkylsulphinyl, C₁₋₂alkylsulphonyl, carbamoyl, N-C₁₋₂alkylcarbamoyl, N,N-di(C₁₋₂alkyl)carbamoyl, aminosulphonyl, N-C₁₋₂alkylaminosulphonyl, N,N-di(C₁₋₂alkyl)aminosulphonyl, N-(C₁₋₂alkylsulphonyl)amino, N-(C₁₋₂alkylsulphonyl)-N-(C₁₋₂alkyl)amino or a C₃₋₇alkylene chain joined to two ring C carbon atoms.

25 7. The use of a compound of the formula I according to any one of the preceding claims wherein n is 0, 1 or 2.

30 8. The use of a compound of the formula I according to any one of the preceding claims wherein m is 1 or 2.

9. A compound of the formula II:



(II)

- 10 [wherein:
 ring C, R^1 , R^2 and n are as defined in claim 1, Zb is $-O-$ or $-S-$ and R^{2a} represents hydrogen,
 halogeno, C_{1-3} alkyl, trifluoromethyl, C_{1-3} alkoxy, C_{1-3} alkylsulphanyl, $-NR^{3a}R^{4a}$ (wherein R^{3a} and
 R^{4a} , which may be the same or different, each represents hydrogen or C_{1-3} alkyl), or
 $R^{5a}(CH_2)_{za}X^{1a}$ (wherein R^{5a} is a 5- or 6-membered saturated heterocyclic group with 1-2
 15 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or
 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-4} cyanoalkyl, C_{1-4} alkyl, C_{1-4}
 hydroxyalkyl, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, C_{1-4} alkylsulphonyl C_{1-4} alkyl, C_{1-4}
 alkoxy carbonyl, C_{1-4} aminoalkyl, C_{1-4} alkylamino, di(C_{1-4} alkyl)amino, C_{1-4} alkylamino C_{1-4} alkyl,
 di(C_{1-4} alkyl)amino C_{1-4} alkyl, C_{1-4} alkylamino C_{1-4} alkoxy, di(C_{1-4} alkyl)amino C_{1-4} alkoxy and a
 20 group $-(O-)_f(C_{1-4}alkyl)_g$ ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 5-6-membered
 saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N,
 which cyclic group may bear one or more substituents selected from C_{1-4} alkyl), za is an
 integer from 0 to 4 and X^{1a} represents a direct bond, $-O-$, $-CH_2-$, $-S-$, $-SO-$, $-SO_2-$, $-NR^{6a}C(O)-$,
 $-C(O)NR^{7a}-$, $-SO_2NR^{8a}-$, $-NR^{9a}SO_2-$ or $-NR^{10a}-$ (wherein R^{6a} , R^{7a} , R^{8a} , R^{9a} and R^{10a} each
 25 independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));
 with the proviso that R^2 is not hydrogen and excluding the compounds:
 6,7-dimethoxy-4-(1-naphthylsulphanyl)quinazoline, 6,7-dimethoxy-4-(2-
 naphthylsulphanyl)quinazoline, 6,7-dimethoxy-4-(1-naphthyloxy)quinazoline and 6,7-
 dimethoxy-4-(2-naphthyloxy)quinazoline;
 30 or a salt thereof.

10. A compound of the formula II according to claim 9 wherein R² represents hydroxy, halogeno, cyano, nitro, trifluoromethyl, C₁₋₃alkyl, amino or R⁵X¹- [wherein X¹ is as defined in claim 1 and R⁵ is selected from one of the following twenty-two groups:

- 1) C₁₋₄alkyl which may be unsubstituted or which may be substituted with one or more groups
5 selected from fluoro, chloro and bromo, or C₂₋₃alkyl which may be unsubstituted or substituted with one or more groups selected from hydroxy and amino;
- 2) C₂₋₃alkylX²C(O)R¹¹ (wherein X² is as defined in claim 1 and R¹¹ represents -NR¹³R¹⁴ or -OR¹⁵ (wherein R¹³, R¹⁴ and R¹⁵ which may be the same or different are each C₁₋₄alkyl or C₁₋₂alkoxyethyl));
- 10 3) C₂₋₄alkylX³R¹⁶ (wherein X³ is as defined in claim 1 and R¹⁶ is a group selected from C₁₋₃alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny and tetrahydropyranyl, which C₁₋₃alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C₁₋₂alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny or tetrahydropyranyl group may bear 1 or 2
15 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group -(-O-)_f(C₁₋₃alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group
20 selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl));
- 4) C₂₋₃alkylX⁴C₂₋₃alkylX⁵R²² (wherein X⁴ and X⁵ are as defined in claim 1 and R²² represents hydrogen or C₁₋₃alkyl);
- 25 5) R²⁸ (wherein R²⁸ is as defined in claim 1);
- 6) C₁₋₄alkylR¹¹⁰ (wherein R¹¹⁰ is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidin-1-yl, azetidiny, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to C₁₋₄alkyl through a carbon atom and which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group -(-
- 30

- O-)_f(C₁₋₃alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl)) or C₂₋₄alkylR¹¹¹ (wherein R¹¹¹ is a group selected from morpholino, thiomorpholino, azetidin-1-yl, pyrrolidin-1-yl, piperazin-1-yl and piperidino which group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C₁₋₃cyanoalkyl, C₁₋₃alkyl, C₁₋₃hydroxyalkyl, C₁₋₃alkoxy, C₁₋₂alkoxyC₁₋₃alkyl, C₁₋₂alkylsulphonylC₁₋₃alkyl, C₁₋₃alkoxycarbonyl, C₁₋₃alkylamino, di(C₁₋₃alkyl)amino, C₁₋₃alkylaminoC₁₋₃alkyl, di(C₁₋₃alkyl)aminoC₁₋₃alkyl, C₁₋₃alkylaminoC₁₋₃alkoxy, di(C₁₋₃alkyl)aminoC₁₋₃alkoxy and a group -(-
- 10 O-)_f(C₁₋₃alkyl)_gringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a heterocyclic group selected from pyrrolidinyl, piperazinyl, piperidinyl, imidazolidinyl, azetidiny, morpholino and thiomorpholino, which cyclic group may bear one or more substituents selected from C₁₋₃alkyl));
- 7) C₃₋₄alkenylR¹¹² (wherein R¹¹² represents R¹¹⁰ or R¹¹¹ as defined herein);
- 15 8) C₃₋₄alkynylR¹¹² (wherein R¹¹² represents R¹¹⁰ or R¹¹¹ as defined herein);
- 9) R²⁹ (wherein R²⁹ is as defined in claim 1);
- 10) C₁₋₄alkylR²⁹ (wherein R²⁹ is as defined in claim 1);
- 11) 1-R²⁹prop-1-en-3-yl or 1-R²⁹but-2-en-4-yl (wherein R²⁹ is as defined in claim 1 with the proviso that when R⁵ is 1-R²⁹prop-1-en-3-yl, R²⁹ is linked to the alkenyl group via a carbon
- 20 atom);
- 12) 1-R²⁹prop-1-yn-3-yl or 1-R²⁹but-2-yn-4-yl (wherein R²⁹ is as defined in claim 1 with the proviso that when R⁵ is 1-R²⁹prop-1-yn-3-yl, R²⁹ is linked to the alkynyl group via a carbon atom);
- 13) C₁₋₃alkylX⁶R²⁹ (wherein X⁶ and R²⁹ are as defined in claim 1);
- 25 14) 1-(R²⁹X⁷)but-2-en-4-yl (wherein X⁷ and R²⁹ are as defined in claim 1);
- 15) 1-(R²⁹X⁸)but-2-yn-4-yl (wherein X⁸ and R²⁹ are as defined in claim 1);
- 16) C₂₋₃alkylX⁹C₁₋₃alkylR²⁹ (wherein X⁹ and R²⁹ are as defined in claim 1);
- 17) C₂₋₃alkylX⁹C₁₋₃alkylR²⁸ (wherein X⁹ and R²⁸ are as defined in claim 1);
- 18) C₂₋₃alkenyl which may be unsubstituted or which may be substituted with one or more
- 30 fluorine atoms or with one or two groups selected from hydroxy, fluoro, amino, C₁₋₄alkylamino, N,N-di(C₁₋₄alkyl)amino, aminosulphonyl, N-C₁₋₄alkylaminosulphonyl and N,N-di(C₁₋₄alkyl)aminosulphonyl;

19) $C_{2,5}$ alkynyl which may be unsubstituted or which may be substituted with one or more fluorine atoms or with one or two groups selected from hydroxy, fluoro, amino, $C_{1,4}$ alkylamino, N,N -di($C_{1,4}$ alkyl)amino, aminosulphonyl, N - $C_{1,4}$ alkylaminosulphonyl and N,N -di($C_{1,4}$ alkyl)aminosulphonyl;

5 20) $C_{2,4}$ alkenyl $X^9C_{1,3}$ alkyl R^{28} (wherein X^9 and R^{28} are as defined in claim 1);

21) $C_{2,4}$ alkenyl $X^9C_{1,3}$ alkyl R^{28} (wherein X^9 and R^{28} are as defined in claim 1); and

22) $C_{1,3}$ alkyl $R^{54}(C_{1,3}$ alkyl) $_q(X^9)_rR^{55}$ (wherein X^9 , q , r , R^{54} and R^{55} are as defined in claim 1);

and additionally wherein any $C_{1,5}$ alkyl, $C_{2,5}$ alkenyl or $C_{2,5}$ alkynyl group in R^5X^1 - may bear one or more substituents selected from hydroxy, halogeno and amino].

10

11. A compound according to any one of claims 9 and 10 wherein Zb is -O-.

12. A compound according to any one of claims 9, 10 and 11 wherein ring C is a 9-10-membered heteroaromatic bicyclic moiety which contains 1-3 heteroatoms selected

15 independently from O, N and S.


13. A compound according to any one of claims 9, 10, 11 and 12 wherein R^1 represents oxo, halogeno, hydroxy, $C_{1,2}$ alkoxy, $C_{1,2}$ alkyl, $C_{1,2}$ alkoxymethyl, $C_{2,3}$ alkanoyl, $C_{1,2}$ haloalkyl, cyano, amino, $C_{2,4}$ alkenyl, $C_{2,4}$ alkynyl, $C_{2,3}$ alkanoyloxy, nitro, $C_{2,3}$ alkanoylamino, $C_{1,2}$ alkoxycarbonyl, $C_{1,2}$ alkylsulphanyl, $C_{1,2}$ alkylsulphinyl, $C_{1,2}$ alkylsulphonyl, carbamoyl, N - $C_{1,2}$ alkylcarbamoyl, N,N -di($C_{1,2}$ alkyl)carbamoyl, aminosulphonyl, N - $C_{1,2}$ alkylaminosulphonyl, N,N -di($C_{1,2}$ alkyl)aminosulphonyl, N -($C_{1,2}$ alkylsulphonyl)amino, N -($C_{1,2}$ alkylsulphonyl)- N -($C_{1,2}$ alkyl)amino or a $C_{3,7}$ alkylene chain joined to two ring C carbon atoms.

25

14. A compound according to any one of claims 9, 10, 11, 12 and 13 wherein n is 0, 1 or 2.

15. A compound of the formula IIb:

30



(IIb)

15 [wherein:

ring C, R^1 , R^2 and n are as defined in claim 1, Zb is -O- and R^{2a} is as defined in claim 9 with the proviso that R^2 does not have any of the following values:

hydrogen, substituted or unsubstituted C_{1-5} alkyl, halogeno, C_{1-5} alkoxy, C_{2-5} alkenyl, phenoxy or phenyl C_{1-5} alkoxy;

20 or a salt thereof.

16. A compound according to claim 9 selected from

6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(2-naphthyloxy)quinazoline,

6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,

25 7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(quinolin-7-yloxy)quinazoline,

6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)-4-(quinolin-7-yloxy)quinazoline,

6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,

4-(4-chloroquinolin-7-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,

6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)-4-(4-methylquinolin-7-yloxy)quinazoline,

30 6-methoxy-4-(4-methylquinolin-7-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,

6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(quinolin-7-yloxy)quinazoline,

- 6-methoxy-7-((1-(2-methylsulphonyl)ethyl)piperidin-4-yl)methoxy)-4-(quinolin-7-yloxy)quinazoline,
4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline,
4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,
5 6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline,
6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)-4-(2-trifluoromethylindol-5-yloxy)quinazoline,
(*R,S*)-4-(3-fluoroquinolin-7-yloxy)-6-methoxy-7-((1-methylpiperidin-3-yl)methoxy)quinazoline,
10 4-(indol-5-yloxy)-6-methoxy-7-(3-methylsulphonylpropoxy)quinazoline,
7-(3-N,N-dimethylaminopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(2-morpholinoethoxy)ethoxy)quinazoline,
7-(2-(N,N-diethylamino)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-7-(3-piperidinopropoxy)-4-(quinolin-7-yloxy)quinazoline,
15 4-(2-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(2-(piperidin-1-yl)ethoxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(2-(1*H*-1,2,4-triazol-1-yl)ethoxy)quinazoline,
6-methoxy-7-(3-piperidinopropoxy)-4-(6-trifluoromethylindol-5-yloxy)quinazoline,
7-(3-(methylsulphonyl)propoxy)-4-(2-methylindol-5-yloxy)quinazoline,
20 7-(3-(N,N-dimethylamino)propoxy)-4-(2,3-dimethylindol-5-yloxy)-6-methoxyquinazoline,
4-(2,3-dimethylindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-3-ylmethoxy)quinazoline,
7-(2-(N,N-diethylamino)ethoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(2-(piperidin-2-yl)ethoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(2-(piperidin-1-yl)ethoxy)quinazoline,
25 4-(indol-6-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
7-(3-(ethylsulphonyl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-4-(3-methylindol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline,
7-(2-hydroxy-3-piperidinopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
7-(2-hydroxy-3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
30 yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methylamino)ethoxy)quinazoline, and
7-(2-hydroxy-3-(isopropylamino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,

or a salt thereof.

17. A compound according to claim 9 selected from

- 6-methoxy-7-(3-morpholinopropoxy)-4-(quinolin-7-yloxy)quinazoline,
5 6-methoxy-4-(2-methylindol-5-yloxy)-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-methylsulphonylpropoxy)quinazoline,
7-((1-cyanomethyl)piperidin-4-ylmethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
10 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-morpholinoethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-pyrrolidin-1-ylethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-methylpiperidin-3-ylmethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-piperidinoethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-(4-
15 pyridyl)amino)ethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-morpholinopropoxy)quinazoline,
6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(1*H*-1,2,4-triazol-1-yl)ethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(2-(4-methylpiperazin-1-
20 yl)ethoxy)ethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-piperidinopropoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline
6-methoxy-7-(1-(2-methoxyethyl)piperidin-4-ylmethoxy)-4-(2-methylindol-5-
yloxy)quinazoline,
25 6-methoxy-4-(2-methylindol-5-yloxy)-7-((2-(2-pyrrolidin-1-
ylethyl)carbamoyl)vinyl)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(piperidin-4-yloxy)ethoxy)quinazoline,
30 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(N-methyl-N-
methylsulphonylamino)ethoxy)quinazoline,

- 7-(2-(1-(2-cyanoethyl)piperidin-4-yloxy)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(3-(pyrrolidin-yl)propoxy)quinazoline,
4-(2-methylindol-5-yloxy)-7-(3-(1,1-dioxothiomorpholino)propoxy)quinazoline,
5 4-(2-methylindol-5-yloxy)-7-(piperidin-4-ylmethoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(2-(2-methoxyethoxy)ethoxy)quinazoline,
7-(3-(N,N-dimethylamino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
7-(3-(N,N-diethylamino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
7-(3-(1,1-dioxothiomorpholino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
10 4-(indol-5-yloxy)-6-methoxy-7-(2-(4-pyridyloxy)ethoxy)quinazoline,
4-(indol-6-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
7-(1-(2-methoxyethyl)piperidin-4-ylmethoxy)-4-(2-methylindol-5-yloxy)quinazoline,
7-(2-hydroxy-3-morpholinopropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
7-(2-(1-(2-methoxyethyl)piperidin-4-yl)ethoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
15 7-(2-hydroxy-3-pyrrolidin-1-ylpropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
7-(3-(N,N-diethylamino)-2-hydroxypropoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
7-(3-(1,1-dioxothiomorpholino)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
20 6-methoxy-4-(2-methylindol-5-yloxy)-7-(2-(4-pyridyloxy)ethoxy)quinazoline,
4-(indol-5-yloxy)-6-methoxy-7-(3-morpholinopropoxy)quinazoline,
(2*R*)-6-methoxy-(2-methyl-1*H*-indol-5-yloxy)-7-(2-hydroxy-3-piperidinopropoxy)quinazoline,
(5*R*)-6-methoxy-4-(2-methyl-1*H*-indol-5-yloxy)-7-(2-oxopyrrolidin-5-ylmethoxy)quinazoline,
25 4-(4-bromoindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-(2-(pyrrolidin-1-yl)ethyl)-piperidin-4-ylmethoxy)quinazoline,
(2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
(2*R*)-7-(2-hydroxy-3-morpholinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
30 (2*R*)-7-(2-hydroxy-3-piperidinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
(2*S*)-7-(2-hydroxy-3-(N,N-diisopropyl)amino)propoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,

- (2*S*)-7-(2-hydroxy-3-piperidinopropoxy)-4-(indol-5-yloxy)-6-methoxyquinazoline,
(2*R*)-7-(2-hydroxy-3-piperidinopropoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline,
(2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-4-(3-methylindol-5-yloxy)quinazoline,
5 (2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
(2*R*)-7-(2-hydroxy-3-(4-methylpiperazin-1-yl)propoxy)-6-methoxy-4-(2-methylindol-5-yloxy)quinazoline,
6-methoxy-4-(2-methylindol-5-yloxy)-7-(1-(2-morpholinoethyl)piperidin-4-ylmethoxy)quinazoline,
10 4-(3-fluoro-quinolin-7-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
4-(3-fluoro-quinolin-7-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)-4-(1*H*-pyrrolo[2,3-*b*]pyridin-5-yloxy)quinazoline,
(2*S*)-6-methoxy-(2-methyl-1*H*-indol-5-yloxy)-7-(2-hydroxy-3-piperidinopropoxy)quinazoline,
15 and
4-(6-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
or a salt thereof.

18. A compound according to claim 9 selected from

- 20 6-methoxy-4-(2-methylindol-5-yloxy)-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline,
4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline,
4-(6-fluoroindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
25 4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(pyrrolidin-1-yl)propoxy)quinazoline,
4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-piperidinopropoxy)quinazoline,
4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-((1-methylpiperidin-4-yl)methoxy)quinazoline,
30 4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-(4-methylpiperazin-1-yl)propoxy)quinazoline,
4-(4-fluoroindol-5-yloxy)-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline,

(2*R*)-7-(2-hydroxy-3-(pyrrolidin-1-yl)propoxy)-4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxyquinazoline, and


4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(2-(1-methylpiperidin-4-yl)ethoxy)quinazoline,

5 or a salt thereof.

19. A compound according to claim 9 in the form of a pharmaceutically acceptable salt.

10 20. A process for the preparation of a compound of formula I or salt thereof which comprises:


(a) the reaction of a compound of the formula III:



(III)

20

(wherein R^2 and m are as defined in claim 1 and L^1 is a displaceable moiety), with a compound of the formula IV:



25


(IV)

(wherein ring C, R^1 , Z and n are as defined in claim 1);

(b) a compound of formula I or a salt thereof wherein at least one R^2 is R^5X^1 wherein R^5 is as defined in claim 1 and X^1 is -O-, -S-, -OC(O)- or -NR¹⁰- (wherein R^{10} independently

- 338 -

represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) may be prepared by the reaction of a compound of the formula V:



(V)


(wherein ring C, Z, R¹, R² and n are as defined in claim 1 and X¹ is as herein defined in this section and s is an integer from 0 to 2) with a compound of formula VI:



(VI)


(wherein R⁵ is as defined in claim 1 and L¹ is as herein defined);

(c) a compound of the formula I or a salt thereof wherein at least one R² is R⁵X¹ wherein R⁵ is as defined in claim 1 and X¹ is -O-, -S-, -OC(O)- or -NR¹⁰- (wherein R¹⁰ represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) may be prepared by the reaction of a compound of the formula VII:



(VII)


with a compound of the formula VIII:



(VIII)


(wherein R^1 , R^2 , R^5 , ring C, Z and n are as defined in claim 1 and L^1 , s and X^1 are as herein defined);

- 5 (d) a compound of the formula I or a salt thereof wherein at least one R^2 is R^5X^1 wherein X^1 is as defined in claim 1 and R^5 is $C_{1-5}alkylR^{113}$, wherein R^{113} is selected from one of the following nine groups:
- 1) $X^{19}C_{1-3}alkyl$ (wherein X^{19} represents -O-, -S-, -SO₂-, -NR¹¹⁴C(O)- or -NR¹¹⁵SO₂- (wherein R^{114} and R^{115} which may be the same or different are each hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$);
 - 10 2) $NR^{116}R^{117}$ (wherein R^{116} and R^{117} which may be the same or different are each hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$);
 - 3) $X^{20}C_{1-5}alkylX^5R^{22}$ (wherein X^{20} represents -O-, -S-, -SO₂-, -NR¹¹⁸C(O)-, -NR¹¹⁹SO₂- or -NR¹²⁰- (wherein R^{118} , R^{119} , and R^{120} which may be the same or different are each hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$) and X^5 and R^{22} are as defined in claim 1);
 - 15 4) R^{28} (wherein R^{28} is as defined in claim 1);
 - 5) $X^{21}R^{29}$ (wherein X^{21} represents -O-, -S-, -SO₂-, -NR¹²¹C(O)-, -NR¹²²SO₂-, or -NR¹²³- (wherein R^{121} , R^{122} , and R^{123} which may be the same or different are each hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$) and R^{29} is as defined in claim 1); and
 - 20 6) $X^{22}C_{1-3}alkylR^{29}$ (wherein X^{22} represents -O-, -S-, -SO₂-, -NR¹²⁴C(O)-, -NR¹²⁵SO₂- or -NR¹²⁶- (wherein R^{124} , R^{125} and R^{126} each independently represents hydrogen, $C_{1-3}alkyl$ or $C_{1-3}alkoxyC_{2-3}alkyl$) and R^{29} is as defined in claim 1);
 - 7) R^{29} (wherein R^{29} is as defined in claim 1);
 - 8) $X^{22}C_{1-4}alkylR^{28}$ (wherein X^{22} and R^{28} are as defined in claim 1); and
 - 25 9) $R^{54}(C_{1-4}alkyl)_q(X^9)_rR^{55}$ (wherein q, r, X^9 , R^{54} and R^{55} are as defined in claim 1);
- may be prepared by reacting a compound of the formula IX:



(IX)

- 10 (wherein X^1 , R^1 , R^2 , ring C, Z and n are as defined in claim 1 and L^1 and s are as herein defined) with a compound of the formula X:



(X)

(wherein R^{113} is as defined herein);

- 15 (e) a compound of the formula I or a salt thereof wherein one or more of the substituents $(R^2)_m$ is represented by $-NR^{127}R^{128}$, where one (and the other is hydrogen) or both of R^{127} and R^{128} are C_{1-3} alkyl, may be effected by the reaction of compounds of formula I wherein the substituent $(R^2)_m$ is an amino group and an alkylating agent; or
- (f) a compound of the formula I or a salt thereof wherein X^1 is $-SO-$ or $-SO_2-$ may be
- 20 prepared by oxidation from the corresponding compound in which X^1 is $-S-$ or $-SO-$; and when a salt of a compound of formula I is required, reaction of the compound obtained with an acid or base whereby to obtain the desired salt.

21. A pharmaceutical composition which comprises as active ingredient a compound
- 25 of formula I or a pharmaceutically acceptable salt thereof according to claim 9 in association with a pharmaceutically acceptable excipient or carrier.

22. A method for producing an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal in need of such treatment which comprises administering to
- 30 said animal an effective amount of a compound of formula I as defined in claim 1 or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB99/00373

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/505 C07D401/14 C07D413/14 C07D417/12 C07D405/12
C07D401/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 29301 A (AGREVO UK LTD ; CORNELL CLIVE LEONARD (GB); RICHARDS IAN CHRISTOPHE) 26 September 1996 (1996-09-26) see compounds 67 and 83 ---	9-11, 14
X	WO 95 15758 A (HSU CHIN YI JENNY ; ZILBERSTEIN ASHER (US); JOHNSON SUSAN E (US); M) 15 June 1995 (1995-06-15) see compound on top of page 17 the whole document ---	9-11, 14, 19-21
Y	---	1-22
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

7 April 2000

Date of mailing of the international search report

18.05.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Scruton-Evans, I

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 00/00373

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	HENNEQUIN L F ET AL: "Design and Structure-Activity Relationship of a New Class of Potent VEGF Receptor Tyrosine Kinase Inhibitors" J. MED. CHEM. (JMCMAR,00222623);1999; VOL.42 (26); PP.5369-5389, XP000197180 AstraZeneca Zeneca Pharma Centre de Recherches Z.I. La Pompelle;Reims; 51689; Fr. (FR) the whole document	1-22
Y	BRIDGES A J ET AL: "Enantioselective inhibition of the epidermal growth factor receptor tyrosine kinase by 4-(.alpha.-phenethylamino)quinazolines" BIOORG. MED. CHEM. (BMECEP,09680896);1995; VOL.3 (12); PP.1651-6, XP002134974 Parke-Davis Pharmaceutical Research Division;Ann Arbor; 48105; MI; USA (US) the whole document	1-22
Y	GIBSON K H ET AL: "Epidermal growth factor receptor tyrosine kinase: structure-activity relationships and antitumor activity of novel quinazolines" BIOORG. MED. CHEM. LETT. (BMCLE8,0960894X);1997; VOL.7 (21); PP.2723-2728, XP002134975 Zeneca Pharmaceuticals;Research Dep. Cancer, Metabolism and Endocrine; Alderley Park, Macclesfield, Cheshire; SK10 4TG; UK (GB) the whole document	1-22
Y	EP 0 602 851 A (ZENECA LTD) 22 June 1994 (1994-06-22) the whole document	1-22
X	WO 97 42187 A (LOHMANN JEAN JACQUES MARCEL ;PLE PATRICK (FR); HENNEQUIN LAURENT F) 13 November 1997 (1997-11-13) see the general formula I and pages 1-3	1-3,6-8, 20
P,X	WO 99 10349 A (LOHMANN JEAN JACQUES MARCEL ;PLE PATRICK (FR); HENNEQUIN LAURENT F) 4 March 1999 (1999-03-04) the whole document	1-3,6-8, 20
Y	WO 97 22596 A (ZENECA LTD ;ZENECA PHARMA SA (FR); LOHMANN JEAN JACQUES MARCEL (FR) 26 June 1997 (1997-06-26) the whole document	1-22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 00/00373

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 22 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/00373

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9629301 A	26-09-1996	AU 5116096 A EP 0819111 A JP 11503114 T ZA 9602319 A	08-10-1996 21-01-1998 23-03-1999 30-10-1996
WO 9515758 A	15-06-1995	US 5480883 A US 5710158 A AU 1305095 A EP 0871448 A SG 54172 A US 5795889 A US 5646153 A US 5721237 A US 5714493 A US RE36256 E	02-01-1996 20-01-1998 27-06-1995 21-10-1998 16-11-1998 18-08-1998 08-07-1997 24-02-1998 03-02-1998 20-07-1999
EP 0602851 A	22-06-1994	AT 143956 T AU 664496 B AU 5072893 A CA 2103383 A CN 1094043 A CZ 9302651 A DE 69305310 D DE 69305310 T DK 602851 T ES 2093367 T FI 935431 A GR 3021326 T HU 65622 A IL 107678 A JP 6336481 A MX 9307778 A NO 934504 A NZ 250218 A US 5580870 A ZA 9308594 A	15-10-1996 16-11-1995 23-06-1994 11-06-1994 26-10-1994 15-06-1994 14-11-1996 13-02-1997 24-03-1997 16-12-1996 11-06-1994 31-01-1997 28-07-1994 12-03-1999 06-12-1994 29-07-1994 13-06-1994 25-06-1996 03-12-1996 10-06-1994
WO 9742187 A	13-11-1997	AU 2647597 A EP 0912557 A	26-11-1997 06-05-1999
WO 9910349 A	04-03-1999	AU 8816298 A	16-03-1999
WO 9722596 A	26-06-1997	AU 712370 B AU 1106197 A CZ 9801882 A EP 0873319 A HU 9901243 A NO 982784 A PL 327310 A SK 82898 A US 5962458 A	04-11-1999 14-07-1997 16-09-1998 28-10-1998 30-08-1999 17-08-1998 07-12-1998 04-11-1998 05-10-1999